

Reproductive biology of *Tephroseris longifolia* subsp. *moravica*, an endemic taxon of European importance

Monika Janišová*, Iveta Škodová and Katarína Hegedúšová

Institute of Botany, Slovak Academy of Sciences, Dúbravská cesta 9, 845 23 Bratislava, Slovakia

(Received 22 December 2010; accepted after revision 14 November 2011; first published online 24 January 2012)

Abstract

Tephroseris longifolia subsp. *moravica* (Asteraceae) is an endangered narrow endemic taxon recently recorded from nine sites in Slovakia and the Czech Republic. We provide the first consistent information on its reproductive biology – mating system, seed output, germination, seedling emergence and survival. Based on results from isolation, hybridization and emasculatation experiments on different subspecies, *T. longifolia* s.l. should be treated as an allogamous taxon without apomictic reproduction. Five populations of *T. l. moravica* were studied in detail for seed output and germination characteristics. The output of well-developed seeds per flowering shoot was estimated to be 592 (1125 seeds per flowering plant). A granivorous butterfly, *Phycitodes albatella* (family Pyralidae), was identified as a pest, reducing seed output by 18–28%. Germination (reaching 30–62% after dry storage for 4 months) rapidly decreased with storage in room conditions. Deep freeze storage increased germination, although not significantly. Seeds obtained from natural populations had higher germination percentages and germinated faster than seeds from cultivated plants. Seedling emergence *in situ* was very low (0.9–2.06% of the sown seeds) without significant differences among sites.

Keywords: mating system, *Phycitodes albatella*, seed germination, seed output, *Tephroseris longifolia*

Introduction

Knowledge of the reproductive biology of a species is important for its successful conservation. In many cases, an indication of the factors responsible for

the rarity of a species may help to set appropriate conservation measures. Demographic and reproductive traits such as low colonization ability (Byers and Meagher, 1997), reduced seed output or hampered seedling establishment due to herbivory (Münzbergová, 2005; Scheidel and Bruelheide, 2005) often differentiate rare plants from their widespread congeners (Kunin and Gaston, 1993; Farnsworth, 2007). Investigation of mating system and seed output may thus provide valuable indications of recent status and perspectives of species populations (Purdy *et al.*, 1994; Isaksson, 2009).

Tephroseris is a genus of approximately 50 species primarily distributed in northern Eurasia (Jeffrey and Chen, 1984; Nordenstam, 2007). From the chromosome numbers previously reported for many species of this genus, it is most likely that $x = 24$ is at least a common, or even the exclusive, basic chromosome number for the genus (Liu and Yang, 2011). In the family Asteraceae and tribe Senecioneae, a wide variety of mating systems and reproductive strategies has been recorded (Czapik, 1996; Nordenstam, 2007; Noyes, 2007; Cron *et al.*, 2009). The self-incompatibility systems studied within the family Asteraceae, documented also in *Tephroseris* and the closely related genus *Senecio* (Hiscock and Tabah, 2003; Isaksson, 2009), were found to be sporophytic, which is a multi-allelic one-locus system, often with complicated hierarchical dominance relationships between alleles.

Our study focuses on the reproductive biology of a narrow endemic subspecies, *Tephroseris longifolia* subsp. *moravica* Holub (Asteraceae). This taxon belongs to the group of *Tephroseris longifolia* (Jacq.) Griseb. *et* Schenk s.l., which has the centre of its distribution area in the eastern Alps (Austria, Slovenia, northern Italy) but reaches also France, Switzerland, Germany, Czech Republic, Slovakia, Hungary, Serbia, Croatia, Bosnia and Albania (Greuter, 2006–2009). Five subspecies can be distinguished within *T. longifolia* s.l. (Greuter, 2006–2009), differing in both their distribution areas and habitat characteristics (Wagenitz, 1987; Aeschmann *et al.*, 2004). *T. l. moravica* is considered

*Correspondence
Fax: + 421 2 54771948
Email: monika.janisova@savba.sk

to be endemic to the western Carpathians (Kliment, 1999). Recently, nine populations of *T. l. moravica* were recorded in western Slovakia and eastern Czech Republic, and seven populations are considered to be extinct (Fig. 1). The taxon is included in the European list of important species (Council Directive 92/43/EHS on the conservation of natural habitats and of wild fauna and flora, Annex II: Animal and plant species of community interest whose conservation requires the designation of special areas of conservation; Decree of Ministry of Environment of SR No. 24/2003 implementing the Act No. 543/2002 Coll. on Nature and Landscape Protection, Annex IV and Annex V) and the Red List of endangered and rare plant species of the Slovak and Czech Republic (Holub, 1999). Its populations have been monitored annually since 2004 for the number of both flowering and vegetative individuals, while a strong inter-annual dynamics has been recorded (Janišová *et al.*, 2005; Chmelová, 2007; Gbelcová, 2010). The main aim of this paper is to provide basic information on the reproductive biology of *T. l. moravica*, namely: (1) to clarify its mating system; (2) to estimate the seed output; (3) to determine germination percentages in cultivation; (4) to assess seedling emergence and survival in natural conditions. The investigations on the mating system were studied simultaneously in three subspecies of *T. longifolia* s.l. (*T. l. moravica*, *T. l. longifolia* and *T. l. pseudocrispa*).

Materials and methods

T. l. moravica is a long-lived perennial plant. It flowers in May and the seeds ripen in June. The inflorescence is a corymb consisting usually of 5–10 capitula (one terminal and the others located on primary or secondary branches). The 4-mm-long achenes (seeds

hereafter) have a 5–6-mm-long pappus. *T. l. moravica* is diploid $2n = 48$ (with the basic chromosome number $x = 24$; Kochjarová, 1997; Liu and Yang, 2011). The studied population sites differed in population size, their area and management (Table 1). *T. l. moravica* occupied mainly ecotone habitats, forest clearings, shrubby forest margins and shaded parts of the sub-xerophilous to mesophilous grasslands (Janišová *et al.*, 2012).

Plants of *T. l. moravica* (grown from the seeds), *T. l. longifolia* and *T. l. pseudocrispa* (both grown from transplanted vegetative rosettes) were cultivated in two experimental gardens (field conditions) in Banská Bystrica [central Slovakia, 375 m above sea level (asl)] and Bratislava (western Slovakia, 329 m asl). They were grown in pots (diameter 12–14 cm) in a substrate of commercial garden soil mixture. The location of source populations of *T. l. longifolia* and *T. l. pseudocrispa* can be found in Appendix 1.

To examine the mating system we performed isolation and emasculation experiments on cultivated plants in 2009 and 2010. Altogether, 86 corymbs (16 plants of *T. l. moravica*, 65 plants of *T. l. longifolia* and 5 plants of *T. l. pseudocrispa*) were isolated by covering them with empty tea bags before and during anthesis to prevent cross-pollination (test for autogamy). To confirm or exclude an aposporic formation of seeds in *T. l. moravica* we performed emasculation by cutting off the upper part of the capitulum containing the anthers just before anthesis, using a razor blade. In cultivation as well as in the field, seed output from open pollination was recorded. An artificial hybridization was performed between cultivated individuals of *T. l. moravica* and two other subspecies of *T. longifolia* s.l. The corymbs of the crossed plants were enclosed together within a tea bag prior to flowering. No human activity (rubbing the inflorescences together) supported the pollen exchange. The bags were




Figure 1. Distribution of *Tephrosieris longifolia* subsp. *moravica* in the Czech Republic and Slovakia. Solid circles show sites with recent evidence of taxon occurrence. Open circles show extinct populations where the historic taxon occurrence was documented by a herbarium specimen.

Table 1. Population sites of *T. longifolia* subsp. *moravica*: location, management and population size

Population site	Altitude (m asl)	Area (m ²)	Current management	Population size (number of all individuals/number of flowering shoots)		
				2008	2009	2010
Čavoj	560–585	20,000	Mostly abandoned, various stages of succession	743/737	399/345	331/170
Lysá	740–780	12,000	Long time without management or irregularly grazed	379/559	448/597	299/370
Omšenie	570–670	11,000	Site of one subpopulation mown, two subpopulations without management	122/90	188/196	169/171
Radobica	480–560	15,000	Mostly mown, partly without management	1810/2518	2048/781	2032/472
Stráž	770–780	5000	Abandoned for decades and covered by large shrubs	68/67	197/112	246/13

removed after ripening of seeds. Mature seeds were collected and classified as well-developed or undeveloped. Although most of the tea bags were undamaged at the time of collection, entry of insects (mostly earwigs, ants and aphids) into the bags was quite frequent and some seeds were even damaged by granivory.

As *T. l. moravica* populations are limited in number and size, seeds for estimation of seed output and germination were collected only from the two biggest natural populations (Čavoj/66 plants, Radobica/98 plants) and from a cultivated population of self-established plants originating from four individuals of the Omšenie population (Omšenie II). Seed output was characterized by three variables, total number of seeds per capitulum, number and percentage of well-developed seeds per capitulum. Well-developed capitula with undispersed seeds were collected on 23 June 2008, while the type of habitat (ecotone versus grassland) and position of capitulum within the corymb (terminal versus primary branch versus secondary branch) were recorded. Several morphological characteristics related to the size of the mother plant were recorded: number of both flowering and vegetative shoots, height of stem and number of capitula in the corymb. Capitula were dried in a warm place for a week. Number of seeds per capitulum was recorded while distinguishing the well-developed, undeveloped and damaged seeds. While the well-developed seeds were large in size, plump and well pigmented (dark brown), poorly developed seeds were narrow, flimsy and weakly pigmented (usually yellowish or greenish). Poorly developed seeds always lack embryos while well-developed seeds may or may not contain an embryo, therefore the well-developed seeds were checked by pinching them gently with forceps and only the firm ones were selected for the experiments. In capitula heavily damaged by granivorous insects it was impossible to distinguish well-developed seeds from undeveloped ones, thus the seed output was counted only in undamaged or slightly damaged capitula. Two-way ANOVA was

used to study differences among habitats and capitulum position, while the effect of population (Čavoj versus Radobica) was analysed separately by a *t*-test on arcsine transformed data.

Germination experiments were conducted on seed samples originating from two natural populations (Čavoj, Radobica) and cultivated population Omšenie II. For each population, two sets of 300 well-developed seeds were selected at random. One set was frozen at -25°C for testing later (after 3 months) to investigate the effect of deep freeze storage on seed germination. The second set was stored in dry and warm conditions (room temperature) for 4 months. Both treatments were germinated in Petri dishes (6 replicates with 50 seeds) on wet filter paper for 12 weeks. Petri dishes with a diameter of 9 cm and distilled water (4.5 ml per dish) were used. Prior to sowing, the seeds were disinfected for 5 min in a weak solution of permanganate and then washed with distilled water. The germination experiment was performed from November 2008 to March 2009 in the laboratory with a natural daily photoperiod, the dark period temperatures $16\text{--}19^{\circ}\text{C}$ and the light period temperatures $19\text{--}24^{\circ}\text{C}$. Emergent seedlings were counted and removed every 1 or 2 days. Based on visual inspection and pinching, all well-developed seeds were assumed to contain embryos; however, the seed viability was not tested. Germination was measured as cumulative percentage of the well-developed seeds that germinated. Data on germination percentages were square-root and arcsin transformed. Two-way ANOVA included the fixed factors population (3 levels) and type of storage (2 levels), respectively. Tukey's post-hoc tests were used for pairwise comparisons. Starting day of germination was analysed by two-way ANOVA without previous transformation.

Seedling emergence and survival of *T. l. moravica* in field conditions was studied in fifty 25×25 cm plots arranged in transects. In each of five investigated population sites (Čavoj, Lysá, Omšenie, Radobica and Stráž), two transects were established with five experimental plots each (ten plots per site). The control

plots of the same size were situated in the adjacent area to each of the plots. In June 2009, three capitula with well-developed seeds (i.e. approximately $3 \times 63 = 189$ seeds, calculated from the average number of well-developed undamaged seeds per capitulum) were spread in each plot. The seedlings were checked in May 2010 and in August 2010 and their position within the plot was recorded on a map. Two variables were distinguished: (1) number of seedlings that emerged since sowing (calculated as cumulative number of seedlings recorded in the plot in May and August 2010 minus number of seedlings found in the adjacent control plot); (2) number of seedlings that survived until August 2010 (calculated as number of seedlings and juveniles recorded during the August census minus number of seedlings and juveniles recorded in the adjacent control plot). One-way ANOVA was used to check differences among sites and individual transects (after adding 0.5 to each recorded value, square-root and then arcsin transformation was used).

Results

Mating system and hybridization in *T. longifolia* s.l.

In the isolation experiments, well-developed seeds were recorded in 4 from 16 cultivated plants (25%) and 0 from 5 plants in a natural population (0%) of *T. l. moravica*; 7 from 65 cultivated plants (11%) of *T. l. longifolia*; and 1 from 5 cultivated plants (20%) of *T. l. pseudocrispa*. In *T. l. moravica* and *T. l. longifolia*, the number of well-developed seeds per corymb did not exceed two (maximum of 0.25% of well-developed seeds per corymb). In *T. l. pseudocrispa*, 28 well-developed seeds (3.26%) were recorded in one plant; the other four plants had all undeveloped seeds. Emasculation of five plants of *T. l. moravica* in a natural population and one plant of *T. l. moravica* in cultivation revealed that none of the plants were able to form well-developed seeds. Under open pollination in *T. l. moravica* (non-isolated corymbs) we observed high seed output during both years of experiments: all of 112 investigated plants in natural populations and 25 cultivated plants formed well-developed seeds, the percentage of well-developed seeds per capitulum being from 21 to 95% (average 75%) in field conditions and from 15 to 88% (average 49%) in cultivation (Table 2). In hybridization experiments, all controlled pollinated (two pairs of plants) as well as cross-pollinated plant combinations (seven pairs of plants) produced well-developed seeds (Table 2). The percentage of well-developed seeds per capitulum (between 0.32 and 3.65%) was markedly lower than that recorded for open pollination, but higher than in the isolation experiments.

Seed output in *T. l. moravica*

There was no difference in total number of seeds per capitulum between the populations (Table 3), mean values being 95 in Čavoj and 101 in Radobica populations. However, there was a significant difference between habitats (three-way ANOVA, $P = 0.005$), capitula had more seeds in ecotone habitats (mean value of 103 seeds) than in grassland (92 seeds). Capitula with different positions within a corymb differed significantly in total number of seeds ($P = 0.002$), terminal capitula having the highest number (mean value of 113 seeds). Capitula on primary branches had an intermediate number (97) and capitula on secondary branches had the smallest total number of seeds (87). The effect of factor interactions was non-significant (population \times position: $P = 0.611$, habitat \times position: $P = 0.301$, population \times habitat \times position: $P = 0.600$), although the interaction population \times habitat had the borderline value $P = 0.055$, indicating that the difference in total number of seeds per capitulum between habitats was dependent on population (valid for Radobica and not for Čavoj population). Total number of seeds per capitulum was not significantly correlated with morphological characteristics of the mother plants (number of flowering and vegetative shoots, height of stem and number of capitula in the corymb).

Mean percentage of well-developed seeds per capitulum was 66.23% in Čavoj and 74.18% in Radobica populations (t -test, $P = 0.076$). Neither habitat and position of capitulum, nor their interaction, had a significant effect on percentage of well-developed seeds (two-way ANOVA, habitat: $P = 0.615$, position: $P = 0.257$, habitat \times position: $P = 0.626$). The populations differed in number of well-developed seeds per capitulum (t -test, $P = 0.003$) with higher values in Radobica (mean 80 seeds) than in Čavoj (63 seeds). The effect of habitat on number of well-developed seeds was not significant (two-way ANOVA, $P = 0.104$). The position of a capitulum within the corymb showed a significant effect on the number of well-developed seeds ($P = 0.011$), terminal capitula showed the highest values and capitula on primary and secondary corymb branches did not differ (Tukey's multiple comparison test: terminal versus primary branch $P = 0.041$, terminal versus secondary branch $P = 0.009$ and primary branch versus secondary branch $P = 0.994$). The interactions of habitat and capitulum position had no significant effects on percentage and number of well-developed seeds ($P = 0.834$). The percentage and number of well-developed seeds were not significantly correlated with morphological characteristics of the mother plants (number of flowering and vegetative shoots, height of stem and number of capitula in the corymb). They were positively correlated with total number of seeds

Table 2. Results of experiments focusing on mating system in the studied populations of *Tephrosieris longifolia* subsp. *moravica*, *T. l.* subsp. *longifolia* and *T. l.* subsp. *pseudocrispa*

Population	Source data	Year	Number of investigated plants/capitula	Mean number of capitula per corymb	Mean total number of seeds per investigated capitulum	Mean % of well-developed seeds per plant (% of plants with well-developed seeds)
Open pollination						
<i>Tephrosieris longifolia</i> subsp. <i>moravica</i>						
Čavoj	Field conditions	2008	40/40	9	95	66.23 (100)
Čavoj	Field conditions	2009	13/13	?	74	63.83 (100)
Lysá	Field conditions	2009	21/21	?	88	78.78 (100)
Radobica	Field conditions	2008	19/19	9	101	74.18 (100)
Stráž	Field conditions	2009	19/19	?	96	77.85 (100)
Omšenie II	Cultivation	2010	16/16	?	132	51.04 (100)
Omšenie	Cultivation	2008	4/9	?	137	42.08 (100)
Isolation experiments						
<i>Tephrosieris longifolia</i> subsp. <i>moravica</i>						
Čavoj	Cultivation	2010	4/23	6	102*	0.13 (50)
Omšenie	Cultivation	2007	1/4	4	?	0 (0)
Omšenie II	Cultivation	2010	8/26	8	108*	0.04 (11)
Radobica	Cultivation	2010	3/16	6	119*	0.05 (33)
<i>Tephrosieris longifolia</i> subsp. <i>longifolia</i>						
Eberstein	Cultivation	2010	7/44	6	79*	0.03 (14)
Hirskeuche	Cultivation	2010	6/39	7	91*	0.08 (33)
Loiblpass	Cultivation	2010	16/121	8	95*	0.17 (6)
Pitten	Cultivation	2010	4/19	5	83*	0 (0)
Sv. Jacob	Cultivation	2010	7/44	6	67*	0 (0)
Sv. Lorenz	Cultivation	2010	12/96	8	85*	0.04 (8)
Trdinov vrh	Cultivation	2010	13/76	6	55*	0.15 (15)
<i>T. longifolia</i> subsp. <i>pseudocrispa</i>						
Predil	Cultivation	2010	5/47	10	86*	0.69 (20)
Controlled pollination and cross-pollination experiments						
Parent population 1	Parent population 2	Year	Number of capitula in parent 1	Number of capitula in parent 2	% of well-developed seeds in both parents	Notes
<i>Tephrosieris longifolia</i> subsp. <i>moravica</i> × <i>Tephrosieris longifolia</i> subsp. <i>moravica</i>						
Radobica	Omšenie II	2010	7	9	0.89	
<i>Tephrosieris longifolia</i> subsp. <i>moravica</i> × <i>Tephrosieris longifolia</i> subsp. <i>longifolia</i>						
Omšenie II	Pitten	2010	7	6	0.32	
Omšenie II	Eberstein	2010	4	4	2.41	Bag penetrated by insects
Omšenie II	Eberstein	2010	6	6	3.65	
Omšenie II	Eberstein	2010	8	7	0.36	Bag penetrated by insects
Omšenie II	Eberstein	2010	4	10	1.15	Bag penetrated by insects
Omšenie II	Hirskeuche	2010	7	9	0.76	Bag penetrated by insects
<i>Tephrosieris longifolia</i> subsp. <i>longifolia</i> × <i>Tephrosieris longifolia</i> subsp. <i>longifolia</i>						
Pitten	Eberstein	2010	4	5	6.46	
<i>Tephrosieris longifolia</i> subsp. <i>longifolia</i> × <i>Tephrosieris longifolia</i> subsp. <i>pseudocrispa</i>						
Pitten	Predil	2010	6	10	1.88	

*Means calculated from subsample of 10 capitula.

Table 3. Characteristics of *T. longifolia* subsp. *moravica* individuals related to their size and seed output in two studied populations

Measured characteristics	All investigated individuals Mean \pm SD (min.–max.)	Čavoj (C) Mean	Radobica (R) Mean	Differences (C versus R)
Total number of seeds per capitulum	98 \pm 26 (44–178)	95	101	n.s.
Number of well-developed seeds per capitulum	68 \pm 22 (9–123)	63	80	C < R
Percentage of well-developed seeds per capitulum	69 \pm 18 (9–92)	66	74	n.s.
Percentage of damaged seeds per capitulum	24 \pm 27 (0–100)	18	28	C < R
Percentage of damaged capitula in a population		39	81	C < R
Number of capitula per corymb	8.73 \pm 4.17 (3–31)	8.62	8.8	n.s.
Number of flowering shoots per flowering plant	1.9 \pm 1.57 (1–9)	1.36	2.27	C < R
Number of vegetative shoots per plant	3.48 \pm 2.8 (1–14)	2.27	4.29	C < R
Height of a flowering stem (cm)	71 \pm 14 (38–117)	65	75	C < R

per capitulum (Pearson correlation coefficient, $r = 0.59$ for number of well-developed seeds and $r = 0.65$ for percentage of well-developed seeds). In the population of cultivated plants (Omšenie II), total number of seeds per capitulum was larger (mean number 141, range 109–179 seeds), but the percentage of well-developed seeds was lower (mean value 42.08%, range 30–63%) than in natural populations in Čavoj and Radobica. The output of well-developed seeds per flowering shoot estimated from mean number of well-developed seeds per capitulum (68 seeds) and mean number of capitula per corymb (8.7 capitula) was 592 (1125 seeds per flowering plant, calculated as 592×1.9 which is the mean number of flowering shoots per flowering plant).

In all studied populations, a certain proportion of seeds were damaged by the caterpillar of a granivorous butterfly from the family Pyralidae. The butterfly has been identified as *Phycitodes albatella* Ragonot 1887. Adult butterflies (wing span 16–21 mm) deposit eggs in the upper stem and leaf-axils. The orange-coloured larva tunnels into the capitulum and out to the developing seeds where it feeds. Mature larvae exit host capitula in summer to pupate. Evidence of infestation is obvious: the pappus of seeds is threaded with a web. Larval feeding frequently destroys most seeds within the mined capitulum. Seeds damaged by the granivorous butterfly in 2008 represented 18% of

seed output in the population of Čavoj and 28% in the population of Radobica. The proportion of capitula attacked was significantly higher in Radobica (80.6%) than in Čavoj (39.4%, $P = 0.027$), and it was higher in grasslands (71%) than in ecotone habitats (59%), although this difference was non-significant ($P = 0.058$). The probability of attack by granivores was not affected by the position of the capitulum within the corymbs. Once the capitulum was attacked by the granivorous butterfly, the number of damaged seeds ranged from 1 to 87 (mean 33 seeds). As a consequence, the mean reduction of seed number in attacked capitula was 37% (range 0.85–100%).

Germination in laboratory conditions

Seeds from natural populations in Čavoj and Radobica had significantly higher germination percentages (30–62%, mean 50% for Čavoj and 51% for Radobica) than seeds from cultivated plants (population Omšenie II, 18–42%, mean 29%, two-way ANOVA, $P < 0.001$; Table 4). There was no difference between Radobica and Čavoj in germination percentages of seeds stored dry at room temperatures. Deep freeze storage for 3 months appeared to stimulate germination in seeds from the two natural populations (Čavoj and Radobica,

Table 4. Germination in laboratory conditions. Final germination percentages and average starting day of germination for three populations of *T. longifolia* subsp. *moravica* under dry storage in room temperature for 4 months and deep freeze storage for 3 months

Population and storage type	Final germination percentages Mean (%) \pm SD	Average starting day of germination Mean (days) \pm SD
Čavoj, dry storage	49.7 \pm 14.8	11.9 \pm 1.1
Čavoj, deep freeze storage	55.3 \pm 5.6	12.3 \pm 1.1
Radobica, dry storage	50.7 \pm 8.1	13.0 \pm 2.2
Radobica, deep freeze storage	65.0 \pm 10.7	14.7 \pm 2.4
Omšenie II, dry storage	29.0 \pm 8.0	20.3 \pm 0.9
Omšenie II, deep freeze storage	28.0 \pm 5.7	19.5 \pm 2.2

Table 4), although its effect was non-significant ($P = 0.052$). The interaction of population and deep freeze storage was also non-significant ($P = 0.154$).

Seeds stored in dry and warm conditions started to germinate on days 5–8 after sowing, while seeds stored in deep freeze started to germinate a few days later (on the seventh day in Čavoj and between days 8 and 11 in Radobica). Seeds from the population in Čavoj germinated slightly earlier in all treatments than seeds from Radobica (Table 4). Seeds from the cultivated population Omšenie II started to germinate the latest (on days 10–11 after sowing in the two studied treatments). Most seeds germinated within 30 days of sowing. Deep freeze storage and its interaction with population had a non-significant effect on the average starting day of germination ($P = 0.446$ and $P = 0.248$, respectively).

Seedling emergence and survival in field conditions

In each plot, 0–31 seedlings were recorded in May 2009 (11 months since sowing) and 0–25 seedlings or juveniles survived until the census in August 2010 (14 months since sowing). If expressed as proportion of the sown seeds, the maximum percentage of emerged seedlings was 16.4% and maximum percentage of survived seedlings was 13.23%. The studied populations did not differ significantly in percentage of emerged seedlings (one-way ANOVA, $P = 0.109$), or in percentage of survived seedlings (one-way ANOVA, $P = 0.167$). Seedling emergence (Table 5) was the highest in habitats with open soil surface (forest and bush understorey), and the lowest in habitats with high litter or moss layers. Seedling survival (Table 5) was the highest in transects least disturbed by man and animals (dense bush stands and remote areas). The highest seedling survival was

recorded in populations of Stráž where all seedlings emerged by May survived until the August census. On the other hand, mortality of seedlings was the highest in populations of Radobica and Lysá, where experimental plots were strongly disturbed by moles and game animals (Radobica) and by cattle grazing (Lysá). Seedling emergence and establishment were much slower in natural conditions than in cultivation (14 months after sowing *in situ* the seedlings had hardly developed true leaves and were still very tiny).

Discussion

Mating system and hybridization

Our results of isolation experiments showed that the studied subspecies of *T. longifolia* s.l. are not autogamous taxa. The low number of well-developed seeds obtained in some isolated capitula (Table 2) could be the result of accidental pollen transfer (insect penetration) or sporadic break-down of the self-incompatibility system. These sporadic break-downs can be attributed to specific characteristics of sporophytic self-incompatibility systems, the so-called 'leaking' (Lewis, 1994; Hiscock, 2000), when even self-pollination produces a small number of seeds. Similar to our results, few seeds were set by self-pollinated plants of closely related *Tephroseris integrifolia* (Isaksson, 2009). Based on our results, *T. l. moravica* should be treated as an allogamous taxon with a sporophytic self-incompatibility system. However, it is necessary to test these results on a larger data set. The results of our emasculation experiment (in spite of the small sample size) suggest that apomictic reproduction does not occur in the studied taxon. There are no records in the scientific literature of apomixis in the genus *Tephroseris* (Czapik, 1996; Noyes, 2007).

Table 5. Seedling emergence and survival of *Tephroseris longifolia* subsp. *moravica* in field conditions. Number of seedlings emerged in plots between the sowing in June 2009 and census in August 2010 and number of seedlings survived until August 2010 are shown for each transect consisting of five experimental plots

Transect	Seedlings emerged	Seedlings survived	Survival %	Habitat characteristics
	mean \pm SD (min.–max.)	mean \pm SD (min.–max.)		
Čavoj 1	0	0		Abandoned grassland (high moss and litter cover)
Čavoj 2	3.4 \pm 4.9 (0–12)	2.2 \pm 4.9 (0–11)	64	Patchy grassland under a cherry tree
Lysá 1	3.4 \pm 3.0 (0–8)	1.6 \pm 3.6 (0–8)	47	Mesophilous grassland (occasional grazing by cattle)
Lysá 2	0	0		Forest margin (high moss and litter cover)
Omšenie 1	0.8 \pm 1.3 (0–3)	0.6 \pm 1.3 (0–3)	75	Mesophilous grassland
Omšenie 2	7.0 \pm 13.5 (0–31)	5.6 \pm 10.9 (0–25)	80	Shrub understorey (open soil surface)
Radobica 1	6.0 \pm 5.5 (0–11)	1.9 \pm 3.3 (0–8)	56	Shrub understorey (open soil surface, high disturbance)
Radobica 2	1.1 \pm 2.5 (0–6)	0	0	Forest margin (high moss cover)
Stráž 1	2.0 \pm 2.5 (0–6)	2.0 \pm 2.5 (0–6)	100	Shrub understorey (open soil surface)
Stráž 2	5.8 \pm 4.3 (0–11)	5.8 \pm 4.3 (0–11)	100	Mesophilous grassland
All transects	3.0 \pm 5.4 (0–31)	2.0 \pm 4.5 (0–25)	65	

The low percentage of well-developed seeds in both controlled pollination and cross-pollination experiments could be a consequence of differences in phenology of cross-pollinated populations. The number of well-developed seeds was higher in the controlled pollination, 16 (0.89%) for *T. l. moravica* and 47 (6.46%) for *T. l. longifolia*, where the phenological development of pollinated plants was most similar. Another possible reason for the low seed production is the methodology used (the corymbs were enclosed without rubbing). However, similar to our results, a markedly decreased number of well-developed seeds in controlled pollination and cross-pollination experiments in comparison to that obtained from open pollination was recorded in the *Cyanus triumfetti* group, in spite of rubbing the capitula together (K. Olšavská, unpublished results). Considering the low number of investigated plants and the methodology used, our results should be taken as preliminary.

Seed output

Population, habitat and position of capitula within corymbs had no effect on percentage of well-developed seeds. As a consequence of terminal capitula being the largest, the number of well-developed seeds ripened in them was higher than in the capitula on side branches. The possible explanation is that plants invest in the formation of an early flowering large terminal capitulum in order to utilize available pollinators and to prolong the period of seed dispersal.

Mean number of seeds per capitulum, 95 in Čavoj and 101 in Radobica populations, are similar to numbers reported by Chmelová (2007) from the Czech populations of Tratihuš and Hodňov (96 seeds) and Kochjarová (1998) from populations of Lysá, Stráž and Radobica (80–128 seeds). The mean percentage of well-developed seeds (66.23% in Čavoj and 74.18% in Radobica) is similar to that found by Kochjarová (1998; 53.89–78.67%) and slightly lower than that recorded by Chmelová (2007; 84.62%). The estimated mean number of well-developed seeds per plant was higher in our study (1125 seeds) than in Kochjarová (1998; 727–822 seeds).

Along with seed abortions, granivory represents one of the predispersal losses that may cause significant reductions in total seed yields in wild populations. In 2008, the granivorous butterfly *P. albatella* reduced the number of seeds by almost one-third in the Radobica population. According to Tlusták (unpublished), in 1996 two-thirds of seeds were damaged by insects in Czech populations of Tratihuš, Uhličky and Hluboče. Although the granivores alone could hardly cause the extinction of *T. l. moravica* populations, in combination with other

factors (such as unsuitable weather conditions during several subsequent years, low flowering intensity of a population due to demographic fluctuations, etc.) they could reduce population generative reproduction even to zero. It is also known from other species of Asteraceae (genus *Cirsium*), that herbivory in combination with the demographic characteristics may strongly influence population dynamics and species distribution (Münzbergová, 2005).

Germination in laboratory conditions

Dry storage evidently affects germination of *T. l. moravica* by fast reduction of germination percentages with storage duration. Several authors recorded higher germination percentages immediately after seed ripening, e.g. 70% by Kochjarová (1998) and 72% by Chmelová (2007). If stored at low temperature or buried in soil, high germination ability of *T. l. moravica* seeds can be maintained for longer (Chmelová, 2007; personal observations), although the upper survival time limit remains unknown.

Seeds from natural populations (Čavoj and Radobica) had significantly higher germination percentages and germinated faster than plants from a cultivated population (Omšenie II). Together with a higher percentage of undeveloped seeds in the capitula of cultivated plants (Table 2), the poorer germination characteristics can strongly reduce their fitness. The reasons for the reduced fitness in cultivated plants may include, for example, lack of pollinators in non-natural conditions, different climate and soil conditions or inbreeding depression (Jennersten, 1988; Isaksson, 2009).

Seedling emergence and survival in field conditions

It seems that *T. l. moravica* has very low seedling emergence in all natural populations and its recruitment from seeds is much dependent on the availability of microsites suitable for germination. In our experiment, the most suitable habitats for germination included understorey of trees and shrubs with open soil surface and loose, shaded grasslands. Thick moss and litter layers suppressed seedling emergence in both studied habitat types – forest margins (ecotone) and grasslands. The general mechanisms of germination prevention by litter are well understood (Facelli and Pickett, 1991). Another important factor may be the reduced rooting capacity for seedlings within the bryophyte layer, as proposed by van Tooren (1988). The presence of gaps with little competition from other species may also be very important for successful seedling establishment of *T. l. moravica*. However, heavy disturbance (e.g. by grazing or trampling

animals) or increased occurrence of slugs feeding on forb seedlings may affect seedling survival of *T. l. moravica* and strongly reduce the number of established plants even in otherwise suitable microsites.

Conclusion

According to our results, *T. l. moravica* is an allogamous taxon without apomictic reproduction. Each flowering shoot may produce several hundreds of well-developed seeds. This number is usually reduced by the incidence of granivorous pests. The germination rapidly decreases in time, so if unsuitable weather conditions hinder the germination of seeds immediately after their dispersal, their chances of germinating later in the year are low. Seedling emergence *in situ* is very low in all studied sites. Proper conservation management could increase the availability of suitable microsites for germination and thus support seedling recruitment.

Acknowledgements

We are grateful to A. Čarni, J. Kochjarová, K. Krajčovičová, J. Kulfan, K. Olšavská, J. Panigaj, M. Slaničanová, J. Smatanová, L. Sutcliffe, B. Singliarová, D. Treplanová and A. Tribsch. This research was supported financially by grants VEGA 2/0017/08 and VEGA 2/0074/11.

References

- Aeschimann, D., Lauber, K., Moser, D.M. and Theurillat, J.P. (2004) *Flora Alpina. Ein Atlas sämtlicher 4500 Gefäßpflanzen der Alpen*. Bern, Haupt Verlag.
- Byers, D.L. and Meagher, T.R. (1997) A comparison of demographic characteristics in a rare and a common species of *Eupatorium*. *Ecological Applications* **7**, 519–530.
- Chmelová M. (2007) Současný stav populací endemického *Tephroseris longifolia* subsp. *moravica* v Bílých Karpatech. Thesis, Charles University, Prague.
- Cron, G.V., Balkwill, K. and Knox, E.B. (2009) Biogeography, rarity and endemism in *Cineraria* (Asteraceae–Senecioneae). *Botanical Journal of the Linnean Society* **160**, 130–148.
- Czapik, R. (1996) Problems of apomictic reproduction in the families *Compositae* and *Rosaceae*. *Folia Geobotanica et Phytotaxonomica* **31**, 381–387.
- Facelli, J.M. and Pickett, S.T.A. (1991) Plant litter: its dynamics and effects on plant community structure. *Botanical Review* **57**, 1–32.
- Farnsworth, E.J. (2007) Plant life history traits of rare versus frequent plant taxa of sandplains: implications for research and management trials. *Biological Conservation* **136**, 44–52.
- Gbelcová A. (2010) *Tephroseris longifolia* subsp. *moravica* v České republice – populační a ekobiologická studie. Thesis, University of Palacký, Olomouc.
- Greuter W. (2006–2009) *Compositae* (pro parte majore). in Greuter, W.; von Raab-Straube, E. (Eds) *Compositae*. Euro + Med Plantbase – the information resource for Euro-Mediterranean plant diversity. Available at <http://ww2.bgbm.org/EuroPlusMed/query.asp> (accessed 8 December 2010).
- Hiscock, S.J. (2000) Genetic control of self-incompatibility in *Senecio squalidus* L. (Asteraceae): a successful colonizing species. *Heredity* **85**, 10–19.
- Hiscock, S.J. and Tabah, D.A. (2003) The different mechanisms of sporophytic self-incompatibility. *Philosophical Transactions of the Royal Society of London – Series B: Biological Sciences* **358**, 1037–1045.
- Holub, J. (1999) *Tephroseris longifolia* (Jacq.) Griseb. et Schenk subsp. *moravica* Holub. p. 371 in Čerovský, J.; Feráková, V.; Holub, J.; Maglocký, Š.; Procházka, F. (Eds) *Červená kniha ohrozených a vzácných druhov rastlín a živočíchov SR a ČR. Vyššie rastliny*. Bratislava, Príroda.
- Isaksson K. (2009) Investigating genetic factors behind the decline of a threatened plant species – *Tephroseris integrifolia* (Asteraceae). Doctoral thesis, University of Lund, Sweden.
- Janišová, M., Škodová, I., Smatanová, J., Jongepierová, I. and Kochjarová, J. (2005) *Tephroseris longifolia* subsp. *moravica* – population size evaluation and possibilities of its conservation. pp. 29–38 in Franc, V. (Ed.) *Strážovské vrchy Mts – research and conservation of the nature*. Proceedings from the conference, Belušké Slatiny, October 2004. Zvolen, Slovakia.
- Janišová M., Hegedúšová K., Král P. and Škodová I. (2012) Ecology and distribution of *Tephroseris longifolia* subsp. *moravica* in relation to environmental variation at a micro-scale. *Biologia* **67**, 97–109.
- Jeffrey, C. and Chen, Y.L. (1984) Taxonomic studies on the tribe Senecioneae (Compositae) of Eastern Asia. *Kew Bulletin* **39**, 205–446.
- Jennersten, O. (1988) Pollination in *Dianthus deltooides* (Caryophyllaceae): effects of habitat fragmentation on visitation and seed set. *Conservation Biology* **2**, 359–366.
- Kliment, J. (1999) Komentovaný prehľad vyšších rastlín flóry Slovenska, uvádzaných v literatúre ako endemické taxóny. *Bulletin Slovenskej Botanickéj Spoločnosti* **21** (suppl. 4), 325.
- Kochjarová, J. (1997) Náčrt taxonomickej problematiky rodu *Tephroseris* v Západných Karpatoch. *Preslia* **69**, 71–93.
- Kochjarová, J. (1998) Poznámky k rozšíreniu, cenológii a ohrozenosti populácií zástupcov rodu *Tephroseris* (Rchb.) Rchb. na Slovensku II.: *Tephroseris longifolia* subsp. *moravica* v Západných Karpatoch. *Bulletin Slovenskej Botanickéj Spoločnosti* **20**, 69–79.
- Kunin, W.E. and Gaston, K.J. (1993) The biology of rarity: patterns, causes and consequences. *Trends in Ecology and Evolution* **8**, 298–301.
- Lewis, D. (1994) Gametophytic–sporophytic incompatibility. pp. 88–101 in Williams, E.G.; Clarke, A.E.; Knox, R.B. (Eds) *Advances in cellular and molecular biology of plants. Genetic control of self-incompatibility and reproductive development in flowering plants*. Dordrecht, Kluwer Academic Publishers.
- Liu, Y. and Yang, Q.R. (2011) Cytology and its systematic implications in *Sinosenecio* (Senecioneae–Asteraceae) and two closely related genera. *Plant Systematics and Evolution* **291**, 7–24.

- Münzbergová, Z.** (2005) Determinants of species rarity: population growth rates of species sharing the same habitat. *American Journal of Botany* **92**, 1987–1994.
- Nordenstam, B.** (2007) Tribe *Senecioneae*. pp. 208–241 in Kadereit, J.W.; Jeffrey, C. (Eds) *The families and genera of vascular plants*. Vol. 8. Berlin, Springer.
- Noyes, R.D.** (2007) Apomixis in the Asteraceae: diamonds in the rough. *Functional Plant Science and Biotechnology* **1**, 207–222.
- Purdy, B.G., Bayer, R.J. and MacDonald, S.E.** (1994) Genetic variation, breeding system evolution, and conservation of the narrow endemic *Stellaria arenicola* and the widespread *S. longipes* (Caryophyllaceae). *American Journal of Botany* **81**, 904–911.
- Scheidel, U. and Bruelheide, H.** (2005) Effects of slug herbivory on the seedling establishment of two montane Asteraceae species. *Flora* **200**, 309–320.
- van Tooren, B.F.** (1988) The fate of seeds after dispersal in chalk grassland: the role of the bryophyte layer. *Oikos* **53**, 41–48.
- Wagenitz, G.** (1987) *Senecio*. pp. 1374–1383 in Hegi, G. (Ed.) *Illustrierte Flora von Mitteleuropa*. Vol. VI/4. Berlin-Hamburg, Paul Parey.

Appendix 1

Geographical location of source populations of *T. l. longifolia* and *T. l. pseudocrispa* used in isolation and cross-pollination experiments: Eberstein (Austria, 570–622 m asl, 46°47'51"N, 14°33'07"E), Hirskeuche (Austria, 740–775 m asl, 46°28'14"N, 14°29'25"E),

Loiblpass (Austria, 990–1005 m asl, 46°26'41"N, 14°15'28"E), Pitten (Austria, 320–340 m asl, 47°42'28"N, 16°10'53"E), Sv. Jakob (Slovenia, 780–790 m asl, 46°06'19"N, 14°22'11"E), Sv. Lorenz (Slovenia, 780–790 m asl, 46°04'18"N, 14°17'59"E), Trdinov vrh (Slovenia, 1130–1180 m asl, 45°45'35"N, 15°19'22"E), Predil (Italy, 880–907 m asl, 46°27'00"N, 13°34'31"E).