

ALPINE HEATHS IN THE WESTERN CARPATHIANS - A NEW APPROACH TO THEIR CLASSIFICATION

Z. DÚBRAVCOVÁ*, I. JAROLÍMEK**, J. KLIMENT***, A. PETRÍK****,
J. ŠIBÍK**, M. VALACHOVIČ**

*Department of Botany, Faculty of Natural Sciences, Comenius University, Révová 39,
SK-811 02 Bratislava, Slovak Republic, e-mail: dubravcova@fns.uniba.sk

**Institute of Botany, Slovak Academy of Sciences, Dúbravská cesta 14, SK-845 23 Bratislava,
Slovak Republic, e-mail: ivan.jarolimek@savba.sk, jozef.sibik@savba.sk,
milan.valachovic@savba.sk

***Botanical Garden of Comenius University, SK-038 15 Blatnica, Slovak Republic,
e-mail: kliment@rec.uniba.sk

****Botanical Garden of Comenius University, Botanická 3, SK-841 04 Bratislava,
Slovak Republic, e-mail: petrika@rec.uniba.sk

ABSTRACT - Using physiognomic parameters (structure), qualitative and quantitative parameters (presence or absence of significant components (life-forms, chorological elements), abundance of plant species and relation among them (spatial pattern) were classified alpine heaths and contact vegetation habitats. Four clusters were recognized - beside own alpine heath class (*Loiseleurio-Vaccinietea*) also three grassland-like habitats (class *Carici rupestris-Kobresietea bellardii*, class *Caricetea curvulae* and *Nardus stricta*-rich grasslands). Each vegetation type is shortly defined with respect to regional peculiarities in the Western Carpathians.

KEYWORDS - Alpine vegetation, *Loiseleurio-Vaccinietea*, Natura 2000.

INTRODUCTION

In the Western Carpathians alpine heaths were recently classified within the large defined class *Juncetea trifidi*. This class encompassed various stands on soils with acidic to neutral reaction, including alpine grasslands on outcrops and flat to steep slopes, and dwarf scrub heaths forming mosaic vegetation with other above mentioned types. Using floristic criterion there were no sufficient differences, though ecological and physiognomic differences are visible on first look. In the period of mapping programs such as NATURA 2000, there is optimal (and practical) to recognize stands according to their structure and ecological demands, which are connected with following nature protection and management plans.

MATERIAL AND METHODS

The syntaxonomical revision included 1760 phytocoenological relevés of the wind edge naked-rush, dwarf-scrub heaths and grasslands of the mountains of the Western Carpathians. Contextually it relates to the recent study on the Western Carpathians high-mountains communities of the class *Carici rupestris-Kobresietea bellardii* (Petrík *et al.*, 2005).

All relevés used in this study have been collected according to the traditional principles of Zürich-Montpellier school. Different scales of abundance and dominance have been used by various authors: the five- or seven-degree Braun-Blanquet's scale, the 10-degree, or the combined 11-degree Hadač and Domin's scale (cf. Sillinger, 1933; Hadač *et al.*, 1969) and a modified 9-degree scale (Barkman *et al.*, 1964). To obtain the data comparable within the numerical classification, all relevés were transformed into the ordinal scale (van der Maarel, 1979). The taxa determined only at the level of genus were excluded and some taxa were classified within the higher or more broadly defined taxa.

Numerical classification was performed by the program NCLAS from the SYNTAX 2000 package (Podani, 2001). The β -flexible method ($\beta = -0.25$) with Euclidian distance and Jaccard's, Ružička's and Wishart's similarity coefficients were used. Obtained hypotheses were evaluated by comparison and analysis of phytocoenological tables processed by program FYTOPACK (Jarolímek & Schlosser, 1997) and simple statistics using life forms and geo-elements accepted according to Dostál (1989).

The contents of each column of the shortened synoptic table (TABLE 1) comprise the number of relevés used for the synthesis, and the average number of species in the relevant community. Each taxon is characterised by the frequency (in %; + = frequency $< 0.5\%$) and the mean value of abundance (upper index, in ordinal scale) calculated using the FYTOPACK (Jarolímek & Schlosser, 1997). Diagnostically important taxa of individual plant communities are given in bold.

The nomenclature of the taxa generally follows the Checklist of non-vascular and vascular plants of Slovakia (Marhold & Hindák, 1998).

RESULTS AND DISCUSSION

Recently in the class *Juncetea trifidi* and order *Caricetalia curvulae* four alliances were merged - *Juncion trifidi*, *Loiseleurio-Vaccinion*, *Festucion versicoloris* and *Nardion strictae* (cf. Mucina & Maglocký, 1985). First our proposal displaced from this heterogeneous vegetation the alliance *Festucion versicoloris* into the class *Carici rupestris-Kobresietea* (cf. Petrík *et al.*, 2005, p. 39). Now we upgrade alliance *Loiseleurio-Vaccinion* on the level of class along with new content definition of the alliance *Juncion trifidi*, in the recent proposal the only one alliance in the class *Caricetea curvulae*. The rest vegetation with dominance of *Nardus stricta* belongs evidently to the other acidophilous subalpine *Nardetalia* communities.




FIGURE 1 - Simplified dendrogram of the numerical classification of the data set of 1,760 phyto-coenological relevés from the Western Carpathians.

A - *Carici rupestris-Kobresietea bellardii*, B - *Caricetea curvulae* (= *Juncetea trifidi* p. p.), C - *Loiseleurio-Vaccinietea*, D - *Nardus stricta*-group.

The result of numerical classification (FIGURE 1) take into account floristic composition, quantitative values (abundance) of species as well as quantitative occurrence of plants in different layers, mainly relation between herbaceous and woody plants and cover of bryophytes. The four resulting groups (A-D) can be shortly characterised as:

Carici rupestris-Kobresietea bellardii Ohba 1974 (see TABLE 1, column A). Species rich, chionophobous, cryotemperate dwarf scrubs (*Dryas octopetala*, *Salix* spec. div.), cushion-shaped chamaephytes (*Silene acaulis*, *Minuartia* spec. div.) and graminoids (*Festuca versicolor*, *Elyna myosuroides*) on wind-exposed ridges. The soils with neutral pH are obviously calcium-enriched rendzinas. Holarctic distribution and highest summits in Europe, from Pyrenees to mountains of Balkan Peninsula, which manifest in the high number of endemic species.

Caricetea curvulae Br. Bl. 1948 nom. cons. prop. (Syn. *Juncetea trifidi* Hadač 1946 pp.), (TABLE 1, column B)

Alpine grassland-like communities (*Juncus trifidus*, *Agrostis rupestris*) on windy slopes with lower snow cover. The strictly acidic soils are alpine ranker. Boreal distribution and Alpine-Caucasian Mountain system.

Loiseleurio-Vaccinietea Eggler ex Schubert 1960 (TABLE 1, column C), Supratimberline, ericoid dwarf scrubs communities on slopes with various snow cover. Types of soils are rankers and podzols. Circumpolar distribution.

Nardus stricta-rich communities (TABLE 1, column D)

Mountain to alpine grassland-like communities on slopes with moderately long snow cover. The soils are acidic alpine rankers and podzols. Boreal distribution.

The present delimitation of plant communities which follows habitat peculiarities is based according to Theurillat *et al.* (1995) on combination of: 1) the structural homogeneity of all units inside a class, and 2) the floristic similarities. Higher units have to be defined with help of additional criteria such as: horizontal and vertical structure, their dominating life and growth forms, which reflect the distinctive ecological conditions (climate, soil) and succession stage.




FIGURE 2 - A, B, C, Comparison of the average frequency (%) in above species level taxa, chorological elements, and life forms in four selected groups.

Structure in phytosociological sense (Rejmánek, 1977) is defined as system determined by: 1) qualitative presence (or absence) of components belonging to defined class (group), e.g. grasses, arcto-alpine taxa, etc.; 2) quantitative representations of these components, e.g. abundance of plant species; and 3) relation among them, e.g. spatial pattern, etc.

An analyse based on taxa above species-level (e.g. family-level), shows clear predominance of *Salicaceae*, *Saxifragaceae* and *Silenaceae* in the group A (FIGURE 2a). Taxa such as *Saxifraga bryoides*, *S. moschata*, *S. oppositifolia* and others, as well as *Salix reticulata*, *Minuartia sedoides*, *Silene acaulis* manifest absolute preference in the class *Carici rupestris-Kobresietea bellardii* in whole Europe. In term of chorology this group is characterized by typical arcto-alpine elements - they related only this class such as geophytes *Lloydia serotina* and *Bistorta vivipara* or hemicryptophytes *Ligusticum mutellinoides*, *Pedicularis oederi*, *Erigeron uniflorus*, rarely also *Comastoma tenellum* and *Oxytropis halleri*. Boreal elements are typical

for grasslands represented here by mezophilous plant species such as *Anthoxanthum alpinum*, *Luzula sudetica*, *Ligusticum mutellina*, *Potentilla aurea*, *Geum montanum*, *Hypericum maculatum*, *Gentiana asclepiadea* etc. (FIGURE 2b).

The relations between the above mentioned families and genera and ecological conditions influence spatial arrangement of stands. Snow cover determines the distribution of vegetation types and its extreme values are limiting factor for existence of many plant taxa. Rich lichen and mosses layer in first two groups (A-B) is interacting with small chamaephytes, which utilize cryptogams for protection of seedlings (FIGURE 2c).

Absence of own character species was reason for unification of the alpine dwarf heaths into largely defined class *Juncetea trifidi* on the level of alliance in the past. According to pure floristic criteria there was insufficient basis for further division. Using quantitative parameters (dominance and abundance, biomass) and representation of prevailing life forms (dwarf scrubs) is the division into two classes acceptable and the existence of class *Loiseleurio-Vaccinietea* in Western Carpathians is confirmed. Dominance of ericoid dwarf scrubs clearly characterizes the third group C. Vertical structure of plant communities is composed from low scrubs, while role of grasses such as *Juncus trifidus*, *Agrostis rupestris*, *Avenula versicolor* is suppressed. On the other hand the absence or low abundance of grasses such as *Nardus stricta*, *Anthoxanthum alpinum*, *Phleum rhaeticum* and other species give a reason to separate scrubby vegetation from *Nardus stricta*-rich grasslands (group D).

ACKNOWLEDGEMENTS

The research was fully supported by VEGA grant n. 4041.

REFERENCES

- BARKMAN J.J., DOING H. & SEGAL S., 1964 - *Kritische Bemerkungen und Vorschläge zur quantitativen Vegetationsanalyse*. Acta Bot. Neerl. **13**: 394-419.
- DOSTÁL J., 1989 - Nová květena ČSSR 1-2. Academia, Praha.
- HADAČ E., BŘEZINA P., JEŽEK V., KUBIČKA J., HADAČOVÁ V., VONDRAČEK M., 1969 - *Die Pflanzengesellschaften des Tales "Dolina Siedmich prameňov" in der Belauer Tatra*. Vegetácia ČSSR **B2**: 5-343.
- JAROLÍMEK I. & SCHLOSSER G., 1997 - *FYTOPACK - a system of programs to process phytosociological tables*. Biologia (Bratislava) **52**: 53-59.
- MARHOLD, K. & HINDÁK, F. [eds] 1998 - Checklist of non-vascular and vascular plants of Slovakia. Veda, Bratislava, 687 pp.
- MUCINA L. & MAGLOCKÝ Š. (eds.), 1985 - *A list of vegetation units Slovakia*. Docum. Phytosoc., N.S. **9**: 175-220.
- PETRÍK J., ŠIBÍK M. & VALACHOVIČ M., 2005 - *The class Carici rupestris-Kobresietea bellardii Ohba 1974 also in the Western Carpathians*. Hacquetia, Ljubljana **4**: 33-51.
- PODANI J., 2001: SYN-TAX 2000. Computer program for data analysis in ecology and systematics for Windows 95, 98 & NT. User's manual. Scientia Publishing, Budapest, 53 pp.

REJMÁNEK M., 1977 - *The concept of structure in phytosociology with references to classification of plant communities*. Vegetatio **35**: 55-61.

SILLINGER P., 1933 - Monografická studie o vegetaci Nízkych Tater. Orbis, Praha.

THEURILLAT J.P., AESCHIMANN D., KUPFER P. & SPICHIGER R., 1995 - *The higher vegetation units of the Alps*. Coll. Phytosoc. **23**: 189-239.

VAN DER MAAREL E., 1979 - *Transformation of cover-abundance values in phytosociology and its effect on community similarity*. Vegetatio **39**: 97-114.

TABLE 1 - Shortened synoptic table with main differential taxa of the classes Carici rupestris-Kobresietea (A), Caricetea curvulae (B), Loiseleurio-Vaccinietea (C) and Nardus stricta-rich communities (D).

Class	A	B	C	D
Number of relevés	339	926	345	150
Average species number in relevé	42	22	20	20
<i>Silene acaulis</i>	76 ⁵	8 ³	1 ²	2 ²
<i>Bistorta vivipara</i>	75 ³	23 ²	12 ²	5 ²
<i>Primula minima</i>	64 ²	40 ³	15 ²	1 ²
<i>Campanula tatrae</i>	60 ²	16 ²	10 ²	19 ²
<i>Festuca versicolor</i>	56 ⁵	4 ⁵	9 ²	1 ²
<i>Lloydia serotina</i>	56 ³	1 ²	-	-
<i>Pedicularis oederi</i>	56 ²	6 ²	1 ²	-
<i>Saxifraga moschata</i>	55 ²	2 ²	1 ²	-
<i>Bartsia alpina</i>	53 ²	6 ²	8 ²	-
<i>Saxifraga paniculata</i>	51 ³	1 ³	2 ²	-
<i>Salix kitaibeliana</i>	47 ⁵	5 ²	3 ³	-
<i>Ligusticum mutellinoides</i>	47 ²	2 ³	1 ²	-
<i>Minuartia sedoides</i>	46 ³	2 ²	-	-
<i>Pedicularis verticillata</i>	45 ²	6 ²	3 ²	1 ¹
<i>Rhodiola rosea</i>	42 ²	2 ²	1 ²	1 ¹
<i>Luzula * mutabilis</i>	40 ²	2 ²	-	1 ³
<i>Cerastium eriophorum</i>	37 ²	1 ³	1 ²	-
<i>Carex fuliginosa</i>	35 ²	2 ²	-	-
<i>Minuartia gerardii</i>	29 ³	+ ²	-	-
<i>Elyna myosuroides</i>	4 ⁶	-	-	-
<i>Campanula alpina</i>	61 ²	87 ⁵	50 ³	32 ³
<i>Oreochloa disticha</i>	64 ³	78 ⁵	40 ³	4 ³
<i>Juncus trifidus</i>	46 ²	74 ⁵	50 ³	17 ²
<i>Hieracium alpinum</i>	23 ²	62 ²	52 ²	61 ²
<i>Avenula versicolor</i>	29 ²	66 ³	37 ²	53 ³
<i>Agrostis rupestris</i>	21 ²	60 ⁵	29 ³	57 ³
<i>Vaccinium myrtillus</i>	22 ²	45 ³	90 ⁵	80 ³
<i>Vaccinium vitis-idaea</i>	37 ³	44 ³	84 ⁵	27 ²
<i>Vaccinium gaultherioides</i>	13 ³	17 ³	47 ⁶	10 ³
<i>Empetrum hermaphroditum</i>	1 ²	3 ²	48 ⁶	3 ²
<i>Nardus stricta</i>	-	4 ²	8 ³	100 ⁸
<i>Homogyne alpina</i>	15 ²	49 ³	57 ³	84 ³
<i>Ligusticum mutellina</i>	22 ²	50 ³	32 ²	77 ³
<i>Oreogonium montanum</i>	7 ²	30 ²	13 ²	67 ³
<i>Anthoxanthum alpinum</i>	23 ²	19 ²	17 ²	64 ³
<i>Potentilla aurea</i>	13 ²	34 ²	21 ²	61 ³
<i>Luzula *rubella</i>	6 ²	6 ²	28 ²	41 ²
<i>Luzula sudetica</i>	7 ²	1 ³	1 ²	25 ²
<i>Deschampsia cespitosa</i>	-	1 ²	4 ³	21 ²
<i>Phleum rhaeticum</i>	-	2 ²	1 ²	19 ²
<i>Carex nigra</i>	-	2 ²	1 ²	15 ²

Appendix 1. Comparison of the floristic composition of the classes *Carici repertiv-*
Kobresietea (A), *Caricea curvulae* (B), *Louiseuria-facmetea* (C) and *Nardus-*
rich communitities (D) in the Western Carpathians.
 (Other taxa with frequency higher than 1% arranged in alphabetical order from the data set of 1760 relevés
 and 799 taxa)

Column	A	B	C	D
Number of relevés	339	926	345	150
Average species number	42	22	20	20
<i>Acetosa arifolia</i>	-	+ ²	5 ³	S ²
<i>Agronitis alpina</i>	1 ¹ ³	-	1 ²	-
<i>Achillea millefolium</i> agg.	4 ²	1 ²	3 ²	7 ²
<i>Alectoria americana</i> (E_o)	4 ²	1 ²	1 ²	-
<i>Alectoria ochroleuca</i> (E_o)	21 ²	2 ³	-	-
<i>Alchemilla</i> sp. div.	15 ²	2 ²	10 ³	2 ²
<i>Androsace chamaejasme</i>	12 ³	+ ²	1 ²	2 ²
<i>Anemone narcissiflora</i>	27 ²	7 ²	6 ²	-
<i>Antennaria carpatica</i>	24 ²	2 ²	-	-
<i>Antennaria dioica</i>	-	2 ²	3 ²	-
<i>Anthelia jucunda</i> (E_o)	5 ²	2 ²	1 ²	-
<i>Avenella flexuosa</i>	11 ²	40 ³	65 ³	75 ³
<i>Barbilophozia bycopodioides</i> (E_o)	4 ²	2 ²	2 ²	2 ²
<i>Bartramia ithyphyllo</i> (E_o)	9 ²	+ ¹	-	-
<i>Bazzania tricrenata</i> (E_o)	3 ²	-	1 ²	-
<i>Bellidistrium michelianum</i>	8 ²	+ ³	+ ³	-
<i>Blepharostoma trichophyllum</i> (E_o)	12 ²	1 ¹	1 ²	+ ²
<i>Calamagrostis villosa</i>	4 ³	6 ²	28 ³	22 ²
<i>Callianthemum coriandrifolium</i>	10 ²	+ ²	-	-
<i>Calluna vulgaris</i>	2 ¹	4 ²	12 ²	17 ²
<i>Campanula serrata</i>	1 ²	-	5 ²	-
<i>Campylium stellatum</i> (E_o)	21 ³	+ ³	-	-
<i>Cardamineopsis neglecta</i>	19 ²	1 ²	-	-
<i>Carex arcta</i>	-	-	-	-
<i>Carex firma</i>	30 ²	11 ²	2 ²	3 ¹
<i>Carex sempervirens</i>	9 ⁵	-	1 ²	-
<i>Cephalozia bicuspidata</i> (E_o)	45 ³	51 ⁴	27 ²	53 ³
<i>Ceratium *glandulosum</i>	11 ²	1 ²	+ ²	6 ²
<i>Ceratium cucullatum</i> (E_o)	22 ²	17 ²	6 ²	-
<i>Cetraria ericetorum</i> (E_o)	66 ³	2 ²	4 ²	79 ⁵
<i>Cetraria islandica</i> (E_o)	-	-	-	-
Column	C	D	B	A
<i>Centaria nivalis</i> (E_o)	-	-	8 ³	24 ²
<i>Cladonia arbuscula</i> (E_o)	-	-	20 ³	20 ³
<i>Cladonia bellidiflora</i> (E_o)	-	-	13 ³	14 ²
<i>Cladonia coccifera</i> (E_o)	-	-	33 ³	33 ³
<i>Cladonia cornicina</i> (E_o)	-	-	17 ³	4 ²
<i>Cladonia furcata</i> (E_o)	-	-	1 ²	4 ²
<i>Cladonia gracilis</i> (E_o)	-	-	6 ²	3 ²
<i>Cladonia chlorophyllaea</i> (E_o)	-	-	6 ²	3 ²
<i>Cladonia leptophylla</i> (E_o)	-	-	1 ²	1 ²
<i>Cladonia macrophyllodes</i> (E_o)	-	-	1 ²	1 ²
<i>Cladonia mitis</i> (E_o)	-	-	5 ²	1 ²
<i>Cladonia pyxidata</i> (E_o)	-	-	11 ²	26 ²
<i>Cladonia rangiferina</i> (E_o)	-	-	29 ²	21 ²
<i>Cladonia squamosa</i> (E_o)	-	-	31 ³	1 ²
<i>Cladonia stellaris</i> (E_o)	-	-	5 ³	1 ²
<i>Cladonia unciaria</i> (E_o)	-	-	3 ²	1 ²
<i>Coeloglossum viride</i>	-	-	8 ²	-
<i>Dianthus barbatus</i>	-	-	10 ²	+ ²
<i>Dicranum fuscescens</i> (E_o)	-	-	5 ²	1 ²
<i>Dicranum scoparium</i> (E_o)	-	-	1 ²	1 ²
<i>Dicranum spadicium</i> (E_o)	-	-	10 ²	26 ³
<i>Diplophyllum taxifolium</i> (E_o)	-	-	7 ³	5 ²
<i>Districhium capillaceum</i> (E_o)	-	-	3 ²	-
<i>Ditrichum flexicaule</i> (E_o)	-	-	19 ²	14 ²
<i>Doronicum sibiricum</i>	-	-	48 ³	33 ³
<i>Drennanocladius uncinatus</i> (E_o)	-	-	7 ²	2 ²
<i>Dryas octopetala</i>	-	-	4 ²	4 ²
<i>Erigeron uniflorus</i>	-	-	4 ²	-
<i>Euphrasia tanaceti</i>	-	-	14 ²	6 ²
<i>Festuca sapina</i>	-	-	74 ³	74 ⁴
<i>Festuca picturata</i>	-	-	13 ³	7 ²
<i>Gaodium antispodion</i>	-	-	32 ²	1 ²
<i>Genniana asclepiadea</i>	-	-	4 ²	3 ²
<i>Genniana frigida</i>	-	-	10 ²	3 ³
<i>Genista huascensis</i>	-	-	6 ²	1 ²
<i>Genista pannicaria</i>	-	-	9 ²	14 ²
<i>Geum reptans</i>	-	-	10 ³	4 ²
<i>Hedysarum heterophyllum</i>	-	-	11 ⁴	-
<i>Rhodax alpestris</i>	-	-	9 ³	+ ²
<i>Heterocladium dimorphum</i> (E_o)	-	-	4 ¹	1 ¹
<i>Hieracium alpicola</i>	-	-	4 ²	1 ²

Column	A	B	C	D	A	B	C	D
<i>Hypoxis selago</i>	43 ²	22 ²	29 ²	-	Potentilla crantzii	1 ²	+ ³	-
<i>Hylocomium splendens</i> (E _o)	45 ⁴	10 ²	21 ³	1 ¹	Pholidium ciliare (E _o)	12 ²	7 ²	19 ²
<i>Hypericum maculatum</i>	0 ⁰	1 ²	10 ²	14 ²	<i>Pulsatilla alba</i>	42 ³	3 ²	-
<i>Hypnum capressiforme</i> (E _o)	10 ³	+ ⁵	13 ²	5 ²	<i>Pyrola carpatica</i>	42 ³	+ ²	-
<i>Trommsdorffia uniflora</i>	1 ²	6 ²	23 ²	-	<i>Racomitrium canescens</i> (E _b)	9 ²	2 ²	1 ²
<i>Juniperus *alpina</i>	3 ²	4 ²	13 ²	13 ²	<i>Racomitrium heterostichum</i> (E _b)	9 ²	2 ²	1 ²
<i>Lecidea granulosa</i> (E _o)	3 ²	1 ²	1 ²	1 ²	<i>Racomitrium lanuginosum</i> (E _o)	21 ³	9 ¹	4 ²
<i>Leiomodon pseudotaraxaci</i>	15 ²	2 ²	-	-	<i>Racomitrium suecicum</i> (E _o)	1 ²	3 ²	+ ¹
<i>Lepidoma demissum</i> (E _o)	1 ²	5 ²	2 ²	-	<i>Ranunculus alpestris</i>	22 ⁵	+ ³	-
<i>Lewiscarthus albiloba</i>	1 ²	1 ²	1 ²	7 ²	<i>Ranunculus brevirostris</i>	22 ³	1 ²	-
<i>Leucanthemopsis alpina</i>	16 ²	26 ²	3 ¹	1 ¹	<i>Ranunculus pseudomontanus</i>	21 ²	26 ²	6 ²
<i>Lophozia collaris</i> (E _o)	3 ²	3 ²	1 ²	-	<i>Rhytidium rugosum</i> (E _b)	26 ²	1 ²	-
<i>Lophozia ventricosa</i> (E _o)	1 ¹	2 ²	1 ²	-	<i>Rhytidiodelphus squarrosum</i> (E _b)	3 ²	+ ²	10 ²
<i>Luzula alpinopilosa</i>	45 ²	48 ²	3 ²	19 ²	<i>Rhytidiodelphus triquetrus</i> (E _b)	27 ³	2 ²	8 ³
<i>Luzula sylvatica</i>	-	-	7 ²	5 ²	<i>Salix alpina</i>	4 ³	+ ²	-
<i>Melanopyrum syriacum</i>	-	-	7 ²	3 ²	<i>Salix herbacea</i>	17 ²	17 ⁵	9 ³
<i>Mnium thomsonii</i> (E _o)	16 ³	-	-	-	<i>Salix reticulata</i>	19 ⁵	-	1 ³
<i>Myosoton alpestris</i>	21 ²	1 ²	-	-	<i>Salix retusa</i>	9 ³	3 ⁶	2 ⁴
<i>Nardia scalaris</i> (E _o)	3 ²	2 ³	1 ³	-	<i>Salix silesiaca</i>	4 ²	+ ²	4 ²
<i>Omalotheca supina</i>	1 ²	5 ²	+ ¹	6 ²	<i>Saxifraga aizoides</i>	16 ²	3 ²	-
<i>Oxyria digyna</i>	12 ²	1 ³	-	-	<i>Saxifrage androsacea</i>	15 ²	+ ³	-
<i>Paraleucobryum enerve</i> (E _o)	6 ²	4 ²	1 ³	-	<i>Saxifraga bryoides</i>	19 ²	1 ²	-
<i>Panmasia palustris</i>	13 ²	1 ²	2 ²	2 ²	<i>Saxifraga hieracifolia</i>	17 ²	1 ²	-
<i>Peltigera aphraea</i> (E _o)	9 ²	1 ²	2 ²	1 ¹	<i>Saxifraga oppositifolia</i>	23 ³	+ ²	-
<i>Phyteuma orbiculare</i>	27 ²	1 ³	3 ²	3 ²	<i>Saxifraga renata</i>	12 ²	+ ²	-
<i>Picea abies</i>	2 ¹	1 ²	8 ³	-	<i>Saxifraga wahlenbergii</i>	14 ²	+ ⁴	-
<i>Pinus mugo</i>	3 ¹	2 ²	18 ²	10 ²	<i>Sedum alpestre</i>	11 ²	+ ¹	2 ²
<i>Plagiochila porelloides</i> (E _o)	9 ²	+ ³	-	-	<i>Selaginella selaginoides</i>	22 ²	2 ²	4 ²
<i>Pleurozium schreberi</i> (E _o)	19 ³	17 ²	7 ³	27 ³	<i>Sempervivum montanum</i>	3 ²	6 ²	4 ²
<i>Poa alpina</i>	24 ²	2 ²	2 ²	8 ³	<i>Sempervivum tectorum</i>	5 ²	20 ³	6 ²
<i>Poa taxifolia</i>	18 ²	5 ²	-	-	<i>Senecio abrotanifolius</i>	5 ²	4 ²	-
<i>Polygonatum multiflorum</i> (E _o)	23 ²	6 ²	1 ¹	1 ²	<i>Senecio camniolicus</i>	9 ²	9 ²	1 ¹
<i>Polygonatum multiflorum</i> (E _o)	24 ²	1 ²	1 ²	1 ³	<i>Sesleria albicans</i>	11 ²	+ ²	1 ²
<i>Polygonatum multiflorum</i> (E _o)	4 ²	5 ²	2 ²	1 ⁵	<i>Soldanella carpatica</i>	62 ²	35 ²	30 ²
<i>Polytrichum commune</i> (E _o)	45 ³	34 ³	14 ⁵	1 ²	<i>Solidago *minuta</i>	16 ²	12 ²	17 ²
<i>Polytrichum formosum</i> (E _o)	1 ²	4 ³	19 ³	19 ³	<i>Solstoma croceum</i> (E _b)	11 ²	3 ²	-
<i>Polytrichum juniperinum</i> (E _o)	3 ²	9 ³	10 ³	11 ³	<i>Sorbus aucuparia</i>	1 ²	13 ²	-
<i>Polytrichum piliferum</i> (E _o)	22 ²	29 ³	10 ³	5 ³	<i>Sphagnum</i> sp. div. (E _b)	6 ³	6 ³	21 ⁵
<i>Polytrichum sexangulare</i> (E _o)	1 ²	3 ³	3 ³	+ ³	<i>Sphenolobus minutus</i> (E _b)	2 ¹	2 ²	-
<i>Polytrichum strictum</i> (E _o)	5 ²	8 ³	9 ³	-	<i>Sweetia *alpestris</i>	25 ³	2 ²	-
					<i>Taraxacum</i> sp. div.	4 ²	2 ²	3 ³
					<i>Thamnochortus subuliformis</i> (E _b)	6 ²	5 ²	-
					<i>Thamnochortus vernicularis</i> (E _b)	32 ²	20 ³	1 ²
					<i>Thymus alpestris</i>	4 ²	3 ²	5 ²
					<i>Tortella tortuosa</i> (E _b)	45 ²	1 ²	2 ²
					<i>Trifolium pratense</i>	8 ²	1 ²	-
					<i>Valeriana riparia</i>	6 ³	1 ²	1 ²
					<i>Veratrum lobelianum</i>	2 ²	2 ²	15 ²
					<i>Viola biflora</i>	2 ²	1 ²	2 ²
					<i>Viola *sudetica</i>	5 ²	-	-