

**Syntaxonomy and nomenclature of the communities
of the orders *Calamagrostietalia villosae*
and *Adenostyletalia* in Slovakia**

JÁN KLIMENT¹, IVAN JAROLÍMEK², JOZEF ŠIBÍK² & MILAN VALACHOVIČ²

5 ¹Botanical Garden of Comenius University, SK-038 15 Blatnica, Slovak Republic, e-mail:
kliment@rec.uniba.sk

²Institute of Botany, Slovak Academy of Sciences, Dúbravská cesta 14, SK-845 23
Bratislava, Slovak Republic, e-mail: ivan.jarolimek@savba.sk, jozef.sibik@savba.sk,
milan.valachovic@savba.sk

- 10 KLIMENT J., JAROLÍMEK I., ŠIBÍK J. & VALACHOVIČ M. (2004): Syntaxonomy
and nomenclature of the communities of the orders
Calamagrostietalia villosae and *Adenostyletalia* in Slovakia. – Thaiszia – J. Bot. 14: xx-yy. – ISSN 1210-0420.
- 15 Abstract: The most important results of the syntaxonomical revision
of the communities of the orders *Calamagrostietalia villosae*
and *Adenostyletalia alliariae* from the territory of Slovakia are
presented: description of the new associations (*Geranio robertiani-*
Delphinietum elati) and subassociations (*Festuco picturatae-*
Calamagrostietum villosae typicum and *crepidetosum conyzifoliae*,
Vaccinio myrtilli-Calamagrostietum villosae inops and
avenuletosum versicoloris), re-evaluation of the syntaxonomical
position of the associations (*Petasito-Senecietum nemorensis*
HADAČ et al. 1969, *Doronicetum austriaci* HADAČ et al. 1969,
Senecio-Adenostyletum alliariae HADAČ et al. 1969), resulting in the
description of the new subassociations (*Petasito kablikianii-*
Senecietum nemorensis doronicetosum austriaci and *crepidetosum*
mollis), and in the change of the position of the alliance *Delphinion*
elati HADAČ ex HADAČ et al. 1969 into the suballiance of the alliance
Adenostylion alliariae. Authors suppose the new names for the
associations *Chaerophyllum cicutariae* KRAJINA 1933 (*Bryo*
pseudotriquetri-Chaerophyllum hirsuti), *Calamagrostietum variae*
SILLINGER 1932 (*Geranio sylvatici-Calamagrostietum variae*) and
Calamagrostietum variae carpaticum SILLINGER 1933 (*Convallario*
majalis-Calamagrostietum variae). The subassociations *Athyrietum*
alpestris HADAČ 1956 *typicum* W. MATUSZKIEWICZ et A. MATUSZKIEWICZ
1975 and *A. a. deschampsietosum flexuosae* W. MATUSZKIEWICZ

et A. MATUSZKIEWICZ 1975 are ordered within the valid association name *Adenostylo alliariae-Athyrietum alpestris* (ZLATNÍK 1928) JENÍK 1961. Authors also call attention to the illegitimate typification of some names of associations and they supplement description of the less investigated communities.

Keywords: syntaxonomical revision, new syntaxa, mountain tall-herb vegetation, *Mulgedio-Aconitetea*, West Carpathians.

Introduction

The class *Mulgedio-Aconitetea* is represented by three orders in the Slovak part of the Carpathians. The order *Calamagrostietalia villosae* includes tall-herb floriferous subalpine meadows, the order *Adenostyletalia* covers subalpine communities of broad-leaved herbs and ferns, and the order *Petasito-Chaerophylletalia* involves natural tall-herb nitrophilous communities at banks of rivers and streams in the submontane up to supramontane belt.

Yet were published results of partial syntheses and syntaxonomical revisions of the order *Calamagrostietalia villosae* (incl. *Adenostylium alliariae*) from the wider territory of the Tatry Mts. (ŠEPPER, ŠEFFEROVÁ & DÚBRAVCOVÁ 1989), of the order *Petasito-Chaerophylletalia* from Slovakia (KLIMENT & JAROLIMEK 2002; JAROLIMEK, KLIMENT & VALACHOVIČ 2002), and the alliance *Calamagrostion arundinaceae* from Slovak part of the Carpathians (KLIMENT & JAROLIMEK 2004). This work arisen from complex processing of phytocoenological relevés of the class *Mulgedio-Aconitetea* from Slovakia and resumes the most important unpublished results of their syntaxonomical revision.

Material and methods

Results of comparison of 1216 phytocoenological relevés of the class *Mulgedio-Aconitetea* from the Slovak part of the Carpathians are presented. They are partially compared with relevés from similar communities from surrounding mountain ranges (Sudeten, the East Alps). All relevés were obtained by methods of the Zürich-Montpeliér school (BRAUN-BLANQUET 1928, 1964); authors used different scales of cover: 7-degree Braun-Blanquet's scale, their modified 9-degree version (BARKMAN et al. 1964), also 10-degree Domin's scale or 11-degree Domin-Hadač's scale (cf. KLIKA 1948; HADAČ et al. 1969). To obtain better comparability of relevés, before numerical classification relevés were transformed into the ordinal 9-degree scale (VAN DEN MAAREL 1979). Subspecies and more narrowly delimited species by some authors were included into the nearest higher or broadly defined taxa. Taxa determined at tribe level and mosses (excepting equal presence in data set) were excluded. For numerical classification were used program NCLAS from the package of programs SYNTAX 5 (PODANI 1993). β -flexible method ($\beta = -0.25$) and Jaccard's, Ružička's and Wishart's coefficients and Euclidean distance were used for processing of data. Obtained hypotheses were evaluated on the basis of comparison of the phytocoenological tables, processed by program FYTOPACK (JAROLIMEK & SCHLOSSER 1997).

Nomenclature of taxa follows Checklist of non-vascular and vascular plants of Slovakia (MARHOLD & HINDÁK 1998); exceptions are added by author's abbreviation. In the tables subspecies (without the name of species) are marked by asterisk (*). Frequencies of taxa in % (99 = 100%) are supplemented by average value of cover (upper index). Headings to the columns content: abbreviated citation of the source (in the course of unpublished data only name of the author/authors of relevé), number of relevés and their localization on the level of the orographic units follows the map of Databank of fauna of Slovakia in the scale 1: 500 000.

The names of syntaxa in the text are completed by the author's citation when they are used for the first time only.

Diagnostic taxa of the class *Mulgedio-Aconitetea* and lower syntaxa are indicated following synoptic table (KLIMENT, JAROLÍMEK & ŠIBÍK ined.), treating for prepared book Plant communities of Slovakia 4; its shorted form is in the table 6. Other taxa are ordered in harmony with recent relevant publications. In the Tables 1–5 the names of syntaxa are abbreviated as follows: aa *Adenosty whole alliariae*, ac *Acerenion pseudoplatani*, ai *Alnion incanae*, ca *Calamagrostion arundinaceae*, cl *Calthion*, cr *Calamagrostion variae*, Cv *Calamagrostietalia villosae*, cv *Calamagrostion villosae*, de *Delphinienion elati*, EA *Epilobietea angustifolii*, ES *Elyno-Seslerietea*, fc *Festucion carpatica*, fp *Festucion pictae*, Fs *Fagetalia sylvaticae*, fs *Fagion sylvaticae*, fv *Festucion versicoloris*, JT *Juncetea trifidi*, jt *Juncion trifidi*, lv *Loiseleurio-Vaccinion*, MU *Mulgedio-Aconitetea*, Ns *Nardetalia strictae*, Pc *Petasito-Chaerophylletalia*, QF *Querco-Fagetea*, Sc *Seslerietalia coerulae*, SH *Salicetea herbaceae*, tf *Trisetion fuscum*, VP *Vaccinio-Piceetea*. Diagnostic taxa of communities are bolded in the tables.

In the description of communities were used following abbreviations: baz.: bazionym, C = characteristic taxon (in the tables), comb. nov. = new combination, art. = article of the International code of phytocoenological nomenclature (ICPN; WEBER, MORAVEC & THEURILLAT 2000), D = differential taxon (in the tables), dom. = dominant species, incl. = inclusive, DGT = diagnostic group of taxa, ined. = ineditus (not published data), nom. corr. = corrected name, nom. ined. = invalidly published (not published) name, OFN = original form of name, p. p. = pro parte (partially), prov. = provisional (name), pseu. = pseudonym, subdom. = subdominant species, syn. = synonym, syntax. syn. = syntaxonomical synonym, transgr., t = transgressive taxon (t – used in the tables), r. = relevé.

Descriptions of communities well-known from literature include only the basic data. More information is given in characteristics of less known and newly described syntaxa.

Results and discussion

Calamagrostion villosae PAWŁOWSKI, SOKOŁOWSKI et WALLISCH 1928

130 **Festuco picturatae-Calamagrostietum villosae** PAWŁOWSKI in PAWŁOWSKI,
SOKOŁOWSKI et WALLISCH 1928 nom. corr. hoc loco

(Tab. 1, columns 1a, 1b, 1)

OFN: *Calamagrostis villosa*-*Festuca picta*-Ass. PAWŁOWSKI in PAWŁOWSKI,
SOKOŁOWSKI et WALLISCH 1928

135 **Syn.:** *Calamagrostidetum villosae* SZAFAER, PAWŁOWSKI et KULCZYŃSKI 1923 (art. 31),
Calamagrostidetum tetricum SZAFAER, PAWŁOWSKI et KULCZYŃSKI 1927 (art. 2b, 34a),
Calamagrostidetum villosae tetricum PAWŁOWSKI in PAWŁOWSKI, SOKOŁOWSKI
et WALLISCH 1928 (art. 3a, 34a), *Calamagrostidetum villosae tetricum* KRAJINA 1933
(art. 31), *Calamagrostidetum villosae altherbosum* SILLINGER 1933 (art. 34a),

140 *Calamagrostidetum villosae carpaticum* WALAS 1933 (art. 34a)

Non: *Calamagrostidetum villosae* SCHMID 1923

Differential taxa: *Festuca picturata*, *Acetosa arifolia*, *Adenostyles alliariae*,
*Bistorta major*¹, *Gentiana punctata*¹, *Veratrum album* subsp. *lobelianum*¹

145 **Constant taxa:** *Calamagrostis villosa* (dom.), *Anthoxanthum alpinum*, *Avenella*
flexuosa, *Homogyne alpina*, *Ligusticum mutellina*, *Oreogonium montanum*,
Potentilla aurea, *Soldanella carpatica*, *Solidago virgaurea* subsp. *alpestris*,
Vaccinium myrtillus

¹ against the association *Vaccinio myrtilli-Calamagrostietum villosae*

150 **Nomenclatural type:** PAWŁOWSKI, SOKOŁOWSKI & WALLISCH 1928, Tab. VIII, r. 9,
lectotypus hoc loco

155 *Festuco-Calamagrostietum* is the most spread community of the alliance
Calamagrostion villosae. It prefers moist rocky shoots, bottoms of glacial cirques,
stabilized screes, debris cones, and also free places among stands of dwarf pine
in subalpine and alpine belts in the Nízke, Západné and Vysoké Tatry Mts. It
thrives also at open habitats, exposed to wind. With average number of species
(25) it belongs to the group of floristically rich communities in the granite
bedrock; the number of taxa in individual relevés varies in large range [11–49
(65)]. Mosses and lichens cover less than 10%, they are often entirely absent.

160 Detailed description of the association was brought in several published and
unpublished works (PAWŁOWSKI, SOKOŁOWSKI & WALLISCH 1928; BRAUN-BLANQUET
1930; KRAJINA 1933; SILLINGER 1933; HADÁČ 1956; ŠOMŠÁK et al. 1981; UNAR,
UNAROVÁ & ŠMARDA 1984, 1985; ŠEFFEROVÁ 1984; DÚBRAVCOVÁ et al. 1990; MIADOK
1995, etc.). Differences in the floristic composition and ecological valence of
stands resulted to the distinguishing of following subassociations:

**Festuco picturatae-Calamagrostietum villosae typicum subass. nov. hoc
loco**

(Tab. 1, column 1a)

170 **Nomenclatural type:** identical with the type of name of the association

The community doubles floristically poorer stands (20 taxa in average) on

granite bedrock.

175 **Festuco picturatae-Calamagrostietum villosae crepidetosum conyzifoliae**
Hrabovcová ex Kliment et al. subass. nov. hoc loco
(Tab. 1, column 1b)

Syn: *Calamagrostietum villosae crepidetosum conyzifoliae* Hrabovcová 1976
nom. ined. p. p. (art. 1)

180 **Differential taxa:** *Crepis conyzifolia*, *Geranium sylvaticum*, *Hylotelephium argutum*, *Hypericum maculatum*, *Myosotis scorpioides* agg., *Phyteuma spicatum*, *Ranunculus platanifolius*, *Valeriana tripteris*

Nomenclatural type: Pawłowski, Sokołowski & Wallisch 1928, Tab. VIII, r. 13, holotypus

185 The subassociation is represented by floristically richer stands with average number of taxa 33 [22–49 (69)]. It grows in sunny habitats on mylonites at deeper and moister soils. The stands are usually protected from the wind by rocky walls and by stands of dwarf pine. Against to the typical subassociation this is differentiated by varicoloured flowering herbs demanding more nutritive and moist soils. Occurrence of the species *Carex aterrima*, *Taraxacum alpinum* and *Trisetum fuscum* indicates the relation of these stands to the communities of the alliance *Trisetion fusci*.

195 **Note 1:** Pawłowski, Sokołowski & Wallisch (1928: 248) selected the species *Festuca picta* Kit. ex Schult. for the name of the association, which was described in 1814. This name of the species is younger homonym of the name *F. picta* F. Gmelin 1792, hence Pils (1980: 93) replaced it by the name *Festuca picturata* G. Pils. This replacement constrained the correction of the name of association.

200 **Vaccinio myrtilli-Calamagrostietum villosae** SILLINGER 1933
(Tab. 1, columns 2a, 2b, 2)

205 **Incl.:** *Calamagrostidetum villosae arundinacetosum* Br.-Bl. 1930;
Calamagrostidetum villosae vaccinietosum myrtilli Br.-Bl. 1930; *Myrtleto-*
Calamagrostidetum villosae carpaticum pinetosum mugo Krajina 1933, facies
with *Juniperus nana* p. p.

Characteristic taxa: *Avenula versicolor* (transgr.)

Differential taxa: *Vaccinium vitis-idaea*

210 **Constant taxa:** *Calamagrostis villosa* (dom.), *Vaccinium myrtillus* (subdom.),
Avenella flexuosa, *Carex sempervirens* subsp. *silicicola* Holub ined., *Homogyne alpina*, *Ligisticum mutellina*, *Luzula luzuloides*, *Oreogezum montanum*, *Potentilla aurea*

Nomenclatural type: Sillinger 1933: 276, r. 1, lectotypus hoc loco

215 **Vaccinio-Calamagrostietum** is usually closed two-layered community of dwarf shrubs and grasses. Its floristic composition, synecology and synchorology were only scarcely mentioned in present publications (Sillinger 1933, Krajina 1933, Miadok 1995); manuscripts held slightly more data (Treskoňová 1972, Kremlová

1974, DÚBRAVCOVÁ et al. 1976, ALTMANNOVÁ 1983). The community is formed beside
220 the dominant *Calamagrostis villosa* by smaller graminoids (*Anthoxanthum alpinum*,
Avenella flexuosa, *Carex *silicicola*, *Luzula luzuloides*) and small shrubs
(*Vaccinium myrtillus*, *V. vitis-idaea*). The most of bright flowering plants are
concentrated to the ground layer. Cover of cryptogams varies between 0–10%, it
moister habitats it reaches 50%. Stands of the community are recently known
225 from the gaps in dwarf pine stands or above these stands on slopes up to 45°
and on the bottoms of glacial cirques in the subalpine to alpine belts in the Nízke,
Západné and Vysoké Tatry Mts., approximately between 1530–2000 m a. s. l.
Within the association two floristically, synmorphologically, syncologically and
syngenetically different subassociations were distinguished:

230 ***Vaccinio myrtilli-Calamagrostietum villosae inops* subass. nov. hoc loco**

(Tab. 1, column 2a)

Nomenclatural type: MIADOK 1995: 38–39, r. 3, holotypus

235 The subassociation joints floristically poorer (14–23, in average 17 taxa),
monotonous small shrubby-grass stands with subdominant to dominant
presence of the species *Vaccinium myrtillus*, which prefers elevated parts of the
terrain with accumulated raw humus. Prevailingly secondary stands arise after
the clearing dwarf pine (KREMLOVÁ 1974, MIADOK 1995). Higher moisture of
240 habitats induces higher frequency of mosses.

Vaccinio myrtilli-Calamagrostietosum villosae avenuletosum versicoloris
subass. nov. hoc loco

(Tab. 1, column 2b)

245 **Differential taxa:** *Agrostis pyrenaica*, *Anthoxanthum alpinum*, *Avenula*
versicolor, *Campanula alpina*, *C. tatrae*, *Carex sempervirens* subsp. *silicicola*,
Festuca supina, *Hieracium alpinum* agg., *Juncus trifidus*, *Pulsatilla scherfelii*,
Ranunculus pseudomontanus, *Sempervivum carpathicum* WETTST. ex PRODAN
subsp. *carpathicum*, *Trommsdorffia uniflora*, *Viola lutea* subsp. *sudetica*

250 **Nomenclatural type:** identical with the type of association

255 Physiognomically conspicuous flower-rich and species richer plant community
with average number of taxa 24 (18–33). Its centre of distribution lies in the lower
part of the alpine belt on sunny south oriented (SW–SE) slopes. Floristic
composition is similar to the association *Agrostio rupestris-Caricetum*
sempervirentis SILLINGER 1933 (alliance *Juncion trifidi* KRAJINA 1933), which is
reflected also in the list of differential taxa.

260 **Note 2:** Abundant relevé data of the communities with *Calamagrostis villosa*
from the Tatry Mts. region contain relatively high number of transition stands
between the associations *Festuco pictae-Calamagrostietum villosae*
and *Vaccinio myrtilli-Calamagrostietum villosae*. This fact complicates the
mutual differentiation both associations and also unambiguous classification of
these stands.

Trisetion fusti KRAJINA 1933

(Tab. 2, columns 1–5)

Syn.: *Deschampsion caespitosae* BORZA 1934 (art. 29c, 31), *Phleo alpini-*

Deschampsion caespitosae (BORZA 1934) ŠT. CSÚRÓS, GERGELY & M. CSÚRÓS

(art. 29c)

Syntax. syn.: *Aconition firmi* KRAJINA 1933

Nomenclatural type: *Rhodiolo-Deschampsietum caespitosae* KRAJINA 1933,

lectotypus

275 The alliance *Trisetion fusti* associates mosaic, chiono- and hygrophilous communities of tall grasses, dicotyledonous herbs and small shrubs. They are among the species richest high mountains communities on granite bedrock. They occur at bottoms of the small terrain depressions with alluvial fine soil and humus near the high mountain torrents and tarns; water flows among rocks deeper below the soil surface. Configuration of terrain causes thick snow layer.

280 BORZA (1934: 34) described the association *Deschampsietum caespitosae transsilvanicum* from the Retezat Mts. (Romania), and created the new alliance *Deschampsion caespitosae*. ŠT. CSÚRÓS, GERGELY & M. CSÚRÓS (1985: 144) within the order *Adenostyletalia* delimited the alliance *Phleo alpini-Deschampsion* all. nov.; based on synonyms at pages 144 a 155 (*Deschampsion caespitosae* BORZA 34, non HORVATIĆ 30) we deduced, that in fact this is the new name for the alliance *Deschampsion caespitosae* BORZA 1934. Since the both authors qualified as the synonym of both alliances (*Deschampsion caespitosae*, *Phleo alpini-Deschampsion caespitosae*) the name *Trisetion fusti* KRAJINA 1933, in accordance to the art. 29c ICPN these names were illegitimate superfluous names for the alliance *Trisetion fusti*.

285 KRAJINA (1933) described from the moisture habitats near the small springs and stream banks in subalpine belt of the Vysoké Tatry Mts. the alliance *Aconition firmi*. The alliance covered two associations *Aconitetum firmi* and *Chaerophylletum cicutariae*, with diagnostic taxa *Chaerophyllum hirsutum*, *Chrysosplenium alternifolium*, *Geum rivale*, *Myosotis laxiflora*, *Stellaria nemorum* and *Bryum weigelii*. Syntaxonomical revision proved relevancy of both associations to the alliance *Trisetion fusti* and confirmed the shift of the lectotype, i. e. the association *Aconitetum firmi* (art. 20) by authors ŠEFFER, ŠEFFEROVÁ & DÚBRAVCOVÁ (1989) into this alliance. Consequently, the alliance *Aconition firmi* KRAJINA 1933 we evaluate as syntaxonomical synonym of the alliance *Trisetion fusti*.

290 *Deschampsio caespitosae-Salicetum helveticae* (KRAJINA 1933) DÚBRAVCOVÁ et ŠEFFER 1992

(Tab. 2, column 1)

Baz.: *Salicetum lapponum tetricum* KRAJINA 1933 (art. 34a)

Syn.: *Salicetum helveticae* (KRAJINA 1933) DÚBRAVCOVÁ et HRABOVCOVÁ in MUCINA et MAGLOCKÝ 1985 (art. 2b, 31)

300 **Non:** *Salicetum helveticae* BRAUN-BLANQUET, PALLMANN et BACH 1954

Nomenclatural type: KRAJINA 1933, Tab. 32, r. 6, lectotypus hoc loco

DÚBRAVCOVÁ & ŠEFFER (1992) published the new name for the association *Salicetum lapporum tetricum* KRAJINA 1933. They typified it by the relevé from HADAČ (1956, Tab. 24, r. 63). In accordance to the art 39 ICPN their typification is illegitimate (cf. THEURILLAT & MORAVEC 1995: 366).

315

Rhodiolo-Deschampsietum caespitosae KRAJINA 1933

(Tab. 2, column 3)

Syn.: *Deschampsietum caespitosae tetricum* HADAČ 1956 (art. 34a)

320

Syntax. syn.: *Trisetetum fuscum* KRAJINA 1933

Nomenclatural type: KRAJINA 1933, Tab. 30, r. 4, lectotypus hoc loco

325

Syntaxonomical revision confirmed the Šeffer's opinion (ŠEFFER 1984: 24), that the association *Trisetetum fuscum* KRAJINA 1933 is only part of variability of the association *Rhodiolo-Deschampsietum caespitosae*; by floristic composition it is in harmony with the subassociation *swertietosum alpestris* ŠEFFER 1991. Following the art. 20 ICPN the association *Rhodiolo-Deschampsietum* represents lectotype of the alliance *Trisetion fuscum*.

330

***Phleo rhaetici-Deschampsietum caespitosae* (KRAJINA 1933) COLDEA 1983**

nom. corr. hoc loco

(Tab. 2, column 3)

Baz.: *Deschampsietum caespitosae* KRAJINA 1933 (art. 31)

335

Syn.: *Aconito firmi-Deschampsietum alpicola* (KRAJINA 1933) HADAČ in MUCINA et MAGLOCKÝ 1985 (art. 2b), *Phleo alpini-Deschampsietum caespitosae* (KRAJINA 1933) COLDEA 1983 (art. 43)

Nomenclatural type (lectotypus): KRAJINA 1933, Tab. 33, r. 4 (UNAR, UNAROVÁ & ŠMARDA 1985: 55)

340

The community with prevalence of species *Deschampsia cespitosa* from the subalpine belt of the Vysoké Tatry Mts. described KRAJINA (1933) with illegitimate name *Deschampsietum caespitosae*. In the latter phytocoenological works from the territory of Slovakia the relevés of this community are the most frequently ordered into the invalidly published new name *Aconito firmi-Deschampsietum alpicola* (KRAJINA 1933) HADAČ in MUCINA et MAGLOCKÝ 1985. However COLDEA (1983) published the name *Deschampsietum caespitosae* KRAJINA 1933 earlier as a basionym of the name „*Phleo alpini-Deschampsietum caespitosae* nom. nov.“. By recent knowledge the name giving species is *Phleum rhaeticum* (HUMPHRIES) RAUSCHERT (syn.: *Ph. alpinum* auct. non L.). Consequently the correction of the name was necessary (art. 43).

345

COLDEA (l. c.) typified the name *Phleo alpini-Deschampsietum caespitosae* by own relevé from the Rodna Mts. in Romania, which is in contradiction with the art. 39a ICPN. The lectotype of this name *Deschampsietum caespitosae* KRAJINA 1933, selected from the original data was published by UNAR, UNAROVÁ & ŠMARDA (1985). This lectotype does not contain the species *Aconitum firmum*, hence also the art. 3f ICPN relates to the Hadač's name *Aconito firmi-Deschampsietum alpicola*.

350

Aconitetum firmi Sokołowski in Pawłowski, Sokołowski et Wallisch 1928

(Tab. 2, column 4)

360 **Syntax. syn.:** *Delphinio oxysepali-Aconitetum firmi* Br.-Bl. 1930

Phantom name: *Aconitetum firmi* PAWŁOWSKI et STECKI 1927 (KLIKA 1948, 1955),
Aconitetum firmi KRAJINA 1933 (MUCINA & MAGLOCKÝ 1985, ŠEFFER, ŠEFFEROVÁ
& DÚBRAVCOVÁ 1989, MIADOK 1995)

365 **Nomenclatural type:** PAWŁOWSKI, SOKOŁOWSKI & WALLISCH 1928, Tab. X, r. 4,
lectotypus hoc loco

In the phytocoenological works and surveys from the territory of Slovakia (ŠEFFER 1984, MUCINA & MAGLOCKÝ 1985, ŠEFFER, ŠEFFEROVÁ & DÚBRAVCOVÁ 1989, MIADOK 1995) the stands dominated by the species *Aconitum firmum* on granite bedrock ordered into the association *Aconitetum firmi* KRAJINA 1933. However, some years earlier the association *Aconitetum firmi* (within the alliance *Adenostylium alliariae*) was validly described by SOKOŁOWSKI (in PAWŁOWSKI, SOKOŁOWSKI & WALLISCH 1928). It was pointed in KRAJINA (1933: 900–901) by authors citations PAWŁOWSKI et STECKI 114, 1927; SOKOŁOWSKI (PAWŁOWSKI, SOKOŁOWSKI, WALLISCH Zesp. rośl. w Tatr. VII, 231, 1928) behind the name of the association, and also by bibliographic reference in the text: „Diese Assoziation wurde ursprünglich von Pawłowski und Stecki (l. c.) auf Kalkunterlage beschrieben, später von Sokołowski aus dem Morskie-Oko-Tal auf Granit.“. PAWŁOWSKI & STECKI (1927: 114–116, Tab. 9) evaluated coenoses with *Aconitum firmum* as vegetation unit without rang (*Aconitum firmum*-Hochstaudenflur; art. 3c), which is quite different in its content from *Aconitum firmum* dominated stands on granite bedrock. KRAJINA (1933: 901) for the first time identified both associations *Aconitetum firmi* and *Aconitum firmum-Delphinium oxysepalum* Br.-Bl. 1930. Results of syntaxonomical revision confirmed this solution.

385 **Bryo pseudotriquetri-Chaerophylletum hirsuti** (KRAJINA 1933) nom. nov. hoc
loco

(Tab. 2, column 5)

Baz.: *Chaerophylletum cicutariae* KRAJINA 1933: 908 (art. 34a)

390 **Non:** *Chaerophylletum cicutariae* ZLATNÍK 1928

Characteristic taxa: *Bryum pseudotriquetrum* (transgr.), *Chiloscyphus polyanthos*, *Plagiomnium medium*

Differential taxa: *Doronicum austriacum*, *Galium anisophyllum*, *Poa pratensis*,
Brachythecium rivulare, *Bryum weigelii*, *Marchantia polymorpha*, *Philonotis seriata*, *Rhizomnium punctatum*

Constant taxa: *Chaerophyllum hirsutum* [syn.: *Ch. cicutaria* VILL.] (dom.),
Aconitum firmum, *Alchemilla* sp. div., *Bistorta major*, *Caltha palustris* subsp.
laeta, *Deschampsia cespitosa*, *Hypericum maculatum*, *Ligusticum mutellina*,
Luzula alpinopilosa subsp. *obscura*, *Potentilla aurea*, *Senecio subalpinus*,
Stellaria nemorum

400 **Nomenclatural type:** KRAJINA 1933, Tab. 27, r. 5, lectotypus

405 Yet relatively recondite and probably neglected community described by
KRAJINA (1933) from slightly sloped (prevailingly up to 5°) banks and alluvium of
the stream Mlynica and its tributaries in the Vysoké Tatry Mts., 1500–1650 m a.
s. l. It was scarcely seen also in the Roháčska dolina Valley in the Západné Tatry
Mts. (KOMÁRKOVÁ 1964). Island-formed stands grow on slightly acid to slightly
410 alkaline shallow soils. Within the alliance *Trisetion fuscii* these stands stay
nearest to the stands of associations *Aconitetum firmi* and *Phleo rhaetici-*
Deschampsietum caespitosae. They can form mutual transitions. Regularly
occurrence of hygrophilous herbs and mosses proves that they are at the same
time close to the spring communities. High number of alliance and class, less of
order species (Tab. 2) justifies their ordination into the tall-herb communities of
415 the *Mulgedio-Aconitetea*.

Calamagrostion variae SILLINGER 1932 emend. HADAČ et al. 1969

(Tab. 3, columns 1, 2)

420 **Phantom name:** *Calamagrostion variae* SILLINGER 1929 (GRABHERR, GREIMLER
& MUCINA 1993), *Calamagrostion variae* SILLINGER ex HADAČ 1962 (MUCINA
& MAGLOCKÝ 1984), *Calamagrostion variae* SILLINGER ex HADAČ 1963 (MUCINA
& MAGLOCKÝ 1985)

425 The alliance *Calamagrostion variae* consists of floristically and
physiognomically very various calciphilous tall-grass communities in steep,
sunny, and wind protected slopes and on the bottom of sheer avalanche
channels on limestone-dolomite bedrock in the mountain to subalpine belts. In
430 the scarped scarred slopes they form vegetation complex with open stands of
relict pine woods. The communities occupy also talus cones in the terminals of
rocky channels in the belt of deciduous or mixed forests; in higher elevation they
occupy also glades in dwarf pine stands. Common occurrence of several
subthermophilous species indicates similar ecology than the alliance
Calamagrostion arundinaceae.

435 SILLINGER (1932: 19) described the alliance *Calamagrostion variae* as substrate
analogue of the alliance *Calamagrostion villosae* on the carbonate bedrock. This
fact result also from communities ordered into the alliance in the original
diagnosis (*Calamagrostietum variae*, *Festucetum carpaticae*, *Adenostyletum*
alliariae calcicolum). In the original range the alliance partially included also later
described alliances *Festucion carpaticae* and *Delphinion elati*. HADAČ et al. (1969:
440 134) considerably reduced range of the alliance by ordering only the association
Calamagrostietum variae carpaticum SILLINGER 1933 into the alliance. However,
automatic lectotype following the article 20 ICPN is the association
Calamagrostietum variae SILLINGER 1932. Its original diagnosis represents only
phytocoenological relevé.

445 Localization of stands, properties of the biotope, and contact phytocoenoses
determined high presence of calciphilous forest species and elements of divet
phytocoenoses. This facts resulted into the ordering of the alliance into the order
Seslerietalia tatrae HADAČ 1962 (HADAČ 1962, HADAČ et al. 1969) or order
Seslerietalia coeruleae Br.-BL. in Br.-BL. et JENNY 1926 (MUCINA & MAGLOCKÝ 1984,
450 1985; GRABHERR, GREIMLER & MUCINA 1993).

Convallario majalis-Calamagrostietum variae (SILLINGER 1933) nom. nov. hoc loco

(Tab. 3, column 1)

Baz.: Calamagrostidetum variae carpaticum SILLINGER 1933: 167 (art. 34a)

455 **Syn.: Carlino-Calamagrostietum variae** (SILLINGER 1933) HADAČ in MUCINA et MAGLOCKÝ 1985 (art. 2b)

Incl.: Carlino-Calamagrostietum variae (SILLINGER 1933) HADAČ in MUCINA et MAGLOCKÝ 1985 *convallarietosum* MUCINA et MAGLOCKÝ ms. (art. 1, 2b)

460 **Characteristic taxa:** *Adenophora liliifolia*, *Anthericum ramosum*, *Bupleurum falcatum*, *Knautia slovaca*, *Polygonatum odoratum*, *Pulsatilla slavica*

Differential taxa: *Achillea stricta*, *Brachypodium pinnatum*, *Campanula rapunculoides*, *Convallaria majalis*, *Galium schultesii*, *Securigera varia*

Constant taxa: *Calamagrostis varia* (dom.), *Carduus glaucinus*, *Cirsium erisithales*, *Laserpitium latifolium*, *Rubus saxatilis*

465 **Nomenclatural type:** SILLINGER 1933: 168, r. 4, lectotypus

The community with *Calamagrostis varia* is open, medium species rich, with average number of taxa 39 (22–53), and with regular presence of several mesophilic to subxerophilic herbs. Its detailed characteristics published SILLINGER (1933). *Convallario-Calamagrostietum* represents the original natural relict community on the steep, south (SW–SE) oriented limestone-dolomite slopes in the forest belt. It forms vegetation complex together with stands of relict pines woods or cover primary forest-free areas on talus cones. It was found in mountain belt (approx. 760–1220 m a. s. l.) in the mountain ranges Muránska planina, Slovenský raj, Veľká Fatra, Chočské vrchy and Nízke Tatry.

470 Occurrence of species *Carduus glaucinus*, *Erysimum witmannii*, *Phyteuma orbiculare*, *Pulsatilla slavica*, *Scabiosa lucida*, *Thesium alpinum* indicates the close relation of the association to the communities of the order *Seslerietalia coerulae*. It caused the original classification of the association within the alliance *Seslerion coerulae* BR.-BL. in BR.-BL. et JENNY 1926 (SILLINGER 1933). On the other hand, more or less regular presence of elements of mountain tall-herb communities (*Astrantia major*, *Cirsium erisithales*, *Cyanus mollis*, *Laserpitium latifolium*, *Pimpinella major* subsp. *rhodochlamys*, *Pyrethrum clusii* etc.) joints this association with the class *Mulgedio-Aconitetea*. Within this class it occupies marginal state towards the boundary of the class *Elyno-Seslerietea* BR.-BL. 1948.

480 **Note 3:** HADAČ (in MUCINA & MAGLOCKÝ 1985) for the association *Calamagrostietum variae carpaticum* SILLINGER 1933 supposed the new name *Carlino (acaulis)-Calamagrostietum variae*. *Carlina acaulis* is typical for the stands on rocky slopes (on the talus cones it almost absent), and in the same time it acts as constant companion of the association *Geranio sylvatici-Calamagrostietum variae*. Based on these facts, for the name of the association were selected differential species, used before in the invalidly published name of the subassociation *convallarietosum* (see the synonyms of the association).

495

**Geranio sylvatici-Calamagrostietum variae (SILLINGER 1932) nom. nov. hoc
loco**

(Tab. 3, column 2)

500 **Baz.: Calamagrostidetum variae** SILLINGER 1932: 4 (art. 31)

Syn.: Calamagrostidetum variae altherbosum SILLINGER 1933 (art. 2b, 34a)

Pseu.: Calamagrostetum variae carpaticum sensu HADAČ et al. 1969 non
SILLINGER 1933

505 **Incl.: Carduo glauci-Caricetum tatrorum calamagrostietum variae** UNAR in UNAR,
UNAROVÁ et ŠMARDA 1985; *Anemono narcissiflorae-Laserpitietum latifolii*
poetosum nemoralis KLIMENT 1995, variant with *Calamagrostis varia*

510 **Differential taxa:** *Achillea millefolium* subsp. *alpestris*, *Astrantia major*,
Campanula elliptica, *C. serrata*, *Carex sempervirens* subsp. *tatrorum*, *Crepis*
mollis, *Geranium sylvaticum*, *Helianthemum grandiflorum*, *Knautia maxima*,
Leontodon hispidus, *Linum extraaxillare*, *Phleum hirsutum*, *Primula elatior*,
Ranunculus nemorosus, *Silene vulgaris*

515 **Constant taxa:** *Calamagrostis varia* (dom.), *Carlina acaulis*, *Cirsium erisithales*,
Galium anisophyllum, *Laserpitium latifolium*, *Leucanthemum vulgare* agg., *Lotus*
corniculatus, *Phyteuma orbiculare*, *Pimpinella major* subsp. *rhodochlamys*,
Scabiosa lucida, *Sesleria albicans*

Nomenclatural type: SILLINGER 1932: 5, r. 6, holotypus

520 Opened to almost close, species rich tall grass community (39–62, 48 taxa in average) is shafted by numerous conspicuous flowering herbs. Cover of moss etage does not exceed 30%. Island-formed stands occupy sunny, from wind protected, south (SW–SE) oriented slopes at the foothill of rocky walls near the timber line, in glades within dwarf pine stands, and also in the ends of steep avalanche glens with centre of distribution in the supramontane belt (approx. 1250–1550 m a. s. l.). They are closely related to the tall-grass communities of 525 the alliance *Calamagrostion arundinaceae* in floristic composition and syncology. They also form transition stands with the similar association *Festucetum carpaticae*. Recently the community is known from the limestone-dolomite peripheries of central mountain ranges of the West Carpathians (Veľká Fatra, Chočské vrchy, Nízke, Západné and Belianske Tatry).

530 The first relevé of the community published SILLINGER (1932) with name „nivové *Calamagrostidetum variae*“ from the rocky channel in the west slopes of Krakova hoľa Mt., 1540 m a. s. l., within the description of stands with *Festuca carpatica* in the Nízke Tatry Mts.; in the later work (SILLINGER 1933) he mentioned it with the name *Calamagrostidetum variae altherbosum*. Phytocoenoses 535 dominated by the species *Calamagrostis varia* from protected habitats in the supramontane to subalpine belt were later ordered to the association *Calamagrostietum variae carpaticum* SILLINGER 1933 (HADAČ et al. 1969), or they were classified in the level of variant or subassociation within other associations (UNAR, UNAROVÁ & ŠMARDA 1984, 1985; KLIMENT 1995). Syntaxonomical revision 540 approved strong differences in their floristic composition against the association *Calamagrostietum variae carpaticum*, which were appointed by UNAR, UNAROVÁ & ŠMARDA (1985) and validity of delimitation of distinctive association.

Festucion carpaticae BĚLOHLÁVKOVÁ et FIŠEROVÁ 1989

(Tab. 3, column 3)

- 545 The community of chiono- and hygrophilous, neutro- to slightly alkaliphilous species dominated by *Festuca carpatica* or scarcely by *Calamagrostis varia*. Stands are species very rich. They prefer protected habitats in steep erosion channels and avalanche glens with thick and long term snow cover in the supramontane to subalpine belt.
- 550 The alliance in Slovakia includes only one association *Festucetum carpaticae* DOMIN 1925. It was from the time of its description gradually ordered into various higher syntaxa. PAWŁOWSKI, SOKOŁOWSKI & WALLISCH (1928), SILLINGER (1933), PAWŁOWSKI (1935, 1956), KLIKA (1948, 1955) and HADAČ (1956) classified this association within the alliance *Calamagrostion villosae*, BRAUN-BLANQUET (1930) and KLIKA (1931) within the alliance *Seslerion coerulæ*, SILLINGER (1932) and ŠOMŠÁK et al. (1980) within the alliance *Calamagrostion variae*, HADAČ et al. (1969), UNAR, UNAROVÁ & ŠMARDA (1984, 1985), MUCINA & MAGLOCKÝ (1985) and DÚBRAVCOVÁ & HAJDÚK (1986) within the alliance *Seslerion tatrae*. This disunity of opinions on classification of the stands of *Festuca carpatica* was stimulus for the re-evaluation of all relevant data including description of the new alliance *Festucion carpaticae* (BĚLOHLÁVKOVÁ & FIŠEROVÁ 1989). In spite of close floristic and synecological relations between the associations *Geranio sylvatici-Calamagrostietum variae* and *Festucetum carpaticae* (cf. SILLINGER 1932), syntaxonomical revision supports their classification into different higher syntaxa and confirms the delimitation of the alliance *Festucion carpaticae*.

Adenostylium alliariae BR.-BL. 1926

- 570 **Syn.:** *Adenostylium* LUQUET 1926 (art. 33), *Adenostylium alliariae* PAWŁOWSKI, SOKOŁOWSKI et WALLISCH 1928 p. p. maj. (art. 31), *Adenostylium alliariae* BR.-BL. 1930 p. p. (art. 31), *Alno-Adenostylium* BR.-BL. 1948 (art. 29); *Dryopterido-Athyriion distentifolii* HOLUB in HOLUB et al. 1967 (art. 3b)
Syntax. syn.: *Dryopterido-Athyriion distentifolii* (HOLUB ex SÝKORA et ŠTURSA 1973) JENÍK, BUREŠ et BUREŠOVÁ 1980
Incl.: *Eu-Adenostylium* KLIKA in KLIKA et HADAČ 1944; *Eu-Adenostylenion alliariae* SÝKORA et ŠTURSA 1973; *Dryopterido-Athyrenion distentifolii* HOLUB ex SÝKORA et ŠTURSA 1973

- 580 The alliance *Adenostylium alliariae* associates mosaic, flowering species rich communities of tall broad-leaved herbs and ferns optimally developed at moist habitats in subalpine belt.

- 585 Communities of ferns from the regions of the Sudeten and the West Carpathians have delimited in present works and surveys into the special suballiance *Dryopterido-Athyrenion distentifolii* (SÝKORA & ŠTURSA 1973; MUCINA & MAGLOCKÝ 1984, 1985), or the alliance *Dryopterido-Athyriion distentifolii* (JENÍK, BUREŠ & BUREŠOVÁ 1980; MORAVEC et al. 1983, 1995; Kočí 2001a). Syntaxonomical revision of the relevés from the Slovak part of the West Carpathians showed that

the associations *Ranunculo platanifolii-Adenostyletum alliariae* and *Adenostylo-Athyrietum alpestris* are linked by many transitions and their classification within separated alliances is not well-founded.

HADAČ et al. (1969) delimited tall herb, chiono- and hygrophilous, hemisciophilous, species rich communities on the limestone bedrock in the subalpine belt of the Belianske Tatry Mts., within the order *Seslerietalia tatrae* HADAČ 1962, class *Aconito-Cardaminetea* HADAČ 1956, into the particular alliance *Delphinion elati*. ŠEFFER, ŠEFFEROVÁ & DÚBRAVCOVÁ (1989) ordered the alliance similarly into the order *Seslerietalia tatrae* (within the class *Mulgedio-Aconitetea*). MUCINA & MAGLOCKÝ (1984, 1985) the alliance rearranged into the order *Adenostyletalia alliariae*. Other authors, e. g. KARNER & MUCINA (1993) regard the *Delphinion elati* as the syntaxonomical synonym of the alliance *Adenostylion alliariae*. Syntaxonomical revision of tall herb communities from Slovakia indicated that both mentioned alliances are lack of characteristic species (excepting of the species *Delphinium elatum*), and they are separated only by several acidophilic or (hemi) calciphilous differential species (Tab. 6). Consequently we prefer to classify the former alliance *Delphinion elati* as the well differentiated suballiance within the alliance *Adenostylion*.

***Adenostylenion alliariae* KLIKA in KLIKA et HADAČ 1944**

OFN: *Eu-Adenostylon* (Krajina 1933) KLIKA in KLIKA et HADAČ 1944

Differential taxa: *Gentiana punctata*, *Luzula alpinopilosa* subsp. *obscura*,
610 *Oreogenum montanum*

The suballiance *Adenostylenion alliariae* combines species rich and flowery communities of tall broad-leaved herbs and ferns in moist habitats in subalpine (to alpine) belt on granite bedrock.

615 The name *Eu-Adenostylon* is recondite in the phytocoenological literature. KLIKA & HADAČ (1944) used it in delimitation of the suballiance *Aconitenion firmi*, SÝKORA & ŠTURSA (1973) in delimitation of the suballiance *Dryopterido-Athyrenion distentifolii*. In this work the suballiance *Adenostylenion alliariae* relates to the range of the suballiance *Eu-Adenostylon* in KLIKA & HADAČ (1944).

620 *Ranunculo platanifolii-Adenostyletum alliariae* (KRAJINA 1933) DÚBRAVCOVÁ et HADAČ ex Koci 2001
(Tab. 4, columns 1a, 1b, 1)

Baz.: *Adenostyletum alliariae tetricum* KRAJINA 1933 (art. 34a)

625 **Syn.:** *Adenostyletum alliariae silicicolum* SILLINGER 1933 (art. 34a), *Ranunculo platanifolii-Adenostyletum alliariae* (KRAJINA 1933) DÚBRAVCOVÁ et HADAČ in MUCINA et MAGLOCKÝ 1985 (art. 2b)

Non: *Adenostyletum alliariae* BR.-BL. 1930

Characteristic taxa: *Poa granitica*

630 **Differential taxa:** *Aconitum firmum*¹, *Doronicum austriacum* (subdom.)¹, *Festuca picturata*, *Phleum rhaeticum*¹, *Potentilla aurea*¹, *Silene dioica*¹, *Viola biflora*¹

Constant taxa: *Adenostyles alliariae* (dom.), *Acetosa arifolia*, *Calamagrostis villosa*, *Gentiana punctata*, *Homogyne alpina*, *Ligusticum mutellina*, *Luzula*

635 *alpinopilosa* subsp. *obscura*, *Oreogeum montanum*, *Ranunculus platanifolius*,
Soldanella carpatica, *Veratrum album* subsp. *lobelianum*

1 against the association *Adenostylo-Athyrietum alpestris*

640 Species medium rich (22 species per relevé in average), closed and physiognomically patchwork community with prevalence of tall broad-leaved herbs forms islets in surrounding vegetation. Cover of mosses in relation to the conditions of habitat ranges 0 to 90%. Mosaic stands occupy moist sites near the banks of brooks, terrain depressions, talus cones below the wet rocky walls with seeping water, all protected from direct influence of wind. They occur optimally developed in the upper part of subalpine to lower part of alpine belts in the Nízke, 645 Západné and Vysoké Tatry Mts., on the granite bedrock; on the mylonites they form transitions to the association *Aconito firmi-Adenostyletum alliariae*. KRAJINA (1933) and ŠEPPER & ŠEFFEROVÁ (1989) published detailed information on structure, synecology and inside classification of the community.

650 ŠEPPER & ŠEFFEROVÁ (l. c.) distinguished within the association four subassociations: *Ranunculo platanifolii-Adenostyletum ranunculetosum pseudomontani* Šeffer et Šeffarová 1989, *R.-A. deschampsietosum alpicolaे* ŠEPPER et ŠEFFEROVÁ 1989, *R.-A. milietosum alpicolaе* ŠEPPER et ŠEFFEROVÁ 1989 and *R.-A. athyrietosum distentifolii* (HADAČ 1956) ŠEPPER et ŠEFFEROVÁ 1989. Actual syntaxonomical revision of the communities of the order *Adenostyletalia* from the 655 Slovak part of the West Carpathians confirmed only two of them, namely *R.-A. milietosum alpicolaе* (Tab. 2, column 1a) and *R.-A. ranunculetosum pseudomontani* (Tab. 2, column 1b). Subassociation *R.-A. deschampsietosum alpicolaе* forms well differentiated variant of the first subassociation (differential species: *Deschampsia cespitosa*, *Primula elatior*, *Brachythecium reflexum*, 660 *Pseudoleskea incurvata*), and stands with *Athyrium distentifolium* were classified as the distinctive association.

Adenostylo alliariae-Athyrietum alpestris (ZLATNÍK 1928) JENÍK 1961

(Tab. 4, columns 2a, 2b, 2)

665 **Baz.: Athyrietum alpestris** ZLATNÍK 1928 (art. 31)

Syn.: Athyrietum alpestris tetricum HADAČ 1956 (art. 34a), **Acetoso alpestris-Athyrietum alpestris** (HADAČ 1956) HADAČ in MUCINA et MAGLOCKÝ 1985 (art. 2b)

Incl.: Ranunculo platanifolii-Adenostyletum alliariae athyrietosum distentifolii (HADAČ 1956) ŠEPPER et ŠEFFEROVÁ 1989

670 **Phantom name:** *Athyrietum alpestris* HADAČ 1956 emend. W. MATUSZKIEWICZ et A. MATUSZKIEWICZ 1960 (MATUSZKIEWICZ W. & MATUSZKIEWICZ A. 1975, JENÍK, BUREŠ & BUREŠOVÁ 1980)

Non: *Athyrietum alpestris* SCHMID 1923

Characteristic taxa: *Athyrium distentifolium* (transgr., dom.)

675 **Differential taxa:** *Oxalis acetosella*, *Rubus idaeus*

Constant taxa: *Adenostyles alliariae* (subdom.), *Acetosa arifolia*, *Calamagrostis villosa*, *Homogyne alpina*, *Milium effusum*, *Veratrum album* subsp. *lobelianum*

Nomenclatural type: JENÍK 1961, Tab. 9, r. 5, neotypus

680 Monotonous physiognomy of closed and floristically relatively poor (6–26, in
average 14 taxa) community of *Athyrium distentifolium* is diversified by several
medium tall to tall broad-leaved herbs. Dense crisscross of dried leaves of the
dominant fern inhibit development of the moss layer and its cover reaches up
maximally to 10%. The community occupies shaded stabilised screes with long
685 term snow and from wind protected by rock walls and by stands of dwarf pine. It
occurs mainly in the lower part of subalpine belt of the Západné and Vysoké
Tatry Mts. and Krivánska Malá Fatra Mts., in the slightly inclined or steep slopes
of various orientation; sparsely it overlaps to the elevation over 1800 m a. s. l.
690 Excepting of several papers (HADAČ 1956; UNAR, UNAROVÁ & ŠMARDA 1984, 1985;
ŠEFFER & ŠEFFEROVÁ 1989) the community is known mostly from manuscripts.

Comparison of relevés of the West Carpathians communities with relevés
from mountains of high Sudeten (JENÍK 1961, SÝKORA & ŠTURSA 1973,
MATUSZKIEWICZ W. & MATUSZKIEWICZ A. 1975, Kočí 2001b) using numerical
classification showed that analysed stands from the West Carpathians form two
695 subassociations:

***Adenostylo alliariae-Athyrietum alpestris typicum* (W. MATUSZKIEWICZ
et A. MATUSZKIEWICZ 1975) comb. nov. hoc loco**

(Tab. 4, column 2a)

700 **Baz.: Athyrietum alpestris typicum** W. MATUSZKIEWICZ et A. MATUSZKIEWICZ 1975: 93

Nomenclatural type: identical with the name of association

Several layered small stands form the transition to the association *Ranunculo
platanifolii-Adenostyletum alliariae*. Transition position of the subassociation is
705 manifested by presence of numerous constant companions of this association
(*Gentiana punctata*, *Luzula alpinopilosa*, *Milium effusum*, *Oreogeum montanum*,
Ranunculus platanifolius, *Soldanella carpatica*).

***Adenostylo alliariae-Athyrietum alpestris avenelletosum flexuosae*
(W. MATUSZKIEWICZ et A. MATUSZKIEWICZ 1975) comb. nov. hoc loco**

(Tab. 4, column 2b)

Baz.: Athyrietum alpestris deschampsietosum W. MATUSZKIEWICZ
et A. MATUSZKIEWICZ 1975: 93

715 **Differential species:** *Oxalis acetosella*, *Dryopteris dilatata*, *Gentiana
asclepiadea*, *Rubus idaeus*, *Vaccinium myrtillus*

Nomenclatural type: MATUSZKIEWICZ W. & MATUSZKIEWICZ A. 1975, Tab. X, r. 11,
lectotypus

720 Stands of the subassociation occupy moist stabilised granite screes in gaps of
the dwarf pine stands or in their margins; secondary communities developing for
the long time after the cutting of the fern spruce forests have more or less
identical species composition.

725 **Note 4:** HADAČ (1956) described his community from the Vysoké Tatry Mts. as the
new association *Athyrietum alpestris tetricum*. Comparison of relevés from high
mountains of the West Carpathians with the association described from the High

Sudeten Mts. showed high similarity of both communities and resulted to including of Carpathian phytocoenoses into the association *Adenostylo-Athyrietum alpestris* described from the High Sudeten Mts. Compared communities differ only by rare presence of some territorially limited taxa, e. g. 730 *Aconitum firmum*, *Luzula alpinopilosa* subsp. *obscura*, *Soldanella carpatica* and *Aconitum hians* Rchb. W. MATUSZKIEWICZ & A. MATUSZKIEWICZ (1975) and Kočí (2001a) also identify both these communities.

735 **Note 5:** Names of the subassociations *Athyrietum alpestris typicum* and *Athyrietum alpestris deschampsietosum* published W. MATUSZKIEWICZ & A. MATUSZKIEWICZ (1975) with the illegitimate association name *Athyrietum alpestris* HADAČ 1955 [recte: 1956] (art. 31). Following the art. 30 ICPN, this fact compelled publishing of the new combinations.

***Delphinienion elati* (HADAČ ex HADAČ et al. 1969) stat. nov. hoc loco**

740 **Baz.:** *Delphinion elati* HADAČ ex HADAČ et al. 1969: 138

Syn.: *Delphinion elati* HADAČ 1962 (art. 8)

Characteristic taxa: *Delphinium elatum*

Differential taxa: *Chaerophyllum hirsutum*, *Cortusa matthioli*, *Epilobium alpestre*, *Galeobdolon luteum* s. l., *Luzula sylvatica*, *Stellaria nemorum*

745 **Nomenclatural type:** *Petasito-Senecietum nemorensis* HADAČ et al. 1969, lectotypus

***Aconito firmi-Adenostyletum alliariae* DOMIN 1930 nom. invers. propos.**

(Tab. 4, column 3)

750 **OFN:** *Adenostyleto-Aconitetum* DOMIN 1930

Syn.: *Adenostyletum alliariae* BR.-BL. 1930 (art. 31), *Adenostyletum alliariae calcicolum* SILLINGER 1933 (art. 34a), *Delphinietum oxysepali* HADAČ et al. 1969 prov. (art. 3b), *Aconiteo-Adenostyletum* DOMIN 1925 (nom. nud.)

Non: *Senecioni-Adenostyletum alliariae* HADAČ et al. 1969

755 **Characteristic taxa:** *Saxifraga rotundifolia*

Differential taxa: *Alchemilla* spec. div., *Deschampsia cespitosa*, *Ligusticum mutellina*¹, *Rhodiola rosea*¹

760 **Constant taxa:** *Adenostyles alliariae* (dom.), *Acetosa arifolia*, *Aconitum firmum*, *Chaerophyllum hirsutum*, *Geranium sylvaticum*, *Hypericum maculatum*, *Primula elatior*, *Senecio nemorensis* agg., *Stellaria nemorum*, *Viola biflora*

¹ against to the communities of the suballiance *Delphinienion elati*

Nomenclatural type: DOMIN 1930: 179, holotypus

765 Varicoloured, physiognomically conspicuous, close, medium species rich to rich (19–60, average 30 taxa) community form islets stands, alternatively dominated by several tall broad-leaved herbs: *Adenostyles alliariae*, *Aconitum firmum*, *Cicerbita alpina* and *Doronicum austriacum*. Among them stems of *Senecio nemorensis* agg. and stalks of several grasses grow. Cover of mosses varies depend on habitat circumstances in range 0–80%.

770 The community occupies stabilised soiled screes on the bottom of shallow depressions below the rocky walls and boulder bottoms of occasional brooks. It

rarely occurs also in natural rocky glades within the crest spruce forests in the supramontane belt. The community prevailingly grows on carbonates, less on melaphyres and other basic bedrocks.

775 The community engaged attention of DOMIN (1930), who published relevé of stand form alpine belt of the Belianske Tatry Mts. with the name *Adenostyleto-Aconitetum*; this valid name was overlooked by later authors. Beside the Belianske Tatry Mts. the community was found also in other mountains (Veporské vrchy, Krivánska Malá Fatra, Veľká Fatra, Nízke, Západné 780 and Vysoké Tatry), and published with different names by BRAUN-BLANQUET (1930), SILLINGER (1933), HADAČ et al. (1969), DÚBRAVCOVÁ & HAJDÚK 1986 and KLIMENT, BERNÁTOVÁ & OBUCH (1994); from some of mentioned mountains only unpublished relevés exist.

785 ***Petasito kablikiani-Senecietum nemorensis* HADAČ et al. 1969**

(Tab. 4, columns 4a, 4b, 4)

Syn.: *Petasito kablikiani-Senecietum jacquinianii* (HADAČ et al. 1969) HADAČ 1987
(art. 29)

790 **Syntax. syn.:** *Senecioni-Adenostyletum alliariae* HADAČ et al. 1969,
Doronicetum austriaci HADAČ et al. 1969

Non: *Arunco-Doronicetum austriaci* KORNÁŠ in KORNÁŠ et MEDWECKA-KORNÁŠ 1967

795 **Differential taxa:** *Calamagrostis arundinacea*, *Cardaminopsis halleri*, *Cirsium erisithales*, *Cystopteris fragilis*, *Dactylis glomerata* subsp. *slovenica*, *Hylotelephium argutum*, *Myosotis alpestris*, *Paris quadrifolia*, *Phyteuma spicatum*, *Pimpinella major* subsp. *rhodochlamys*, *Pulmonaria obscura*, *Scrophularia scopolii*, *Trisetum flavescens* subsp. *taticum*, *Eurhynchium praelongum*, *Mnium spinosum*

800 **Constant taxa:** *Senecio nemorensis* agg., *Acetosa arifolia*, *Bistorta major*, *Carduus personata*, *Chaerophyllum hirsutum*, *Chrysosplenium alternifolium*, *Delphinium elatum*, *Epilobium alpestre*, *Galeobdolon luteum* s. l., *Galium schultesii*, *Geranium sylvaticum*, *Geum rivale*, *Hypericum maculatum*, *Polygonatum verticillatum*, *Primula elatior*, *Silene dioica*, *Thalictrum aquilegiifolium*, *Valeriana excelsa* subsp. *sambucifolia*, *Veratrum album* subsp. *lobelianum*, *Viola biflora*

805 **Nomenclatural type:** HADAČ et al. 1969: 143, r. 263, lectotypus

810 Several layered, closed, islets-forming stands are notable by their highness, high number of flowering species and species richness (average 46 taxa per relevé). They occupy more or less stabilized limestone screes on steep, sunny to slightly shadow slopes in upper parts of avalanche glens, infrequently also on natural glades in dwarf pine stands in the subalpine belt in the Belianske Tatry Mts., 1575–1725 m a. s. l. Detail characteristics of the community was published by HADAČ et al. (1969).

815 Authors described from the Dolina Siedmich prameňov Valley in the Belianske Tatry Mts. three closely related associations: *Petasito-Senecietum nemorensis*, *Senecio-Adenostyletum alliariae* and *Doronicetum austriaci*. Syntaxonomical revision of communities of the order *Adenostyletalia* from the territory of Slovakia

proved that all these associations belong to the only one association. Based on constant occurrence of the species *Senecio nemorensis* agg. in all relevés the name *Petasito-Senecietum nemorensis* was retained for this association.

820 According to the actual knowledge above mentioned aggregate taxon is in the tall herb communities in the subalpine belt of the Belianske Tatry Mts. represented almost exclusively by species *Senecio hercynicus*, in the riparian 825 communities dominated by *Petasites kablikianus* also the species *Senecio germanicus* penetrates into high elevations. The variability of originally described communities responds to the subassociations:

***Petasito kablikiani-Senecietum nemorensis doronicetosum austriaci* (HADAČ et al. 1969) comb. nov. et stat. nov. hoc loco**

830 (Tab. 4, column 4a)

Baz.: *Doronicetum austriaci* HADAČ et al. 1969: 150

Differential taxa: *Doronicum austriacum*, *Filipendula ulmaria*, *Milium effusum*, *Ribes petraeum*

Nomenclatural type: HADAČ et al. 1969: 152, r. 235, lectotypus

835 The subassociation covers phytocoenoses dominated by the species *Doronicum austriacum* in natural glades in dwarf pine stands.

***Petasito kablikiani-Senecietum nemorensis crepidetosum mollis* subass.**

840 **nov. hoc loco**

(Tab. 4, column 4b)

Differential taxa: *Crepis mollis*, *Astrantia major*, *Campanula elliptica*, *Daphne mezereum*, *Festuca carpatica*, *Linum extraaxillare*, *Origanum vulgare*, *Petasites kablikianus*, *Pleurospermum austriacum*, *Poa nemoralis*, *Eurhynchium angustirete*

Nomenclatural type: identical with the type of association name

This subassociation in the original delimitation includes beside the association 845 *Petasito-Senecietum nemorensis* also stands of the association *Seneciono-Adenostyletum alliariae*.

Chaerophyllo hirsuti-Cicerbitetum alpinae (KÄSTNER 1938) SÝKORA et HADAČ 1984
(Tab. 4, column 5)

Baz.: *Mulg Dietum alpini montanum* KÄSTNER 1938 (art. 34a)

Incl.: *Adenostylo-Athyrietum alpestris petasitetosum albi* KOPECKÝ et HEJNÝ 1971

Non: *Cicerbitetum alpinae* BOLLETER 1921

Differential taxa: *Cicerbita alpina*, *Athyrium distentifolium*¹, *Leucanthemum rotundifolium*, *Myosotis nemorosa*, *Petasites albus* (subdom.), *Prenanthes purpurea*

860 **Constant taxa:** *Aconitum firmum*, *Adenostyles alliariae* (dom.), *Calamagrostis villosa*, *Chaerophyllum hirsutum* (subdom.), *Doronicum austriacum*, *Galeobdolon montanum*, *Gentiana asclepiadea*, *Geranium sylvaticum*, *Luzula sylvatica*, *Oxalis acetosella*, *Ranunculus platanifolius*, *Senecio subalpinus*, *Stellaria nemorum*,

Viola biflora

865 ¹ against the other communities of the suballiance *Delphinienion elati*

Mostly closed, floristically medium rich (19–26, average 22 taxa), physiognomically conspicuous community dominated by the species *Adenostyles alliariae* with higher cover of tall herbs and ferns (*Aconitum firmum*,
870 *Athyrium distentifolium*, *Chaerophyllum hirsutum*, *Cicerbita alpina*, *Doronicum austriacum*, *Petasites albus*), occupies banks of rapid creeks in mountain and upper mountain belt. It was found in the middle and upper part of the Roháčsky potok Stream in the Západné Tatry Mts., 1180–1550 m a. s. l. (KOPECKÝ 1971). Analyzed stands this author ordered into the subassociation *Adenostylo-Athyrietum alpestris petasitetosum albi* KOPECKÝ et HEJNÝ 1971. Kočí (2001a)
875 regarded this name as syntaxonomical synonym of the association *Chaerophyllo-Cicerbitetum alpinae*, described from the German side of the Krušné hory Mts. In the original diagnoses of both syntaxa the species *Adenostyles alliariae* was not presented, but in the stands from the Západné Tatry it prevails. Comparison of both communities using numerical classification nevertheless showed their close floristic composition. This fact and identical synecology of both communities permitted ordering of the West Carpathian phytocoenoses into the association *Chaerophyllo-Cicerbitetum alpinae*.
880 Comparison with the other West Carpathian communities of the alliance *Adenostyliion alliariae* indicated that the stands from the Roháčsky potok Stream in spite of granite bedrock are markedly similar to the communities of the suballiance *Delphinienion elati* thanks to the sufficient of nutrients in the soil.

Geranio robertiani-Delphinietum elati ass. nov. hoc loco

890 (Tab. 4, columns 6a, 6b, 6, Tab. 5)

Non: *Delphinietum elati* BEGER ex SUTTER 1978

Characteristic taxa: *Delphinium elatum* (transgr., dom.)

Differential taxa: *Acer pseudoplatanus*, *Carex muricata*, *Clinopodium vulgare*,
895 *Fragaria vesca*, *Geranium robertianum*, *Urtica dioica*
Constant taxa: *Arabis alpina*, *Carduus personata*, *Chaerophyllum hirsutum* (subdom.), *Chrysosplenium alternifolium*, *Heracleum sphondylium*, *Senecio nemorensis* agg., *Silene dioica*, *Valeriana excelsa* subsp. *sambucifolia*, *Viola biflora*

Nomenclatural type: Tab. 5, r. 5 holotypus

900 Two or three layered phytocoenoses dominated by the species *Delphinium elatum*. This species together with the subdominant *Chaerophyllum hirsutum* and other tall herbs (*Aconitum variegatum*, *Carduus personata*, *Heracleum sphondylium*, *Senecio nemorensis* agg., *Urtica dioica*, *Valeriana *sambucifolia*)
905 determines the maximal height of stands 180–200 cm. Species *Arabis alpina*, *Chrysosplenium alternifolium*, *Geranium robertianum*, *Silene dioica*, *Stellaria nemorum* and *Viola biflora* form the ground and medium layer. Cover of mosses varies from 10 to 90%. The community is species rich (average 41, min. 24, max. 57 species per relevé).

910 It occupies boulder colluviums, fixed scree cones and gravel alluviums along

the mountain torrents. Inverse character of microclimate permits occurrence of the community in deep valleys and below the rocky walls. Habitats are partially shaded; soils are skeletal, humus-rich and moist.

According to the actual information the association *Geranio-Delphinietum elati* prefers limestone and dolomite regions in the mountain and upper mountain belt in the West Carpathians. Phytocoenological relevés come from glens, ravines and banks of mountain brooks in the Muránska planina Mts. (JAROLÍMEK, KOCHJAROVÁ, TURIS, VALACHOVIČ ined.), Krivánska Malá Fatra Mts. (JAROLÍMEK, KRAJČIOVÁ ined.), Veľká Fatra Mts. (MUCINA ined.), and v Nízke Tatry Mts. (JAROLÍMEK, MAGLOCKÝ, MUCINA ined.).

Different floristic composition and synecology of stands resulted into the classification of two subassociations:

Geranio robertiani-Delphinietum elati orobanchetosum flavae subass. nov.

hoc loco

(Tab. 4, column 6a; Tab. 5, r. 1–5)

Differential taxa: *Orobanche flava*, *Crepis paludosa*, *Impatiens noli-tangere*, *Myosotis nemorosa*, *Petasites albus*, *P. kablikianus*, *Roegneria canina*, *Stachys sylvatica*, *Stellaria nemorum*

Nomenclatural type: Tab. 5, r. 5, holotypus

This floristically poorer community (24–40, average 35 taxa) has higher number of wetland species. Its stands are in close relation to the riparian communities of the alliance *Petasition officinalis* SILLINGER 1933 by their floristic composition and synecology.

Geranio robertiani-Delphinietum elati ranunculetosum platanifolii subass. nov. hoc loco

(Tab. 4, column 6b; Tab. 5, r. 6–10)

Differential species: *Ranunculus platanifolius*, *Cirsium erisithales*, *Galeobdolon montanum*, *Galium schultesii*, *Geranium sylvaticum*, *Luzula luzuloides*, *Polygonatum verticillatum*

Nomenclatural type: Tab. 5, r. 7, holotypus

Stands of the subassociation contain higher number of forest species and species from mountain alluvium vegetation; they have average number of taxa 47 (39–57) per relevé.

Note 6: Beside some rarely occurred species only several constant companions, such as *Chaerophyllum hirsutum*, *Heracleum sphondylium*, *Silene dioica*, *Urtica dioica* and *Viola biflora* are common for both associations *Geranio-Delphinietum elati* and *Delphinietum elati* BEGER ex SUTTER 1978. The last one was described from the Swiss Alps, 1350–1760 m a. s. l. (SUTTER 1978). On the contrary, comparison of both associations uncovered evident differences between them in their floristic composition. West Carpathian phytocoenoses positively differ from the Alpine ones by occurrence of species *Aconitum variegatum*, *Cardamine*

960 *impatiens*, *Carex muricata*, *Chrysosplenium alternifolium*, *Clinopodium vulgare*,
Cystopteris montana, *Digitalis grandiflora*, *Epilobium montanum*, *Fragaria vesca*,
Galium schultesii, *Geranium robertianum*, *Lathyrus vernus*, *Myosotis scorpioides*
965 agg., *Orobanche flava*, *Oxalis acetosella*, *Petasites albus*, *P. kablikianus*,
Pulmonaria obscura, *Roegneria canina* and *Stachys sylvatica*. At the same time
numerous group of taxa differs the Alpine community against the stands
dominated by *Delphinium elatum* in the West Carpathians: *Acetosa arifolia*,
970 *Achillea macrophylla* L., *Aconitum paniculatum* LAM., *A. platanifolium* DEGEN ex
GÄYER, *Adenostyles alliariae*, *A. glabra* (MILL.) DC., *Alchemilla* spec. div.,
Campanula scheuchzeri VILL., *Carex ferruginea* SCOP., *Centaurea nervosa* WILLD.,
Cirsium oleraceum, *Crepis pyrenaica* (L.) GREUTER, *Epilobium alpestre*, *Festuca*
975 *pratensis*, *Geranium sylvaticum*, *Imperatoria ostruthium* L., *Leucanthemum*
adustum (W. D. J. KOCH) GREMLI, *Lilium martagon*, *Phyteuma ovatum* HONCK., *Poa*
hybrida GAUDIN, *Polystichum aculeatum*, *Pulsatilla alpina* (L.) DELARBRE,
Ranunculus serpens SCHRANK, *Rosa pendulina*, *Saxifraga rotundifolia*, *Veratrum*
album, *Veronica latifolia*.

975 *Daphno mezerei-Dryopteridetum filicis-maris* SÝKORA et ŠTURSA 1973

980 Two or three layered, more or less closed and species rich community is typical by dense cover of leaves of the species *Dryopteris filix-mas* with length up to 90 cm. In the shade of leaves of dominant grows numerous smaller herbs. Physiognomy of stands is diversified by medium to tall flowering herbs (985 *Adenostyles alliariae*, *Carduus personata*, *Digitalis grandiflora*, *Gentiana asclepiadea*, *Pyrethrum clusii*, *Senecio hercynicus*, *Silene dioica*, *Valeriana excelsa* subsp. *sambucifolia*). Mosses are presented scarcely on crop out rocks and boulders.

985 The community occupies more or less stabilized scree below the top rock walls, near the tree limit in the supramontane belt. Till now it was known only from the Sudeten Mountains Krkonoše, Jizerské hory, Králický Sněžník and Hrubý Jeseník. SÝKORA & ŠTURSA (1973) described it based on 22 phytocoenological relevés and detailed study of its ecology. In Slovakia the first relevé comes from the Mačacia Mt. (1410 m) on melaphyre bedrock:

990 **Locality:** Nízke Tatry Mts., Mačacia Mt., stabilized scree with soil in the ravine below rock walls, above the forest; 1310 m a. s. l., WNW, 40°, 4 × 6 m, E_i: 95 %, E_d: do 5 %, 48°57'33,5" north latitude, 19°52'35,7" east longitude, August 5, 2003, J. Kliment & P. Turis.

995 E_i: *Dryopteris filix-mas* 4, *Acetosa arifolia* 2a, *Calamagrostis arundinacea* 2a, *Angelica sylvestris* 1, *Digitalis grandiflora* 1, *Galeobdolon luteum* 1, *Senecio hercynicus* 1, *Stellaria nemorum* 1, *Valeriana excelsa* subsp. *sambucifolia* 1, *Anthriscus sylvestris* +, *Bromus benekenii* +, *Campanula rapunculoides* +, *Cardaminopsis halleri* +, *Carduus personata* +, *Cicerbita alpina* +, *Clinopodium vulgare* +, *Daphne mezereum* +, *Epilobium montanum* +, *Fragaria vesca* +, *Galeopsis pubescens* +, *Galium schultesii* +, *Gentiana asclepiadea* +, *Geranium robertianum* +, *G. sylvaticum* +, *Heracleum sphondylium* +, *Hordelymus europaeus* +, *Hylotelephium argutum* +, *Hypericum maculatum* +, *Luzula luzuloides* +, *Melica nutans* +, *Milium effusum* +, *Myosotis sylvatica* +, *Oxalis acetosella* +, *Paris quadrifolia* +, *Poa chaixii* +, *P. nemoralis* +, *Pulmonaria obscura* +, *Pyrethrum clusii* +, *Rosa pendulina* +, *Rubus idaeus* +, *Silene dioica* +, *Solidago virgaurea* subsp.

1005 *minuta* +, *Thlaspi caerulescens* +, *Urtica dioica* +, *Veronica chamaedrys* +, *Adenostyles alliariae* r, *Cardamine impatiens* r, *Carex muricata* r, *Crepis mollis* r, *Polygonatum verticillatum* r, *Scrophularia scopolii* r.

The association *Daphno-Dryopteridetum* was ordered by authors – within the alliance *Adenostylium alliariae* – into the suballiance *Dryopterido-Athyrenion distentifolii*, later into the individual alliance *Dryopterido-Athyriion distentifolii* (MORAVEC et al. 1995, Kočí 2001a, b). Comparison of relevant West Carpathian communities of the order *Adenostyletalia* and original diagnose of the association *Daphno-Dryopteridetum* indicated, that analysed stand from the Nízke Tatry Mts. in spite of dominance of the species *Dryopteris filix-mas* with respect to its floristic composition clearly resembles to the communities of the suballiance *Delphinienion elati*. Within the association it is the closest to the typical variant, which presents optimally developed phytocoenoses in drier and warmer habitats (cf. SÝKORA & ŠURSA 1973, Tab. 1, r. 8–22).

Acknowledgements

1020 Authors are indebted for collaboration in the field and for providing unpublished phytocoenological relevés to S. Alexyová, D. Bernátová, Z. Dúbravcová, E. Holotová, M. Chytrý, J. Kochjarová, I. Krajčiová, P. Kučera, Z. Kyselová, Š. Maglocký, L. Mucina, L. Paclová, J. Školek, R. Šoltés, A. Šoltés and P. Turis. This work was supported by grants of VEGA 1/9149/02 and 1/0045/03.

Sources to tables:

(used abbreviations: BT=the Belianske Tatry Mts.; KMF=the Krivánska Malá Fatra Mts.; MP=the Muránska planina Mts.; NT=the Nízke Tatry Mts.; VF=the Veľká Fatra Mts.; VT=the Vysoké Tatry Mts.; ZT= the Západné Tatry Mts.;

1030 Tab. 1, the alliance *Calamagrostion villosae*

- 1a KOMÁRKOVÁ 1964, Tab. 30 (*Calamagrostietum villosae*), r. 2, ZT; KREMLOVÁ 1974, Tab. inter p. 48–49 (*Calamagrostietum villosae*), r. 3–6, 8, 9, ZT; DÚBRAVCOVÁ 1976, Tab. 10 (*Calamagrostietum villosae*), r. 4, 6–11, ZT; DÚBRAVCOVÁ et al. 1976, Tab. 18 (*Calamagrostietum villosae*), r. 9, 11, 13, ZT; HRABOVCOVÁ 1976, Tab. 9 (*Calamagrostietum villosae*), r. 2, 3, ZT; KRÁLIK 1979, Tab. 11 (*Calamagrostietum villosae*), r. 1, 2, 5, 7, ZT; ÚNAR, ÚNAROVÁ & ŠMARDA 1984, Tab. 31 (*Calamagrostietum villosae*), r. 1, 2, 4, 5, ZT; DÚBRAVCOVÁ et al. 1990, Tab. 20 (*Festuco pictae-Calamagrostietum villosae*), r. 5, 6, ZT; BRAUN-BLANQUET 1930: 20 (*Calamagrostis villosa-Festuca picta*-Ass.), 1 r., VT; KRAJINA 1933, Tab. 29 (*Calamagrostidetum villosae*), r. 5–8, 10, VT; HADAČ 1956, Tab. 17 (*Calamagrostietum villosae*), r. 6, VT; ŠOMŠÁK et al. 1981, Tab. 10 (*Calamagrostietum villosae*), r. 2, 5, VT; ŠEFFEROVÁ 1984, Tab. 3 (*Festuco pictae-Calamagrostietum villosae*), r. 1–13, VT; DÚBRAVCOVÁ et al. 1990, Tab. 20, r. 1–4, 7, 12–15, 17, VT; SILLINGER 1933: 276 (*Calamagrostis villosa-Vaccinium myrtillus*-Ass.), r. 2, 4, 6, NT; TRESKOŇOVÁ 1972, Tab. inter p. 68–69 (*Calamagrostietum villosae altherbosum*), 6 r., NT; ALTMANNOVÁ 1983, Tab. 19 (*Calamagrostietum villosae*), 9 r., NT; DÚBRAVCOVÁ 1983: 32, 1 r. (*Festuco pictae-Calamagrostietum villosae*), NT; FAJMONOVÁ 1987: 1–2, r. 1 (*Festuco pictae-Calamagrostietum villosae*), NT; MIADOK 1995: 37 (*Festuco pictae-Calamagrostietum villosae*), r. 2, 4, NT.

- 1050 **1b** KOMÁRKOVÁ 1964, Tab. 30 (*Calamagrostietum villosae*), r. 3, ZT; KREMLOVÁ 1974, Tab. inter p. 48–49 (*Calamagrostietum villosae*), r. 10, ZT; DÚBRAVCOVÁ 1976, Tab. 10 (*Calamagrostietum villosae*), r. 5, 12–14, ZT; DÚBRAVCOVÁ et al. 1976, Tab. 18 (*Calamagrostietum villosae*), r. 12, 21, 22, ZT; HRABOVCOVÁ 1976, Tab. 9 (*Calamagrostietum villosae*), r. 10, ZT; KRÁLIK 1979, Tab. 11 (*Calamagrostietum villosae*), r. 3, 4, 6, ZT; PAWŁOWSKI, SOKOŁOWSKI & WALLISCH 1928, Tab. VIII, r. 13, 14, VT; KRAJINA 1933, Tab. 29 (*Calamagrostidetum villosae*), r. 1–4, 11, VT; HADAČ 1956, Tab. 17 (*Calamagrostietum villosae*), r. 1–5, 7–10, VT; ŠEFFEROVÁ 1984, Tab. 3 (*Festuco pictae-Calamagrostietum villosae*), r. 14–20, VT; ŠEFFEROVÁ 1984, Tab. 3, r. 21–24 (*Rumici scutati-Calamagrostietum villosae* prov.), VT; DÚBRAVCOVÁ et al. 1990, Tab. 20 (*Festuco pictae-Calamagrostietum villosae*), r. 10, 16, 18, 19, VT; DÚBRAVCOVÁ et al. 1990, Tab. 20, r. 8, 9, BT; ALTMANNOVÁ 1983, Tab. 13 (*Vaccinio myrtilli-Calamagrostietum villosae*), r. 1, NT; DÚBRAVCOVÁ 1983: 33 (*Festuco pictae-Calamagrostietum villosae*), 1 r., NT.
- 1055 **2a** KREMLOVÁ 1974, Tab. inter p. 34–35 (*Calamagrostis villosa-Vaccinium myrtillus-Ass.*), r. 2–4, Tab. inter p. 48–49 (*Calamagrostietum villosae*), r. 1, ZT; DÚBRAVCOVÁ 1976, Tab. 10 (*Calamagrostietum villosae*), r. 1–3, ZT; DÚBRAVCOVÁ et al. 1976, Tab. 18 (*Calamagrostietum villosae*), r. 28, 33, 36, ZT; KRAJINA 1933, Tab. 62 (*Myrtilleto-Calamagrostidetum villosae carpaticum*), r. 2, VT; ŠOMŠÁK et al. 1981, Tab. 10 (*Calamagrostietum villosae*), r. 3, VT; MIADOK 1995: 37 (*Festuco pictae-Calamagrostietum villosae*), r. 3, p. 38–39 (*Vaccinio myrtilli-Calamagrostietum villosae*), r. 1–3, NT.
- 1060 **2b** KOMÁRKOVÁ 1964, Tab. 30 (*Calamagrostidetum villosae*), r. 1, 4, 5; ZT; KREMLOVÁ 1974, Tab. inter p. 34–35 (*Calamagrostis villosa-Vaccinium myrtillus-Ass.*), r. 1, Tab. inter p. 48–49 (*Calamagrostietum villosae*), r. 2, 7, ZT; DÚBRAVCOVÁ et al. 1976, Tab. 18 (*Calamagrostietum villosae*), r. 20, 34, 35, 37, ZT; HRABOVCOVÁ 1976, Tab. 9 (*Calamagrostietum villosae*), r. 4–9, ZT; KRÁLIK 1979, Tab. 11 (*Calamagrostietum villosae*), r. 8, 9, ZT; UNAR, UNAROVÁ & ŠMARDA 1984, Tab. 31 (*Calamagrostietum villosae*), r. 3, ZT; BRAUN-BLANQUET 1930: 21 (*Calamagrostidetum villosae arundinacetosum*), 1 z, VT; KRAJINA 1933, Tab. 29 (*Calamagrostidetum villosae*), r. 9, VT; ŠOMŠÁK et al. 1981, Tab. 10 (*Calamagrostietum villosae*), r. 1, 4, VT; DÚBRAVCOVÁ et al. 1990, Tab. 20 (*Festuco pictae-Calamagrostietum villosae*), r. 11, VT; SILLINGER 1933: 276 (*Calamagrostis villosa-Vaccinium myrtillus-Ass.*), r. 1, 3, 5, NT; TRESKOŇOVÁ 1972: 49 (*Calamagrostis villosa-Vaccinium myrtillus-Ass.*), 7 r., NT; ALTMANNOVÁ 1983, Tab. 13 (*Vaccinio myrtilli-Calamagrostietum villosae*), r. 2–9, NT; DÚBRAVCOVÁ 1983: 32 (*Festuco pictae-Calamagrostietum villosae*), 1 r., NT; FAJMONOVÁ 1987: 2, r. 2 (*Vaccinio myrtilli-Calamagrostietum villosae*), NT; MIADOK 1995: 37 (*Festuco pictae-Calamagrostietum villosae*), r. 1, NT.
- 1065 **3** KOMÁRKOVÁ 1964, Tab. 36, r. 1–3 (*Salicetum lapponum*), ZT; HORÁK 1971, Tab. 2, r. 51, 52 (*Salicetum lapponae-helveticum*), ZT; DÚBRAVCOVÁ & ŠEFFER 1992, Tab. 1 (*Calamagrostio villosae-Salicetum helveticae*), r. 2, 4, 5, 7–9, 11–14, ZT, r. 10, VT.
- 1070 **Tab. 2, the alliance *Trisetion fuscii***
- 1075 **1** KREMLOVÁ 1974, Tab. inter p. 84–85 (*Salicetum lapponum tetricum*), 7 r., ZT; KRAJINA 1933, Tab. 32 (*Salicetum lapponae tetricum*), 6 r., VT; HADAČ 1956, Tab. 24 (*Salicetum lapponum tetricum*), 3 r., VT; DÚBRAVCOVÁ & ŠEFFER 1992, Tab. 1 (*Deschampsia caespitosa-Salicetum helveticae*), r. 15, 16, 18–20, 23, 30, VT.
- 1080 **2** KOMÁRKOVÁ 1964, Tab. 34 (*Deschampsietum caespitosae tetricum*), r. 1, 2, p. 100 (*Rhodiolo-Deschampsietum caespitosae*), 1 r., ZT; KREMLOVÁ 1974, Tab. inter p. 81–82 (*Rhodiolo-Deschampsietum caespitosae*), 4 r., ZT; DÚBRAVCOVÁ et al. 1976, Tab. 20 (*Rhodiolo-Deschampsietum caespitosae*), r. 5–11, Tab. 21 (*Trisetetum fuscii*), ZT; KRAJINA 1933, Tab. 30 (*Rhodiolo-Deschampsietum caespitosae*), 4 r., Tab. 31
- 1085 **1090**
- 1095 **1100**

(*Trisetetum fuscii*), 6 r., VT; HADAČ 1956: 52 (*Deschampsietum caespitosae tetricum*), 1 r., p. 53 (*Rhodiolo-Deschampsietum caespitosae*), 1 r., VT; ŠEPPER 1984, Tab. 2 (*Aconito firmi-Deschampsietum alpicola*e), r. 8, VT; ŠEPPER 1991, Tab. 1 (*Rhodiolo-Deschampsietum caespitosae*), r. 2–20, VT; MIADOK 1995: 41 (*Trisetetum fuscii*), 1 z, NT.

- 1105 3 KOMÁRKOVÁ 1964, Tab. 34 (*Deschampsietum caespitosae tetricum*), r. 3, ZT; HORÁK 1971, Tab. 4 (*Deschampsia caespitosa*-typ, *Aconitum firmum-Viola biflora*-subtyp), r. 6–9, ZT; KREMLOVÁ 1974, Tab. inter p. 79–80 (*Deschampsietum caespitosae tetricum*), 7 r., ZT; DÚBRAVCOVÁ et al. 1976, Tab. 23 (*Deschampsietum caespitosae tetricum*), r. 1–5, ZT; Krajina 1933, Tab. 33 (*Deschampsietum caespitosae*), 9 r., VT; ŠEPPER 1984, Tab. 2 (*Aconito firmi-Deschampsietum alpicola*e), r. 1–3, 6, 7, VT; MIADOK 1995: 40 (*Deschampsietum caespitosae*), 3 r., NT.
- 1110 4 KOMÁRKOVÁ 1964, Tab. 28 (*Chaerophylletum cicutariae*), r. 1, ZT; BRAUN-BLANQUET 1930: 17 (*Aconitum firmum-Delphinium oxysepalum*-Ass.), 2 r., VT; KRAJINA 1933, Tab. 26 (*Aconitum firmi*), 8 r., VT; ŠEPPER 1984, Tab. 2 (*Aconito firmi-Deschampsietum alpicola*e), r. 4, 5, Tab. 4 (*Aconitum firmi*), 6 r., VT; ALTMANNOVÁ 1983: 92–93 (*Aconitum firmi*), 2 r., NT; FAJMONOVÁ 1987: 3–4, r. 7 (*Aconitum firmi*), NT; MIADOK 1995: 42 (*Aconitum firmi*), 2 r., NT; Jarolímek, Dúbravcová, Šibík ined., 1 r., KMF; Jarolímek ined., 1 r., KMF, 3 r. NT; Jarolímek, Alexyová ined., 3 r., NT; Jarolímek, Kochjarová ined., 1 r., VT; Valachovič ined., 1 r., VT.
- 1115 5 KOMÁRKOVÁ 1964, Tab. 28 (*Chaerophylletum cicutariae*), r. 3, ZT; KRAJINA 1933, Tab. 27 (*Chaerophylletum cicutariae*), 6 r., VT.

Tab. 3, the alliance *Calamagrostion variae* (1, 2), and *Festucion carpaticae* (3)

- 1125 1 SILLINGER 1933: 169 (*Calamagrostietum variae carpaticum*), 12 r., NT; ŠMARDA 1970, Tab. 5 (*Calamagrostietum variae carpaticum*), r. 1–3; Slovenský raj Mts.; Mucina ined., 1 r., Chočské vrchy Mts., 1 r., VF, 8 r., NT, 5 r., MP; Mucina & Valachovič ined., 7 r., MP.
- 1130 2 KLIMENT 1995, Tab. 1 (*Anemono narcissiflorae-Laserpitietum latifolii*), r. 21–23, VF; SILLINGER 1932, Tab. 5, r. 6 (*Calamagrostidetum variae*), NT; UNAR, UNAROVÁ & ŠMARDA 1984, Tab. 19 (*Carduo glauci-Caricetum tatarorum calamagrostietosum variae*), 5 r., ZT; HADAČ et al. 1969: 137–138 (*Calamagrostetum variae carpaticum*), 4 r., BT; Mucina ined., 1 r., Chočské vrchy Mts.; Kliment ined., 8 r., VF; Kliment & Bernátová ined., 1 r., VF; Kliment & Turis ined., 3 r., NT, 1 r., BT.
- 1135 3 ŠIBÍK 2003: 63, r. 65, KMF; VESELÁ 1992, Tab. 2, r. 61, 62, VF; SILLINGER 1932: 5, r. 1–5, NT, r. 7, VF; ŠOMŠÁK et al. 1980, Tab. 4, 4 r., ZT; UNAR, UNAROVÁ & ŠMARDA 1984, Tab. 21, 10 r., Tab. 26 (*Senecio-Adenostyletum alliariae*), 3 r., ZT; DÚBRAVCOVÁ & HAJDÚK 1986: 46, 1 r., ZT; HADAČ 1956, Tab. 18, 3 r., VT; DOMIN 1925: 8, 11 r., BT; PAWŁOWSKI & STECKI 1927, Tab. 8, r. 6, BT; HADAČ et al. 1969: 118, 5 r., BT; ŠMARDA et al. 1971, Tab. 12, 6 r., BT; BÉLOHLÁVKOVÁ & FIŠEROVÁ 1989, Tab. 4, r. 2, 3, Chočské vrchy Mts., r. 4, VF, r. 5, 6, NT, r. 7–16, ZT; Jarolímek ined., 2 r., KMF; Jarolímek & Krajčiová ined., 3 r., KMF; Kliment & Šibík ined., 8 r., KMF; Šibík & Krajčiová ined., 6 r., KMF; Kliment ined., 1 r., VF; Petrík ined., 2 r., ZT, 4 r., BT; Kliment, Turis & Valachovič ined., 2 r., BT.

Tab. 4, the alliance *Adenostylium alliariae*

- 1145 1a KOMÁRKOVÁ 1964, Tab. 32 (*Adenostyletum alliariae*), r. 1, 2, ZT; KREMLOVÁ 1974, Tab. inter p. 44–45 (*Adenostyletum alliariae*), r. 1, 4, ZT; DÚBRAVCOVÁ et al. 1976, Tab. 19 (*Adenostyletum alliariae tetricum*), r. 5–7, 13, 18, ZT; HRABOVCOVÁ 1976, Tab. 11 (*Adenostyletum alliariae*), 7 r., ZT; KRÁLIK 1979, Tab. 12 (*Adenostyletum alliariae*), r. 2–5, ZT; KRAJINA 1933, Tab. 28 (*Adenostyletum alliariae tetricum*), r. 1–3, VT; HADAČ 1956, Tab. 19 (*Adenostyletum alliariae tetricum*), 5 r., VT; ŠOMŠÁK et al. 1981: 197

- (*Adenostyletum alliariae tetricum*), r. 1, VT; ŠEFFER & ŠEFFEROVÁ 1989, Tab. 1 (*Ranunculo platanifolii-Adenostyletum alliariae*), r. 10–15, 25, VT; SILLINGER 1933: 260 (*Adenostyletum alliariae*), r. 3, 4, NT; TRESKOŇOVÁ 1972, Tab. inter p. 72–73 (*Adenostyletum alliariae*), 4 r., NT; SVINČÁK 1975, 1 r. (*Adenostyles alliariae*-typ), NT; ALTMANNOVÁ 1983, Tab. 21 (*Adenostyletum alliariae*), r. 1–3, 5, 7, 8, 10, NT; Jarolímek & Kochjarová ined., 3 r., VT; Šoltésová & Paclová ined., 1 r., VT; Valachovič ined., 1 r., VT, 1 r., NT.
- 1b** KOMÁRKOVÁ 1964, Tab. 32 (*Adenostyletum alliariae*), r. 3, 4, ZT; KREMLOVÁ 1974, Tab. inter p. 44–45 (*Adenostyletum alliariae*), r. 2, 3, ZT; DÚBRAVCOVÁ et al. 1976, Tab. 19 (*Adenostyletum alliariae tetricum*), r. 9, 12, ZT; KRÁLIK 1979, Tab. 12 (*Adenostyletum alliariae*), r. 1, ZT; KRAJINA 1933, Tab. 28 (*Adenostyletum alliariae tetricum*), r. 4–8, VT; ŠOMŠÁK et al. 1981: 197 (*Adenostyletum alliariae tetricum*), r. 2, VT; HÁBEROVÁ & ŠOLTÉSOVÁ 1989, Tab. 2 (*Ranunculo platanifolii-Adenostyletum alliariae*), 2 r., VT; ŠEFFER & ŠEFFEROVÁ 1989, Tab. 1 (*Ranunculo platanifolii-Adenostyletum alliariae*), r. 1–4, 7, 8, 17–21, VT; MIADOK 1995: 43 (*Ranunculo platanifolii-Adenostyletum alliariae*), 2 r., NT; Šoltés, Školek & Kyselová ined., 3 r., VT; Šoltésová & Paclová ined., 6 r., VT.
- 2a** KOMÁRKOVÁ 1964, Tab. 32 (*Athyrietum alpestris*), r. 5–7, ZT; KREMLOVÁ 1974, Tab. inter p. 44–45 (*Adenostyletum alliariae*, facies with *Athyrium alpestre*), r. 6, ZT; KRÁLIK 1979: 60 (*Athyrietum*), 1 r., ZT; UNAR, UNAROVÁ & ŠMARDA 1984, Tab. 27 (*Athyrietum alpestris tetricum*), 3 r., ZT; HADAČ 1956, Tab. 20 (*Athyrietum alpestris tetricum*), 3 r., VT; ŠEFFER & ŠEFFEROVÁ 1989, Tab. 1, r. 23, 24, 30 (*Ranunculo platanifolii-Adenostyletum alliariae athyrietosum distentifolii*), VT; Chytrý ined., 1 r., VT.
- 2b** KREMLOVÁ 1974, Tab. inter p. 44–45 (*Adenostyletum alliariae*, facies with *Athyrium alpestre*), r. 9, ZT; BĚLOHLÁVKOVÁ 1980, Tab. 11 (*Athyrietum alpestris*), 6 r., KMF; Krajčiová & Šibík ined., 2 r., KMF.
- 3** KLIMENT, BERNÁTOVÁ & OBUCH 1994: 15, r. 3 (*Senecioni-Adenostyletum alliariae*), VF; SILLINGER 1933: 260 (*Adenostyletum alliariae*), r. 1, 2, NT; ALTMANNOVÁ 1983, Tab. 21 (*Adenostyletum alliariae*), r. 4, 6, NT; DÚBRAVCOVÁ & HAJDÚK 1986: 47 (*Ranunculo platanifolii-Adenostyletum alliariae*), 1 r., ZT; BRAUN-BLANQUET 1930: 16 (*Adenostyletum alliariae*), 1 r., VT; DOMIN 1930: 179 (*Adenostyleto-Aconitetum*), 1 r., BT; HADAČ et al. 1969: 154 (*Delphinietum oxysepali* prov.), 1 r., BT; Jarolímek & Krajčiová ined., 10 r., KMF; Jarolímek, Šibík & Dúbravcová ined., 7 r., KMF; Kliment & Šibík ined., 3 r., KMF; Šibík & Krajčiová ined., 5 r., KMF; Bernátová & P. Kučera ined., 1 r., VF; Jarolímek & Alexyová ined., 1 r., NT; Valachovič ined., 1 r., NT; Valachovič, Turis & Holotová ined., 1 r., NT; Mucina ined., 1 r., ZT; Šoltésová & Paclová ined., 2 r., VT; Kliment & Valachovič ined., 6 r., BT; Jarolímek ined., 1 r., NT, 1 r., Veporské vrchy Mts.; Kliment ined., 2 r., KMF, 3 r., VF.
- 4a** HADAČ et al. 1969: 152 (*Doronicetum austriaci*), r. 1–3, 5, BT; ŠMARDA et al. 1971, Tab. 15 (*Adenostyletum alliariae*), r. 1, BT.
- 4b** HADAČ et al. 1969, p. 143 (*Petasito-Senecietum nemorensis*), 4 r., p. 149 (*Senecio-Adenostyletum alliariae*), 4 r., BT.
- 5** KOPECKÝ 1971, Tab. 1, r. 15–19 (*Adenostyli-Athyrietum alpestris petasitetosum albae*), ZT.
- 6a** Jarolímek ined., 2 r., MP; Jarolímek & Kochjarová ined., 1 r., MP; Valachovič & Turis ined., 2 r., MP.
- 6b** Jarolímek & Krajčiová ined., 1 r., KMF; Mucina ined., 1 r., VF; Mucina, Jarolímek & Maglocký ined., 1 r., NT; Valachovič ined., 2 r., MP.
- 1200 Localities of relevés (Tab. 5):**
- MP, Zlatno, Zlatnica-Kremenina Valley, 837 m a. s. l., N, 5°, 10 m², E₁: 100 %, E₀: 15%, 48° 48' 40", 20° 06' 19", 17. 7. 1996, Valachovič, Turis, author's mark of

- relevé MV 1872.
2. MP, Zlatnica-Kremenina Valley, 828 m a. s. l., N, 5°, 10 m², E₁: 100 %, E₀: 10 %, 48° 48' 40", 20° 06' 19", 17. 7. 1996, Valachovič, Turis, MV 1873.
 3. MP, Trsteník Valley, between the road and brook, 895 m a. s. l., SW, 15°, 4×10 m, E₁: 100 %, E₀: 75 %, 48° 47' 24,5", 20° 05' 32,5", 12. 7. 2001, Jarolímek, Kochjarová, IJ 2928.
 4. MP, Havraník Valley, alluvium of occasional stream, 784 m a. s. l., N, 3°, 3×10 m, E₁: 100 %, E₀: 15 %, 48° 48' 41,6", 20° 04' 19,9", 9. 8. 2003, Jarolímek, IJ 3162.
 5. MP, as rel. 4, S 370 m, 813 m a. s. l., SE, 3°, 3×8 m, E₁: 100 %, E₀: 25 %, 48° 48' 35,6", 20° 04' 08,6", 8. 8. 2003, Jarolímek, IJ 3163.
 6. MP, Nižná Kľaková, mountain meadows, 1257 m a. s. l., W, 50°, 15 m², E₁: 100 %, E₀: 5 %, 48° 46' 16", 19° 57' 26", 11. 7. 2001, Valachovič, Kliment, MV 2393.
 7. NT, Jánska dolina Valley, Ohnište Mt., W, 50°, 3×5 m, E₁: 100 %, E₀: 90 %, 48° 57' 22", 19° 41' 50", 22. 7. 1984, Mucina, Maglocký, Jarolímek, IJ 1110, LM 2997.
 8. MP, Veľká Stožka Mt, 1070 m a. s. l., SW, 35°, 32 m², E₁: 100 %, 48° 46' 00", 19° 57' 20", 9. 7. 1985, Valachovič.
 9. VF, Čierny Kameň Mt., rocks above the shepherd's chalet, 1420 m a. s. l., SE, 25°, 100 m², E₁: 40 %, E₀: 5 %, 48° 56' 42", 19° 09' 13", 16. 7. 1986, Mucina, LM 3164.
 10. KMF, the ridge between pekelník and Veľký Kriváň Mt., small glen below the rocky wallet, 1525 m a. s. l., SSW, 40°, 4×6 m, E₁: 100 %, 49° 11' 23,5", 19° 01' 21,3", 22. 7. 2003 Jarolímek, Krajčiová, IJ 3129.

References

- 1225 ALTMANNOVÁ M. (1983): Subalpínska a alpínska vegetácia Nízkych Tatier a jej hodnotenie pre potreby LANDEP. [Subalpine and alpine vegetation in the Nízke Tatry Mts. and its evaluation for the needs of LANDEP.] – Mscr., PhD thesis, depon. in Ústav krajinnnej ekológie SAV, Bratislava.
- 1230 BARKMAN J. J., DOING H. & SEGAL S. (1964): Kritische Bemerkungen und Vorschläge zur quantitativen Vegetationsanalyse. – Acta Bot. Neerl. 13: 394–419.
- BĚLOHLÁVKOVÁ R. (1980): Rostlinná společenstva alpínského stupně Kriváňské Malé Fatry. [Plant communities in the alpine belt of the Krivánska Malá Fatra Mts.] – Mscr., depon. in Správa NP Malá Fatra, Varín.
- 1235 BORZA A. (1934): Studii fitosociologice in Munții Retezatului. – Bul. Grăd. Bot. Cluj 14: 1–84.
- BRAUN-BLANQUET J. (1928): Pflanzensoziologie. Grundzüge der Vegetationskunde. – Berlin.
- BRAUN-BLANQUET J. (1930): Zentralalpen und Tatra, eine pflanzensoziologische Parallele. – Veröff. Geobot. Inst. Rübel 6: 81–123. [sep. p. 1–43]
- 1240 BRAUN-BLANQUET J. (1964): Pflanzensoziologie. Grundzüge der Vegetationskunde. 3. Aufl. – Springer Verlag, Wien, 865 pp.
- COLDEA Gh. (1983): Precizări sintaxonomice asupra cenozelor cu *Deschampsia caespitosa*. – Stud. Cercet. Biol., Ser. Biol. Veg., 35: 96–100.
- CsURÓS Št., GERGELY I. & CsURÓS M. (1985): Studii fitocenologice asupra pajistilor de *Deschampsia caespitosa* (L.) P. Beauv. din R. S. România. – Contr. Bot. 1985: 141–159.
- DOMIN K. (1930): Zur Soziologie der chionophytischen Pflanzenassoziationen des Tatragebirges. – Veröff. Geobot. Inst. Rübel 6: 167–190.
- DÚBRAVCOVÁ Z. (1976): Subalpínska a alpínska vegetácia Kamenistej a Gáborovej doliny (Západné Tatry). [Subalpine and alpine vegetation in the Kamenistá and Gáborová Valleys (Západné Tatry Mts.).] – Mscr., dissertation, depon. in Prírodovedecká fakulta UK Bratislava.

- 1255 DÚBRAVCOVÁ Z. (1983): Alpínska a subalpínska vegetácia centrálnej časti Nízkych Tatier (žula). [Alpine and subalpine vegetation of the central part of the Nízke Tatry Mts. (granite).] – Mscr., partial report, depon. in Správa NP NT, Banská Bystrica.
- 1260 DÚBRAVCOVÁ Z., BERNÁTOVÁ D., DZUBINOVÁ Ľ., KLIMENT J., LISICKÁ E., MEDOVIČ J., PETRÍK A. & BERNÁT J. (1990): Nelesné rastlinné spoločenstvá Západných Karpát a hodnotenie negatívnych vplyvov na nelesnú vegetáciu Tatier. [Non forest plant communities in the West Carpathians and evaluation of the negative influences to the non forest vegetation in the Tatry Mts.] – Mscr., final report, depon. in Prírodovedecká fakulta UK Bratislava.
- 1265 DÚBRAVCOVÁ Z., FOLTÍNOVÁ J., PACLOVÁ L. & TUREČKOVÁ J. (1976): Vegetácia subalpínskeho a alpínskeho stupňa Západných Tatier. [The vegetation of the subalpine and alpine belt in the Západné Tatry Mts.] – Mscr., final report, depon. in Prírodovedecká fakulta UK Bratislava.
- DÚBRAVCOVÁ Z. & HAJDÚK J. (1986): Príspevok k výskumu vegetácie subalpínskeho stupňa Sivého vrchu v Západných Tatrách. [Contribution to the research of vegetation in the Sivý vrch Mt. in the Západné Tatry Mts.] – Zborn. Slov. Nár. Múz., Prír. Vedy, 32: 33–54.
- 1270 DÚBRAVCOVÁ Z. & ŠEFFER J. (1992): Porasty so *Salix helvetica* v Západných a Vysokých Tatrách. [Stands with the species *Salix helvetica* in the Západné and Vysoké Tatry Mts.] – Biológia (Bratislava) 47: 331–340.
- 1275 FAJMONOVÁ E. (1987): Stručný náčrt vegetácie územia navrhovaného na rozšírenie ŠPR Chabenec-Kotliská. [Outline of the vegetation in the territory proposed for enlargement of the State Nature Reservation Chabenec-Kotliská.] – Mscr., depon. in Správa NP Nízke Tatry, Banská Bystrica.
- GRABHERR G., GREIMLER J. & MUCINA L. (1993): *Seslerietea albantis*. – In: GRABHERR G. & MUCINA L. (eds): Die Pflanzengesellschaften Österreich II, p. 402–446. – Gustav Fischer Verlag, Jena.
- 1280 HÁBEROVÁ I. & ŠOLTÉSOVÁ A. (1989): Alpínska vegetácia Batizovskej doliny vo Vysokých Tatrách (*Montio-Cardaminetea*, *Mulgedio-Aconitetea*, *Salicetea herbaceae*). [Alpine vegetation of the Batizovská dolina Valley in the Vysoké Tatry Mts. (*Montio-Cardaminetea*, *Mulgedio-Aconitetea*, *Salicetea herbaceae*).] – Biológia (Bratislava) 44: 51–60.
- 1285 HADAČ E. (1956): Rostlinná společenstva Temnosmrečinové doliny ve Vysokých Tatrách. [Plant communities in the Temnosmrečinová dolina Valley in the Vysoké Tatry Mts.] – Biol. Práce Slov. Akad. Vied 2/1: 1–78.
- HADAČ E. (1962): Übersicht der höheren Vegetationseinheiten des Tatragebirges. – Vegetatio 11: 46–54.
- 1290 HADAČ E. et al. (1969): Die Pflanzengesellschaften des Tales „Dolina Siedmich prameňov“ in der Belauer Tatra. – Vegetácia ČSSR, Ser. B, 2: 5–343.
- HORÁK J. (1971): Westliche Tatra – Geobiozönosen der oberen Wald- und Krummholzgrenze. – Přír. Práce Ústavů Českoslov. Akad. Věd v Brně 5 (5): 1–47.
- 1295 HRABOVCOVÁ J. (1976): Vegetácia subalpínskeho a alpínskeho stupňa Jamnickej doliny (Západné Tatry). [The vegetation of the subalpine and alpine belt in the Jamnická dolina Valley (Západné Tatry Mts.)] – Mscr., dissertation, depon. in Prírodovedecká fakulta UK Bratislava.
- JAROLÍMEK I., KLIMENT J. & VALACHOVIC M. (2002): The syntaxonomical revision of the riparian plant communities dominated by *Petasites hybridus* in Slovakia. – Biologia (Bratislava) 57: 471–492.
- 1300 JAROLÍMEK I. & SCHLOSSER G. (1997): FYTOPACK – a system of programs to process phytosociological tables. – Biologia (Bratislava) 52: 53–59.
- JENÍK J. (1961): Alpinská vegetace Krkonoš, Králického Sněžníku a Hrubého Jeseníku. – Nakladatelství ČSAV, Praha, 409 pp.

- 1305 JENÍK J., BUREŠ L. & BUREŠOVÁ Z. (1980): Syntaxonomic study of vegetation in Velká Kotlina Cirque, the Sudeten Mountains. – *Folia Geobot. Phytotax.* 15: 1–28.
- KARNER P. & MUCINA L. (1993): *Mulgedio-Aconitetea*. – In: GRABHERR G., MUCINA L. (eds): Die Pflanzengesellschaften Österreich II, pp. 468–505. – Gustav Fischer Verlag, Jena.
- 1310 KÄSTNER M. (1938): Die Pflanzengesellschaften der Quellfluren und Bachufer und der Verband der Schwarzerlen-Gesellschaften. – Veröff. Landesver. Sächs. Heimatsch. 9: 69–118.
- KLIKA J. (1931): K výskytu *Festuca versicolor* a *F. carpatica* ve Veľké Fatře na Slovensku. [On the occurrence of the species *Festuca versicolor* and *F. carpatica* in the Veľká Fatra Mts. in Slovakia.] – Veda Přír. 12: 251–253.
- 1315 KLIKA J. (1948): Rostlinná sociologie. [Phytosociology.] – Melantrich, Praha, 382 pp.
- KLIKA J. (1955): Nauka o rostlinných společenstvech. [Teachings on plant communities.] – Nakladatelství ČSAV, Praha, 362 pp.
- KLIKA J. & HADAČ E. (1944): Rostlinná společenstva střední Evropy. [Plant communities of the Central Europe.] – *Příroda* 36: 281–295. [sep. p. 1–26]
- 1320 KLIMENT J. (1995): *Anemono narcissiflorae-Laserpitietum latifolii* Grebenščikov et al. 1956 – ozdoba hôľ Veľkej Fatry. [*Anemono narcissiflorae-Laserpitietum latifolii* Grebenščikov et al. 1956 – flourish of the natural subalpine meadows in the Veľká Fatra Mts.] – *Bull. Slov. Bot. Spoločn.* 17: 104–11.
- KLIMENT J., BERNÁTOVÁ D. & OBUCH J. (1994): On the coenology of *Conioselinum tataricum* Hoffm. in Slovakia. – *Biologia* (Bratislava) 49: 13–18.
- 1325 KLIMENT J. & JAROLÍMEK I. (2002): Syntaxonomical revision of the *Petasites kablikianus* communities (*Petasition officinalis*) in the West Carpathians. – *Biologia* (Bratislava) 57: 101–118.
- KLIMENT J. & JAROLÍMEK I. (2004): Syntaxonomical revision of the plant communities dominated by *Calamagrostis arundinacea* (alliance *Calamagrostion arundinaceae*) in Slovakia. – *Thaiszia-J. Bot.*, 13 (2003): 135–158.
- Kočí M. (2001a): Subalpine tall-forb vegetation (*Mulgedio-Aconitetea*) in the Czech Republic – syntaxonomical revision. – *Preslia* 73: 289–331.
- 1330 Kočí M. (2001b): Společenstva vysokobylinných niv (*Mulgedio-Aconitetea*) v Hrubém Jeseníku. [Tall herb communities (*Mulgedio-Aconitetea*) in the Hrubý Jeseník Mts.] – Čas. Slez. Mus. v Opavě, Ser. A, 50: 175–191.
- KOMÁRKOVÁ V. (1964): Alpinská vegetace Roháčů. [Alpine vegetation in the Roháče Mts.] – Mscr., diploma thesis, depon. in Přírodovedecká fakulta KU, Praha.
- 1335 KOPECKÝ K. (1971): Proměnlivost druhového složení nitrofilních lemových společenstev v údolí Studeného a Roháčského potoka v Liptovských holích. [Variability of the species composition of the nitrophilous fringe communities in the valleys of Roháčsky and Studený potok Brook in the Liptovké hole Mts.] – *Preslia* 43: 344–365.
- KOPECKÝ K. & HEJNÝ S. (1971): Nitrofilní lemová společenstva víceletých rostlin severovýchodních a středních Čech. [Nitrophilous communities of the perennial herbs in the north-east and central Bohemia.] – Rozpr. Českoslov. Akad. Věd, Praha, Vědy Mat.-Přír., 81/9: 1–126.
- KRAJINA V. (1933): Die Pflanzengesellschaften des Mlynica-Tales in den Vysoké Tatry (Hohe Tatra). – Beih. Bot. Cbl. 50B: 774–957, 51B: 1–224.
- 1340 KRÁLIK T. (1979): Rastlinné spoločenstvá Spálenej doliny. [Plant communities of the Spálená dolina Valley.] – Mscr., diploma thesis, depon. in Prírodovedecká fakulta UK, Bratislava.
- KREMLOVÁ R. (1974): Alpínska a subalpínska vegetácia Žiarskej doliny (Západné Tatry). [Alpine and subalpine vegetation of the Žiarska dolina Valley (Západné Tatry Mts.).] – Mscr., diploma thesis, depon. in Prírodovedecká fakulta UK, Bratislava.
- 1345 MAAREL VAN DEN E. (1979): Transformation of cover-abundance values in phytosociology and its effect on community similarity. – *Vegetatio* 39: 97–114.

- MARHOLD K. & HINDÁK F. [eds] (1998): Zoznam nižších a vyšších rastlín Slovenska. [Checklist of nonvascular and vascular plants of Slovakia.] – Veda, Bratislava, 687 pp.
- 1360 MATUSZKIEWICZ W. & MATUSZKIEWICZ A. (1975): Mapa zborowisk roślinnych Karkonoskiego Parku Narodowego. – Ochr. Przyr. 40: 45–112.
- MIADOK D. (1995): Vegetácia ŠPR Ďumbier. [Vegetation of State Nature Reservation Ďumbier.] – Univerzita Komenského, Bratislava, 70 pp.
- 1365 MORAVEC J. et al. (1983): Rostlinná společenstva České socialistické republiky a jejich ohrožení. [Plant communities of the Czech Republic and their threat.] – Severočeskou Přír., append. 1983/1: 5–110.
- MORAVEC J. et al. (1995): Rostlinná společenstva České republiky a jejich ohrožení. Ed. 2. [Plant communities of the Czech Republic and their threat. Ed. 2.] – Severočeskou Přír., append. 1995: 3–206.
- 1370 MUCINA L. & MAGLOCKÝ Š. (1984): A list of higher syntaxonomical units of Slovakia. – Tuxenia 4: 31–38.
- MUCINA L. & MAGLOCKÝ Š. [eds] (1985): A list of vegetation units of Slovakia. – Doc. Phytosoc., N. S., 9: 175–220.
- PAWŁOWSKI B. (1935): Über die Klimaxassoziationen in der alpinen Stufe der Tatra. – Bull. Int. Acad. Polon. Sci., Cl. Sci. mat., Ser. B., Sci. Nat., pp. 115–146.
- 1375 PAWŁOWSKI B. (1956): Flora Tatr I. Rośliny naczynowe. – Polska akademia nauk, Warszawa, 669 pp.
- PAWŁOWSKI B., SOKOŁOWSKI M. & WALLISCH K. (1928): Die Pflanzenassoziationen des Tatra-Gebirges. VII. Teil. Die Pflanzenassoziationen und die Flora des Morskie Oko-Tales. – Bull. Int. Acad. Polon. Sci., Cl. Sci. Math., Ser. B., Suppl. 2: 205–272.
- 1380 PILS G. (1980): Systematik, Verbreitung und Karyologie der *Festuca violacea*-Gruppe (Poaceae) in Ostalpenraum. – Pl. Syst. Evol. 136: 73–124.
- PODANI J. (1993): SYN-TAX-pc. Computer Programs for Multivariate Data Analysis in Ecology and Systematics. Ver. 5.0. – Scientia Publ., 104 pp.
- 1385 SILLINGER P. (1932): *Festucetum carpaticae* v Nízkých Tatrách ve srovnání s analogickou asociací v jiných částech oblasti západokarpatské. [The association *Festucetum carpaticae* in the Nízke Tatry Mts. and its comparison with analogous association in other parts of the West Carpathian region.] – Rozpr. České Akad. Věd, Tř. 2, Vědy Mat.-Přír., 41 (1931)/16: 1–21.
- 1390 SILLINGER P. (1933): Monografická studie o vegetaci Nízkých Tater. [Monographic study on vegetation of the Nízke Tatry Mts.] – Orbis, Praha, 339 pp.
- SUTTER R. (1978): Sind die *Centaurea rhipontica*- und die *Delphinium elatum*-Hochstaudenfluren Assoziationen? – Mitt. Ostalp.-Dinar. Ges. Vegetkde. 14: 375–385.
- 1395 SVINČÁK I. (1975): Spoločenstvá nad hornou hranicou lesa na severnej strane masívu Prašivá v Nízkych Tatrách. [Communities above the timber line in the north side of the massif Prašivá in the Nízke Tatry Mts.] – Mscr., diploma thesis, depon. in Lesnícka fakulta TU, Zvolen.
- 1400 SÝKORA T. & ŠTURSA J. (1973): Vysokostébelné nivy s dominancí kapradin v sudetských karech – *Daphno (mezereo)-Dryopteridetum filix-mas* ass. nova. [Tall herb communities dominated by ferns in the Sudeten cirques – *Daphno (mezereo)-Dryopteridetum filix-mas* ass. nova.] – Preslia 45: 338–357.
- 1405 ŠEFFER J. (1984): Spoločenstvá nív a horských lúk vo Vysokých Tatrách I. [Communities on mountain alluviums and meadows in the Vysoké Tatry Mts. I.] – Mscr., diploma thesis, depon. in Prírodovedecká fakulta UK Bratislava.
- ŠEFFER J. (1991): Spoločenstvá asociácie *Rhodiolo-Deschampsietum caespitosae* Krajina 1933 vo Vysokých Tatrách. [Communities of the association *Rhodiolo-Deschampsietum caespitosae* Krajina 1933 in the Vysoké Tatry Mts.] – Biológia (Bratislava) 46: 831–837.

- 1410 ŠEFFER J. & ŠEFFEROVÁ E. (1989): Spoločenstvá s *Adenostyles alliariae* vo Vysokých Tatrách – numerické priblíženie. [Communities with *Adenostyles alliariae* in the Vysoké Tatry Mts. – numerical approach.] – Biológia (Bratislava) 44: 43–50.
- ŠEFFER J., ŠEFFEROVÁ E. & DÚBRAVCOVÁ Z. (1989): Numerical syntaxonomy of the tall-forbs and tall-grass communities in the Tatra Mountains. – Vegetatio 81: 181–187.
- 1415 ŠEFFEROVÁ E. (1984): Spoločenstvá nív a horských lúk vo Vysokých Tatrách II. [Communities on mountain alluviums and meadows in the Vysoké Tatry Mts. II.] – Mscr., diploma thesis, depon. in Prírodovedecká fakulta UK, Bratislava.
- ŠMARDA J. (1970): Flóra a vegetace Slovenského ráje. [Flora and vegetation of the Slovenský raj Mts.] – Práce Štúd. Českoslov. Ochr. Prír., Ser. II., 4: 5–41.
- 1420 ŠOMŠÁK L., KUBÍČEK F., JURKO A., HÁBEROVÁ I., ŠIMONOVÍC V., MAJZLÁNOVÁ E., ŠOLTÉSOVÁ A., ŠOLTÉS R. & RYBÁRSKA V. (1981): Vplyv zošľapovania na vegetáciu okolia Skalnatého plesa a Hrebienka vo Vysokých Tatrách. [Trampling effect on vegetation of surrounding of the Skalnaté pleso Blind Lake and Hrebienok in the Vysoké Tatry Mts.] – Zborn. Prác Tatransk. Nár. Parku 22: 145–292.
- 1425 THEURILLAT J.-P. & MORAVEC J. (1995): Index of names of syntaxa typified in 1992. – Folia Geobot. Phytotax. 30: 363–369.
- TRESKOŇOVÁ M. (1972): Hole strednej a západnej časti Nízkych Tatier. [Mountain meadows (hole) in the west and central part of the Nízke Tatry Mts.] – Mscr., diploma thesis, depon. in Prírodovedecká fakulta UK, Bratislava.
- 1430 UNAR J., UNAROVÁ M. & ŠMARDA J. (1984): Vegetační poměry Tomanovy doliny a Žlebu spod Diery v Západních Tatrách. 1. Fytocenologické tabulky. [Vegetation of the Tomanova dolina Valley and Žlab spod Diery Ravine in the Západné Tatry Mts. 1. Phytocoenological tables.] – Folia Fac. Sci. Nat. Univ. Purkynianae Brun., Ser. Biol. 25/10: 5–101.
- 1435 UNAR J., UNAROVÁ M. & ŠMARDA J. (1985): Vegetační poměry Tomanovy doliny a Žlebu spod Diery v Západních Tatrách. 2. Charakteristika přírodních poměrů a rostlinných společenstev. [Vegetation of the Tomanova dolina Valley and Žlab spod Diery Ravine in the Západné Tatry Mts. 2. Characteristics of the natural conditions and plant communities.] – Folia Fac. Sci. Nat. Univ. Purkynianae Brun., Ser. Biol. 26/14: 5–78.
- 1440 WEBER H. E., MORAVEC J. & THEURILLAT J.-P. (2000): International Code of Phytosociological Nomenclature. 3rd edition. – J. Veg. Sci. 11: 739–768.

Received: 18 October 2004
 Revised: 14 December 2004
 Accepted: 14 December 2004

Tab. 1. Communities of the alliance *Calamagrostion villosae* in Slovakia.

Festuco picturatae-Calamagrostietum villosae (1), subass. typicum (1a), subass. crepidetosum conyzifoliae (1b); Vaccinio myrtilli-Calamagrostietum villosae (2), subass. inops (2a), subass. avenuletosum versicoloris (2b); Calamagrostio villosae-Salicetum helveticae (3)

1450 1455	Community	1a	1b	1	2a	2b	2	3		
	Number of relevés	83	48	131	16	45	61	16		
	Average number of taxa	20	33	25	17	24	22	20		
Diagnostic taxa of the associations										
Cv,fp	<i>Festuca picturata</i>	D1	89 ⁴	90 ⁴	89⁴	25 ³	38 ²	34 ²	38 ²	
MU	<i>Veratrum *lobelianum</i>	D1*	75 ³	79 ³	76³	25 ²	35 ²	31 ²	50 ³	
1460	Cv	<i>Gentiana punctata</i>	D1*	72 ³	58 ³	67³	61	38 ²	30 ²	88 ³
MU	<i>Bistorta major</i>	D1*	42 ³	56 ³	47³	19 ²	11 ²	13 ²	75 ³	
MU	<i>Acetosa arifolia</i>	D1	35 ³	52 ²	41³	6 ²	9 ²	8 ²	6 ²	
Aa	<i>Adenostyles alliariae</i>	D1	36 ³	42 ²	38²	.	.	.	13 ³	
Iv	<i>Vaccinium vitis-idaea</i>	D2	2 ²	13 ²	6 ²	50 ³	58³	56³	13 ⁴	
1465		<i>Salix helvetica</i>	D3	99⁹	
		<i>Dryopteris dilatata</i>	D3	.	2 ²	1 ²	.	.	63²	
		<i>Hylocomium splendens</i> (E ₀)	D3	13 ²	15 ³	14 ²	25 ²	4 ⁴	10 ³	63³
MU	<i>Aconitum firmum</i>	D3	8 ²	25 ²	15 ²	6 ²	.	2 ²	56²	
1470	Iv	<i>Dicranum scoparium</i> (E ₀)	D3	11 ²	17 ²	13 ²	38 ²	7 ²	15 ²	44²
		<i>Salix kitaibeliana</i>	D3	1 ¹	6 ²	3 ²	.	.	.	25²
Differential taxa of the subassociations										
		<i>Hypericum maculatum</i>	D1b	17 ²	81³	40 ³	50 ³	22 ³	30 ³	6 ²
Fs		<i>Phyteuma spicatum</i>	D1b	12 ²	79²	37 ²	.	13 ²	10 ²	.
MU		<i>Geranium sylvaticum</i>	D1b	8 ²	63³	28 ³	13 ²	4 ²	7 ²	13 ²
1475	Cv	<i>Valeriana tripteris</i>	D1b	5 ²	56²	24 ²	.	2 ³	2 ³	13 ²
	Aa	<i>Crepis conyzifolia</i>	D1b	6 ¹	56³	24 ³	19 ³	40 ³	34 ³	.
	Ranunculus platanifolius		D1b	17 ²	50²	29 ²	.	9 ²	7 ²	.
		<i>Myosotis scorpioides</i> agg.	D1b	.	25²	9 ²
1480		<i>Hylotelephium argutum</i>	D1b	.	23²	8 ²
	Ns	<i>Anthoxanthum alpinum</i>	D2b	57 ³	67 ³	60 ³	19 ²	71³	57 ³	25 ²
JT		<i>Ranunculus pseudomontanus</i>	D2b	39 ²	25 ²	34 ²	6 ²	64²	49 ²	13 ³
Cv,JT		<i>Agrostis pyrenaica</i>	D2b	13 ²	13 ²	13 ²	.	47²	34 ²	6 ²
JT		<i>Campanula tatrae</i>	D2b	31 ²	56 ²	40 ²	19 ²	44²	38 ²	.
1485	JT	<i>Campanula alpina</i>	D2b	17 ²	6 ²	13 ²	.	40³	30 ³	.
	JT	<i>Festuca supina</i>	D2b	4 ²	.	2 ²	.	40²	30 ²	.
jt		<i>Juncus trifidus</i>	D2b	16 ²	15 ²	15 ²	.	38²	28 ²	6 ²
Ns		<i>Viola *sudetica</i>	D2b	5 ²	6 ⁴	5 ³	.	38²	28 ²	.
Calamagrostion villosae										
1490		<i>Calamagrostis villosa</i>	t	99 ⁸	99 ⁸	99 ⁸	99 ⁷	99 ⁷	99 ⁷	99 ⁶
jt		<i>Carex *silicicola</i>	t,D2b	39 ²	50 ³	43 ³	19 ²	84⁴	67 ⁴	13 ³
jt		<i>Hieracium alpinum</i> agg.	C,D2b	27 ²	35 ²	30 ²	13 ²	62²	49 ²	.
JT		<i>Avenula versicolor</i>	t2,D2b	25 ²	10 ²	20 ²	.	71²	52²	.
		<i>Sempervivum *carpathicum</i>	C,D2b	19 ²	29 ²	23 ²	.	36²	26 ²	.
JT		<i>Pulsatilla scherfelii</i>	D	29 ²	33 ³	31 ³	6 ³	44³	34 ³	13 ²

1495	Calamagrostietalia villosae							
	<i>Luzula *rubella</i>	41 ³	77 ³	54 ³	99 ³	69 ³	77 ³	19 ³
	<i>Campanula serratula</i>	4 ¹	6 ²	5 ²	6 ²	24 ²	20 ²	6 ²
	<i>Achillea *alpestris</i>	6 ²	17 ³	10 ²	6 ²	7 ²	7 ²	.
1500	ca <i>Calamagrostis arundinacea</i>	2 ²	4 ³	3 ³	19 ²	4 ³	8 ²	.
	tf <i>Rhodiola rosea</i>	5 ²	33 ²	15 ²	.	4 ²	3 ²	38 ²
	<i>Trommsdorffia uniflora</i>	19 ²	35 ³	25 ²	.	53 ²	39 ²	.
	ca <i>Hieracium prenanthoides</i>	.	4 ²	2 ²	6 ²	2 ²	3 ²	.
	<i>Anemone narcissiflora</i>	2 ³	8 ²	5 ²	.	2 ²	2 ²	.
	<i>Linum extraaxillare</i>	.	6 ⁴	2 ⁴	.	2 ²	2 ²	.
1505	tf <i>Trisetum fuscum</i>	.	13 ²	5 ²	.	.	.	6 ²
	tf <i>Carex aterrima</i>	.	15 ²	5 ²
	tf <i>Taraxacum alpinum</i>	.	10 ²	4 ²
	<i>Knautia maxima</i>	.	2 ²	1 ²
	<i>Crepis mollis</i>	.	2 ²	1 ²
1510	Mulgedio-Aconitetea							
	<i>Solidago *minuta</i>	64 ³	71 ³	66 ³	38 ²	62 ²	56 ²	13 ²
	<i>Gentiana asclepiadea</i>	14 ²	29 ²	20 ²	38 ²	33 ²	34 ²	6 ²
	<i>Poa chaixii</i>	12 ²	29 ³	18 ²	38 ²	4 ²	13 ²	.
1515	aa <i>Athyrium distentifolium</i>	7 ²	13 ²	9 ²	19 ²	2 ²	7 ²	6 ²
	aa <i>Milium effusum</i>	12 ²	17 ²	14 ²	6 ²	.	2 ²	.
	aa <i>Doronicum austriacum</i>	16 ²	17 ²	16 ²	.	2 ²	2 ²	.
	aa <i>Silene dioica</i>	7 ²	17 ²	11 ²	.	.	.	6 ¹
	<i>Senecio subalpinus</i>	2 ²	25 ³	11 ³
1520	aa <i>Cicerbita alpina</i>	1 ²	4 ⁴	2 ³
	<i>Valeriana *sambucifolia</i>	1 ²	2 ²	2 ²
	<i>Thalictrum aquilegiifolium</i>	10 ²	4 ²	.	2 ²	2 ²	.	.
	<i>Primula elatior</i>	.	8 ²	3 ²
	Pc <i>Chaerophyllum hirsutum</i>	.	4 ³	2 ³
	Juncetea trifida							
1525	<i>Cetraria islandica</i> (E ₀)	8 ²	6 ¹	8 ²	25 ³	24 ²	25 ³	13 ³
	<i>Oreogenum montanum</i>	92 ⁴	81 ⁴	88 ⁴	25 ³	78 ²	64 ²	81 ³
	<i>Vaccinium gaultherioides</i>	4 ²	2 ²	3 ²	13 ³	9 ⁵	10 ⁴	13 ²
	<i>Senecio *carpaticus</i>	8 ²	2 ²	6 ²	6 ⁵	16 ³	13 ³	6 ²
1530	<i>Polytrichum alpinum</i> (E ₀)	7 ²	6 ³	7 ²	.	2 ⁷	2 ⁷	31 ²
	<i>Oreochloa disticha</i>	1 ³	.	1 ³	.	4 ³	3 ³	13 ²
	<i>Bartsia alpina</i>	6 ¹	10 ²	8 ¹	.	11 ³	8 ³	.
	<i>Euphrasia tatrae</i>	4 ²	.	2 ²	.	9 ²	7 ²	.
	<i>Senecio *carniolicus</i>	.	2 ¹	1 ¹	.	2 ³	2 ³	6 ²
	Salicetea herbaceae							
1535	<i>Luzula alpinopilosa</i>	46 ³	25 ²	38 ³	6 ²	11 ²	10 ²	25 ²
	<i>Sedum alpestre</i>	2 ²	6 ²	4 ²	6 ²	2 ¹	2 ²	13 ¹
	<i>Doronicum stiriacum</i>	16 ²	10 ²	14 ²	.	2 ¹	2 ¹	38 ²
	<i>Omalotheca supina</i>	1 ¹	6 ¹	3 ¹	.	2 ²	2 ²	.
	Nardo-Calunetea							
1540	<i>Soldanella carpatica</i>	64 ²	73 ²	67 ²	50 ²	62 ²	59 ²	75 ²
	<i>Potentilla aurea</i>	77 ³	90 ³	82 ³	44 ²	87 ²	75 ²	19 ²
	<i>Homogyne alpina</i>	90 ³	79 ³	86 ³	81 ³	96 ³	92 ³	88 ³
	<i>Phleum rhaeticum</i>	18 ²	29 ²	22 ²	6 ²	4 ²	5 ²	.

1545	<i>Ligusticum mutellina</i>	96 ³	90 ³	94 ³	63 ³	87 ³	80 ³	75 ²
	<i>Calluna vulgaris</i>	.	4 ²	2 ²	13 ³	16 ³	15 ³	.
	<i>Nardus stricta</i>	.	4 ¹	2 ¹	19 ³	4 ³	8 ³	.
Other taxa								
1550	<i>Avenella flexuosa</i>	67 ³	67 ³	67 ³	99 ⁴	91 ⁴	93 ⁴	99 ⁴
	<i>Vaccinium myrtillus</i>	59 ³	73 ³	64 ³	99 ⁵	93 ⁴	95 ⁴	69 ³
	<i>Luzula sylvatica</i>	8 ²	6 ⁵	8 ³	13 ⁵	2 ³	5 ⁴	13 ³
	<i>Pinus mugo</i>	1 ²	4 ¹	2 ¹	13 ³	4 ²	7 ²	6 ²
	<i>Salix silesiaca</i>	1 ²	19 ²	8 ²	6 ²	.	2 ²	13 ²
	<i>Huperzia selago</i>	1 ¹	.	1 ¹	.	2 ³	2 ³	19 ¹
1555	<i>Deschampsia cespitosa</i>	7 ³	13 ³	9 ³	6 ²	2 ⁵	3 ⁴	.
	<i>Senecio nemorensis</i> agg.	4 ²	29 ²	13 ²	6 ²	7 ²	7 ²	.
	<i>Silene vulgaris</i>	1 ²	15 ⁴	6 ⁴	6 ¹	2 ¹	3 ¹	.
	<i>Hieracium murorum</i>	1 ³	6 ³	3 ³	13 ²	2 ²	5 ²	.
	<i>Polygonatum verticillatum</i>	1 ¹	8 ²	4 ²	6 ³	7 ²	7 ²	.
1560	<i>Melampyrum sylvaticum</i>	2 ¹	.	1 ¹	19 ²	11 ²	13 ²	.
	<i>Oxalis acetosella</i>	1 ²	4 ²	2 ²	6 ¹	.	2 ¹	.
	<i>Omalotheca norvegica</i>	10 ¹	31 ²	18 ²	.	20 ²	15 ²	.
	<i>Alchemilla</i> sp. div.	6 ³	29 ³	15 ³	.	4 ¹	3 ¹	.
	<i>Hieracium</i> sp.	5 ²	8 ³	6 ²	.	16 ²	11 ²	.
1565	<i>Pseudorchis albida</i>	5 ¹	10 ²	7 ¹	.	11 ²	8 ²	.
	<i>Leontodon hispidus</i>	4 ²	4 ²	4 ²	.	2 ¹	2 ¹	.
	<i>Rhinanthus pulcher</i>	4 ²	40 ²	17 ²	.	13 ³	10 ³	.
	<i>Jovibarba globifera</i>	2 ⁴	2 ³	2 ³	.	7 ²	5 ²	.
	<i>Thymus alpestris</i>	1 ²	17 ²	7 ²	.	9 ³	7 ³	.
1570	<i>Pilosella aurantiaca</i>	1 ²	2 ³	2 ³	.	4 ²	3 ²	.
	<i>Galium anisophyllum</i> agg.	1 ¹	13 ²	5 ²	.	7 ¹	5 ¹	.
	<i>Chamerion angustifolium</i>	1 ²	13 ¹	5 ¹	13 ²	.	3 ²	.
	<i>Cardaminopsis halleri</i>	2 ²	.	2 ²	.	2 ⁵	2 ⁵	.
	<i>Antennaria dioica</i>	1 ²	.	1 ²	.	7 ²	5 ²	.
	<i>Lilium martagon</i>	.	10 ²	4 ²	.	2 ²	2 ²	.
1575	<i>Rubus idaeus</i>	.	15 ²	5 ²	13 ³	.	3 ³	.
	<i>Sorbus aucuparia</i>	.	2 ²	1 ²	13 ¹	.	3 ¹	.
	<i>Ranunculus breyninus</i>	.	8 ²	3 ²	.	2 ³	2 ³	.
	<i>Gymnadenia conopsea</i>	.	6 ²	2 ²	.	7 ²	5 ²	.
	<i>Arabis alpina</i>	.	2 ²	1 ²	.	4 ²	3 ²	.
1580	<i>Malianthemum bifolium</i>	.	2 ¹	1 ¹	.	2 ²	2 ²	.
	<i>Phyteuma orbiculare</i>	.	2 ¹	1 ¹	.	2 ²	2 ²	.
	<i>Viola biflora</i>	11 ²	29 ³	18 ²	.	.	.	19 ¹
	<i>Leucanthemum rotundifolium</i>	2 ²	21 ³	9 ³	.	.	.	19 ²
	<i>Pilosella officinarum</i>	1 ³	4 ²	2 ²	.	.	.	6 ²
1585	<i>Picea abies</i>	.	.	.	6 ¹	2 ²	3 ²	6 ²
	<i>Juniperus communis</i>	.	.	.	25 ²	4 ³	10 ²	.
Mosses and lichens								
1590	<i>Rhytidiodelphus triquetrus</i>	4 ²	6 ³	5 ³	.	2 ²	2 ²	38 ³
	<i>Drepanocladus uncinatus</i>	6 ²	13 ²	8 ²	6 ¹	4 ²	5 ¹	.
	<i>Racomitrium canescens</i>	1 ¹	2 ¹	2 ¹	.	7 ²	5 ²	.
	<i>Pleurozium schreberi</i>	56 ⁴	18 ³	28 ⁴	14 ³	15 ²	15 ²	31 ⁴
	<i>Plagiothecium</i> sp.	5 ²	2 ²	4 ²	6 ²	.	2 ²	6 ²

	<i>Polytrichum strictum</i>	1 ³	.	1 ³	13 ³	2 ²	5 ²	31 ³
1595	<i>Rhytidadelphus squarrosus</i>	7 ²	19 ³	11 ³	.	4 ²	3 ²	19 ²
	<i>Brachythecium starkei</i>	8 ²	13 ²	10 ²	.	2 ¹	2 ¹	6 ⁵
	<i>Plagiothecium denticulatum</i>	2 ²	.	1 ²	6 ²	.	2 ²	13 ³
	<i>Barbilophozia barbata</i>	2 ³	.	1 ³	6 ²	.	2 ²	19 ²
	<i>Brachythecium reflexum</i>	11 ²	8 ²	10 ²	.	.	.	19 ²
1600	<i>Brachythecium rutabulum</i>	6 ³	2 ⁷	5 ³	6 ³	2 ¹	3 ²	.
	<i>Polytrichum formosum</i>	6 ²	2 ²	5 ²	6 ²	2 ⁷	3 ⁵	.
	<i>Polytrichum piliferum</i>	1 ²	2 ¹	2 ²	.	9 ³	7 ³	.
	<i>Barbilophozia lycopodioides</i>	6 ³	.	4 ³	.	4 ²	3 ²	.
	<i>Polytrichum</i> sp.	2 ⁴	.	2 ⁴	13 ²	.	3 ²	.
1605	<i>Diplophyllum taxifolium</i>	1 ²	.	1 ²	6 ²	.	2 ²	.
	<i>Racomitrium heterostichum</i>	1 ¹	.	1 ¹	.	7 ²	5 ²	.
	<i>Racomitrium patens</i>	1 ²	.	1 ²	.	2 ¹	2 ¹	.
	<i>Kiarea starkei</i>	1 ²	.	1 ²	.	2 ²	2 ²	.
	<i>Barbilophozia hatcheri</i>	.	2 ¹	1 ¹	6 ²	2 ²	3 ²	.
1610	<i>Polytrichum juniperinum</i>	.	4 ³	2 ³	6 ³	18 ²	15 ²	.
	<i>Tortella tortuosa</i>	.	2 ¹	2 ¹	.	2 ²	2 ²	.
	<i>Brachythecium velutinum</i>	1 ²	2 ²	2 ²	.	.	.	6 ⁵
	<i>Ptilidium ciliare</i>	.	.	.	6 ⁵	2 ²	3 ⁴	6 ¹
1615	<i>Cladonia pyxidata</i>	1 ¹	6 ¹	3 ¹	.	13 ²	11 ²	.
	<i>Cladonia coccifera</i>	1 ²	2 ¹	2 ²	6 ²	2 ²	3 ²	.
	<i>Cladonia</i> sp.	4 ²	10 ²	6 ²	13 ²	11 ²	11 ²	.
	<i>Cladonia ecmocyna</i>	1 ²	.	1 ²	.	2 ¹	2 ¹	.
	<i>Cladonia macrophyloides</i>	1 ²	.	1 ²	.	2 ²	2 ²	.

D1* differential taxa against the association *Vaccinio myrtilli-Calamagrostietum villosae*

1620 Rarely occurring taxa with low constancy:

E₁: *Acetosa scutata* 8³ (1b), 3³ (1); *Agrostis stolonifera* 6² (2a), 2² (2); *Androsace obtusifolia* 2² (1b), 1² (1); *Angelica sylvestris* 2¹ (1b), 1¹ (1); *Avenula pubescens* 1² (1b), 1² (1); *Bistorta vivipara* 4² (1a), 8² (1b), 5² (1); *Botrychium lunaria* 6¹ (1b), 2¹ (1); *Caltha *laeta* 1² (1a, 1); *Carex ovalis* 6² (2a), 2² (2); *Carlina acaulis* 2³ (1b), 1³ (1); *Coeloglossum viride* 2² (1a, 1b, 1); *Crepis paludosa* 8² (1b), 3² (1); *Digitalis grandiflora* 4³ (1b), 2³ (1); *Dryopteris filix mas* 6² (1a), 2² (1); *Empetrum hermaphroditum* 6² (2a), 2² (2); *Epilobium alsinifolium* 4² (1b), 2² (1); *E. montanum* 8² (1b), 3² (1); *Euphrasia* sp. 4² (1b), 2² (1); *Galium schultesii* 2² (1b), 1² (1); *Gentianella lutescens* 2¹ (2b, 2); *Geum rivale* 6³ (1b), 2³ (1); *Heracleum sphondylium* 15² (1b), 5² (1); *Hieracium caesium* 2² (1b), 1² (1); *H. epimedum* 8² (1b), 3² (1); *H. fritzei* 1² (1a, 1); *H. lachenalii* 7² (2b), 5² (2); *Leontodon pseudotaraxaci* 2² (1b), 1² (1); *Leucanthemopsis *tatrae* 1² (1a, 1); *Leucanthemum vulgare* agg. 6² (1b), 2² (1); *Melampyrum pratense* 6¹ (2a), 2¹ (2); *Moneses uniflora* 2² (1b), 1² (1); *Myosotis sylvatica* 2² (1a, 1); *Omalotheca sylvatica* 6¹ (2a), 2¹ (2); *Paris quadrifolia* 2² (2b, 2); *Pilosella alpicola* 2¹ (1b), 1¹ (1); *P. atrata* 2² (2b, 2); *Poa alpina* 2² (1a, 1); *P. nemoralis* 13³ (1b), 5³ (1); *Potentilla erecta* 2² (2b, 2); *Primula minima* 2¹ (1b), 1¹ (1); *Pyrola minor* 6² (3); *Ranunculus acris* 1¹ (1a), 2¹ (1b, 1); *R. nemorosus* 4² (1a, 1b, 1); *Salix caprea* 6⁵ (3); *S. herbacea* 1¹ (1a, 1); *S. phyllicifolia* 2³ (1b), 1³ (1); *S. reticulata* 1² (1a), 4² (1b), 2² (1); *Saxifraga paniculata* 2¹ (1b), 1¹ (1); *Selaginella selaginoides* 4² (1b), 2² (1), 6² (3); *Sesleria tatrae* 4¹ (1a), 2³ (1b), 3² (1); *Soldanella hungarica* 2² (2b, 2); *Sorbus chamaemespilus* 2⁵ (1b), 1⁵ (1); *Swertia *alpestris* 2² (1b), 1² (1); *Tephroseris crispa* 2² (1b), 1² (1); *Thesium alpinum* 4² (1b), 2² (1); *Thymus* sp. 4² (1b), 3² (1); *Traunsteinera globosa* 2² (1b), 1² (1); *Veronica alpina* 1² (1a), 4² (1b), 2² (1); *V. chamaedrys* 2² (2b, 2).

1645 **E.**: *Anastrepta orcadensis* 2² (2b, 2); *Anomodon viticulosus* 2² (1b), 1² (1); *Barbilophozia*
 sp. 1² (1a, 1); *Bazzania trilobata* 4² (1b), 2² (1); *Bazzania* sp. 1² (1a, 1); *Blepharostoma*
trichophyllum 1² (1a, 1), 6² (3); *Brachythecium erythrorrhizon* 2² (1b), 1² (1); *B. glareosum*
 4² (1b), 2² (1); *B. salebrosum* 1² (1a), 8³ (1b), 4² (1); *Bryum elegans* 1¹ (1a, 1);
Calypogeia azurea 2¹ (1b), 1¹ (1); *Cephalozia* sp. 2² (1a, 1); *Ceratodon purpureus* 2² (2b,
 2); *Chiloscyphus pallescens* 2¹ (1b, 1); *Climacium dendroides* 6² (3); *Desmatodon*
 1650 *latifolius* 2¹ (1b), 1¹ (1); *Dicranoweissia crispula* 6¹ (1b), 2¹ (1); *Dicranum fuscesens* 2²
 (2b, 2); *D. montanum* 2² (2b, 2); *D. polysetum* 6³ (2a), 2³ (2); *Errhynchium hians* 1² (1a),
 8⁵ (1b), 4⁴ (1); *Grimmia alpestris* 2² (1b), 1² (1); *Hylocomium umbratum* 2³ (1b), 1³ (1), 6²
 (3); *Hypnum cupressiforme* 4² (2b), 3² (2); *Lophozia wenzelii* 2¹ (1b), 1¹ (1); *Mnium*
 1655 *marginatum* 4² (1b), 2² (1); *M. thomsonii* 4³ (1b), 2³ (1); *Mnium* sp. 4² (1b), 2² (1);
Oncophorus virens 6¹ (1b), 2¹ (1); *Plagiochila asplenoides* 6² (1b), 2² (1); *Plagiommium*
cuspidatum 1² (1a), 10³ (1b), 5³ (1); *Plagiothecium laetum* 10² (1a), 4³ (1b), 8² (1); *P.*
succulentum 2² (2b, 2); *Pohlia cruda* 2¹ (1b), 1¹ (1); *P. nutans* 4² (2b), 3² (2); *Pohlia* sp. 2²
 (2b, 2); *Polytrichum commune* 6⁵ (2a), 2⁵ (2); *Ptilidium pulcherrimum* 6² (2a), 2² (2);
 1660 *Ptilium crista-castrensis* 6² (2a), 2² (2); *Racomitrium aciculare* 2¹ (2b, 2); *R. sudeticum* 2²
 (1a), 2¹ (1b, 1); *Racomitrium* sp. 2² (2b, 2); *Rhizomnium punctatum* 2² (1b), 1² (1), 6² (3);
Rhodobryum roseum 8¹ (1b), 3¹ (1); *Rhytidium rugosum* 2² (2b, 2); *Sphagnum*
girgensohnii 2³ (1a, 1), 6² (3); *S. quinquefarium* 6⁵ (3). – *Cladonia arbuscula* 4² (2b), 3²
 (2); *C. bellidiflora* 2² (2b, 2); *C. furcata* 6² (2a), 2¹ (2b), 3² (2); *C. macrophylla* 2² (2b, 2);
 1665 *C. polydactyla* 6¹ (3); *C. rangiferina* 6³ (2a), 4² (2b), 5² (2); *C. stricta* 2¹ (1b), 1¹ (1); *C.*
uncialis 2¹ (1b), 1¹ (1); *Peltigera aphthosa* 13² (3); *P. canina* 4² (1b), 2² (1); *P. malacea* 2²
 (1b), 1² (1).

Tab. 2. Communities of the alliance *Trisetion fusci* in Slovakia.

Deschamsio caespitosae-Salicetum helveticae (1); *Rhodiolo-Deschampsietum caespitosae* (2); *Phleo rhaetici-Deschampsietum caespitosae* (3); *Aconitetum firmi* (4);
1670 *Bryo pseudotriquetri-Chaerophylletum hirsuti* (5)

Community	1	2	3	4	5
Number of relevés	23	50	34	34	7
Average number of taxa	25	32	22	24	29

1675 **Diagnostic taxa of the associations**

	<i>Sphagnum capillifolium</i>	C1	43⁵	2 ¹	3 ³	.	.
	<i>Salix helvetica</i>	D1	99⁸
	<i>Swertia *alpestris</i>	D2	22 ²	58³	.	.	14 ¹
	<i>Bistorta vivipara</i>	D2	17 ²	48²	15 ²	6 ²	14 ¹
1680	NC <i>Nardus stricta</i>	D3	9 ¹	4 ²	47²	.	.
	MC <i>Chrysosplenium alternifolium</i>	D4	4 ²	14 ²	6 ²	50³	29 ²
	<i>Rumex alpinus</i>	D4	.	.	3 ²	26⁴	.
	<i>Plagiornnium affine</i>	D4	.	.	.	15²	.
1685		C5	71²
	<i>Chiloscyphus polyanthus</i> (E ₀)	C5	.	.	3 ²	3 ⁷	57²
	MC <i>Brachythecium rivulare</i> (E ₀)	D5	.	6 ⁴	6 ³	29 ⁴	86²
	MC <i>Rhizomnium punctatum</i> (E ₀)	D5	17 ³	10 ²	9 ¹	9 ³	71²
	<i>Marchantia polymorpha</i> (E ₀)	D5	13 ¹	.	6 ²	3 ³	71²
1690	aa <i>Doronicum austriacum</i>	D5	9 ³	6 ²	15 ¹	18 ³	57²
	<i>Poa pratensis</i>	D5	.	.	3 ⁵	6 ¹	57¹
	<i>Galium anisophyllum</i>	D5	.	12 ²	6 ²	.	43²
	MC <i>Philonotis seriata</i> (E ₀)	D5	.	.	3 ³	3 ¹	43²
	Mc <i>Bryum weigelii</i> (E ₀)	D5	.	.	.	9³	43²

Trisetion fusci

1695		<i>Trisetum fuscum</i>	tC2	43 ³	58⁵	9 ²	6 ²	14 ²
		<i>Rhodiola rosea</i>	tC2	74 ²	94⁶	24 ²	41 ³	29 ¹
		<i>Taraxacum alpinum</i>	C	17 ²	66 ²	41 ²	21 ²	43 ²
		<i>Cerastium fontanum</i>	tC3	9 ¹	18 ²	53²	9 ²	14 ¹
1700	Mc	<i>Bryum pseudotriquetrum</i> (E ₀)	tC5	13 ¹	28 ²	12 ²	12 ²	86¹
		<i>Carex aterrima</i>	tC2	30 ²	52²	18 ²	.	14 ¹
		<i>Cardaminopsis neglecta</i>	C	.	28 ²	12 ³	3 ²	14 ¹
	MU	<i>Aconitum firmum</i>	D	48 ⁴	82 ³	62 ³	99⁸	86 ²
	MC	<i>Caltha *laeta</i>	D	43 ²	46 ³	32 ³	44 ³	71 ²
	Mc	<i>Cardamine *opicia</i>	D	9 ³	6 ²	12 ²	26 ³	14 ¹
1705		<i>Calamagrostion villosae</i>						
		<i>Calamagrostis villosa</i>		61 ³	54³	9 ²	38 ³	.
	JT	<i>Avenula versicolor</i>		26 ¹	8 ²	3 ¹	.	.
		<i>Sempervivum *carpathicum</i>		4 ²	10 ²	.	.	.
	jt	<i>Hieracium alpinum</i> agg.		.	10 ²	6 ²	.	.
1710		<i>Calamagrostietalia villosae</i>						
	fp	<i>Festuca picturata</i>		65 ²	70 ³	68 ³	47 ³	57 ²
	Cv	<i>Gentiana punctata</i>		61 ²	44 ²	9 ¹	12 ²	14 ¹
	JT	<i>Campanula tatrae</i>		13 ²	28 ²	35 ²	.	14 ¹

	jt	<i>Carex *silicicola</i>	13 ²	46 ²	26 ²	3 ²	.
1715		<i>Luzula *rubella</i>	4 ²	14 ²	26 ²	3 ¹	.
		<i>Solidago *minuta</i>	22 ²	20 ³	26 ²	6 ²	.
		<i>Crepis conyzifolia</i>	9 ¹	4 ²	18 ²	.	.
		<i>Achillea *alpestris</i>	.	4 ³	15 ¹	9 ³	.
1720		<i>Anemone narcissiflora</i>	.	4 ³	.	6 ³	.
		<i>Campanula serratula</i>	.	.	6 ²	3 ³	.
		<i>Trommsdorffia uniflora</i>	.	2 ²	.	.	.
		<i>Adenostyletalia alliariae</i>					
		<i>Adenostyles alliariae</i>	22 ²	28 ³	3 ¹	44 ⁴	29 ²
1725		<i>Milium effusum</i>	4 ³	12 ³	3 ²	26 ³	14 ¹
		<i>Silene dioica</i>	4 ⁵	10 ³	3 ²	12 ³	29 ²
		<i>Athyrium distentifolium</i>	4 ²	18 ²	6 ²	24 ³	.
		<i>Ranunculus platanifolius</i>	9 ³	18 ³	6 ²	3 ²	.
		<i>Cicerbita alpina</i>	.	.	.	9 ³	.
		<i>Petasito-Chaerophylletalia</i>					
1730		<i>Stellaria nemorum</i>	9 ²	8 ²	21 ³	71⁶	99³
		<i>Chaerophyllum hirsutum</i>	17 ⁴	22 ³	9 ²	62⁵	99⁷
		<i>Carduus personata</i>	.	2 ³	.	6 ³	.
		<i>Mulgedio-Aconitetea</i>					
1735		<i>Geranium sylvaticum</i>	39 ⁴	48 ⁴	18 ²	32 ⁴	43 ¹
		<i>Bistorta major</i>	87 ²	70 ³	26 ²	35 ⁵	71 ³
		<i>Veratrum *lobelianum</i>	61 ²	20 ²	21 ²	32 ³	57 ²
		<i>Acetosa arifolia</i>	4 ²	28 ³	21 ³	53 ²	57 ²
		<i>Senecio subalpinus</i>	4 ²	24 ³	35 ³	41 ²	99 ³
1740		<i>Primula elatior</i>	9 ³	16 ²	.	9 ²	14 ⁵
		<i>Thalictrum aquilegiifolium</i>	.	18 ³	3 ¹	6 ²	.
		<i>Gentiana asclepiadea</i>	.	6 ³	3 ¹	18 ³	.
		<i>Poa chaixii</i>	.	8 ²	9 ³	3 ¹	.
		<i>Valeriana *sambucifolia</i>	4 ⁵	.	3 ¹	.	.
		<i>Juncetea trifidi</i>					
1745		<i>Oreogenum montanum</i>	74 ³	66 ³	82 ³	56 ³	43 ²
		<i>Festuca supina</i>	9 ³	.	12 ³	3 ²	.
		<i>Agrostis pyrenaica</i>	17 ²	32 ²	35 ³	.	.
		<i>Polytrichum alpinum (E₀)</i>	39 ²	20 ²	3 ³	.	.
1750		<i>Juncus trifidus</i>	13 ²	10 ²	9 ²	.	.
		<i>Campanula alpina</i>	4 ²	4 ²	9 ²	.	.
		<i>Oreochloa disticha</i>	13 ²	2 ¹	6 ²	.	.
		<i>Saxifraga moschata</i>	.	4 ²	9 ³	3 ²	.
		<i>Salix kitaibeliana</i>	22 ³	14 ³	.	.	.
1755		<i>Pulsatilla scherfelii</i>	17 ¹	16 ²	.	.	.
		<i>Cladonia gracilis (E₀)</i>	9 ¹	4 ¹	.	.	.
		<i>Vaccinium vitis-idaea</i>	13 ²	.	3 ²	.	.
		<i>Cetraria islandica (E₀)</i>	9 ²	.	3 ²	.	.
		<i>Euphrasia tatrae</i>	.	2 ¹	9 ²	.	.
1760		<i>Luzula sudetica</i>	.	4 ²	9 ¹	.	.
		<i>Primula minima</i>	.	2 ¹	9 ³	.	.
		<i>Pedicularis oederi</i>	.	20 ²	.	3 ²	.

	<i>Pedicularis verticillata</i>	.	4 ²	.	3 ²	.
	<i>Gentiana frigida</i>	4 ²
	<i>Saussurea alpina</i>	.	8 ³	.	.	.
1765	<i>Lloydia serotina</i>	.	2 ¹	.	.	.
	<i>Silene acaulis</i>	.	.	9 ³	.	.
	<i>Antennaria *carpatica</i>	.	.	3 ²	.	.
	<i>Vaccinium gaultherioides</i>	.	.	3 ²	.	.
	Salicetea herbaceae					
1770	<i>Luzula *obscura</i>	61 ²	56 ³	38 ²	41 ³	86 ²
	<i>Sedum alpestre</i>	9 ¹	12 ²	3 ¹	3 ¹	.
	<i>Salix herbacea</i>	22 ²	8 ⁴	3 ²	.	.
	<i>Veronica alpina</i>	9 ²	4 ²	15 ²	.	.
1775	<i>Omalotheca supina</i>	4 ¹	2 ²	6 ³	.	.
	<i>Myosotis alpestris</i>	.	8 ²	.	12 ³	.
	<i>Doronicum stiriacum</i>	13 ²	2 ²	.	.	.
	<i>Leontodon pseudotaraxaci</i>	.	10 ²	.	.	.
	<i>Saxifraga androsacea</i>	.	6 ¹	.	.	.
	Thlaspietea rotundifolii					
1780	<i>Arabis alpina</i>	9 ³	10 ³	6 ³	15 ²	29 ²
	<i>Delphinium oxysepalum</i>	9 ²	12 ⁴	.	18 ³	.
	<i>Poa granitica</i>	.	4 ²	6 ²	6 ³	.
	<i>Oxyria digyna</i>	.	2 ³	.	.	14 ¹
	<i>Cerastium *glandulosum</i>	.	.	6 ⁴	.	.
1785	Nardo-Callunetea					
	<i>Ligusticum mutellina</i>	74 ³	98 ³	82 ³	71 ³	86 ²
	<i>Soldanella carpatica</i>	65 ²	56 ²	38 ²	32 ²	29 ²
	<i>Phleum rhaeticum</i>	26 ²	32 ²	56 ²	15 ²	43 ¹
	<i>Potentilla aurea</i>	22 ²	76 ²	74 ²	18 ²	71 ¹
1790	<i>Homogyne alpina</i>	52 ²	48 ²	41 ²	26 ²	.
	<i>Ranunculus pseudomontanus</i>	13 ²	36 ²	24 ³	21 ²	.
	Montio-Cardaminetea					
	<i>Epilobium anagallidifolium</i>	13 ¹	2 ²	6 ³	.	.
1795	<i>Philonotis fontana</i> (E ₀)	.	2 ²	6 ³	3 ³	.
	<i>Palustriella commutata</i> (E ₀)	.	2 ²	3 ³	.	14 ³
	<i>Epilobium alsinifolium</i>	.	.	15 ²	18 ²	14 ¹
	<i>Pohlia wahlenbergii</i> (E ₀)	.	8 ³	.	.	.
	<i>Scapania undulata</i> (E ₀)	.	8 ²	.	.	.
1800	<i>Bryum schleicheri</i> (E ₀)	.	2 ⁵	.	.	.
	<i>Saxifraga aizoides</i>	.	.	3 ⁵	.	.
	<i>Cratoneurum filicinum</i> (E ₀)	.	.	3 ¹	.	.
	<i>Stellaria alsine</i>	.	.	.	3 ¹	.
	<i>Pellia neesiana</i> (E ₀)	.	.	.	3 ²	.
	Other taxa					
1805	<i>Deschampsia cespitosa</i>	57 ⁴	84 ⁶	99 ⁸	82 ⁴	99 ⁴
	<i>Viola biflora</i>	52 ³	84 ³	53 ³	71 ³	57 ³
	<i>Alchemilla</i> sp. div.	57 ³	80 ³	79 ⁴	85 ³	99 ⁵
	<i>Anthoxanthum alpinum</i>	52 ²	52 ²	26 ²	9 ²	14 ²
	<i>Leucanthemum rotundifolium</i>	30 ³	20 ³	9 ⁴	29 ³	29 ⁴

1810	<i>Poa alpina</i>	9 ²	50 ²	32 ³	21 ⁴	43 ¹
	<i>Myosotis scorpioides</i> agg.	4 ⁵	14 ²	12 ²	35 ³	71 ²
	<i>Geum rivale</i>	17 ²	4 ²	.	35 ³	29 ³
	<i>Phyteuma spicatum</i>	4 ²	22 ²	6 ²	.	43 ¹
1815	<i>Avenella flexuosa</i>	30 ²	36 ²	41 ³	12 ²	.
	<i>Vaccinium myrtillus</i>	13 ²	12 ²	6 ¹	3 ³	.
	<i>Hypericum maculatum</i>	.	34 ³	24 ³	26 ³	71 ¹
	<i>Senecio nemorensis</i> agg.	.	8 ³	9 ³	32 ³	29 ¹
	<i>Poa nemoralis</i>	.	6 ²	3 ³	12 ³	14 ²
1820	<i>Rubus idaeus</i>	.	6 ²	6 ²	9 ²	14 ¹
	<i>Agrostis capillaris</i>	.	4 ²	18 ³	9 ²	14 ²
	<i>Crepis paludosa</i>	.	4 ²	3 ²	6 ²	14 ²
	<i>Leontodon hispidus</i>	.	2 ²	12 ²	3 ¹	29 ¹
	<i>Archangelica officinalis</i>	17 ²	20 ⁴	.	12 ⁵	.
1825	<i>Myosotis</i> sp.	4 ²	2 ²	.	3 ²	.
	<i>Ranunculus nemorosus</i>	17 ²	10 ²	12 ²	.	.
	<i>Dryopteris filix-mas</i>	9 ²	4 ²	6 ²	.	.
	<i>Dryopteris carthusiana</i>	4 ¹	6 ²	3 ¹	.	.
	<i>Juncus filiformis</i>	4 ¹	2 ²	3 ⁵	.	.
1830	<i>Heracleum sphondylium</i>	.	8 ²	3 ²	12 ²	.
	<i>Chamerion angustifolium</i>	.	8 ²	3 ¹	3 ¹	.
	<i>Luzula sylvatica</i>	.	4 ³	6 ²	3 ²	.
	<i>Eriophorum vaginatum</i>	4 ²	.	3 ²	.	.
	<i>Thymus alpestris</i>	.	20 ³	21 ³	.	.
Mosses and lichens						
1835	<i>Drepanocladus uncinatus</i>	26 ²	36 ²	18 ²	12 ²	.
	<i>Brachythecium starkei</i>	17 ³	10 ²	6 ²	9 ³	.
	<i>Plagiothecium</i> sp.	9 ²	4 ²	.	3 ³	.
	<i>Brachythecium reflexum</i>	13 ⁴	2 ³	.	6 ³	.
1840	<i>Scapania subalpina</i>	13 ²	4 ²	.	3 ¹	.
	<i>Scapania helvetica</i>	9 ²	2 ¹	.	3 ³	.
	<i>Plagiothecium denticulatum</i>	4 ²	.	.	12 ¹	14 ¹
	<i>Dicranum scoparium</i>	13 ²	4 ²	3 ⁵	.	.
	<i>Pleurozium schreberi</i>	13 ²	4 ³	15 ²	.	.
1845	<i>Hylocomium splendens</i>	13 ²	4 ¹	3 ²	.	.
	<i>Polytrichum strictum</i>	9 ²	2 ²	3 ⁵	.	.
	<i>Oncophorus virens</i>	4 ¹	16 ²	3 ³	.	.
	<i>Pohlia drummondii</i>	4 ¹	6 ¹	3 ²	.	.
	<i>Aulacomnium palustre</i>	4 ¹	4 ²	3 ²	.	.
1850	<i>Campylium stellatum</i>	.	24 ²	9 ²	3 ²	14 ²
	<i>Brachythecium salebrosum</i>	.	8 ²	26 ²	6 ²	.
	<i>Brachythecium</i> sp.	.	6 ²	9 ²	9 ⁴	.
	<i>Mnium thomsonii</i>	.	4 ¹	3 ¹	3 ³	.

D2* differential taxa against herb-grass communities of the alliance (i. e. excluding *Deschampsio-Salicetum helveticae*)

1855 D4* differential taxa against the associations *Phleo rhaetici-Deschampsietum caespitosae* and *Bryo pseudotriquetri-Chaerophylletum hirsuti*

Rarely occurring taxa with low constancy:

E₁: *Adoxa moschatellina* 3¹ (3), 3³ (4); *Allium *sibiricum* 2⁵ (2); *Alopecurus pratensis* 3²

1860	(4); <i>Athyrium filix-femina</i> 3 ² (4); <i>Bartsia alpina</i> 10 ² (2); <i>Botrychium lunaria</i> 3 ¹ (3, 4); <i>Briza media</i> 6 ¹ (3); <i>Calamagrostis arundinacea</i> 4 ⁵ (1); <i>Callianthemum coriandriifolium</i> 8 ² (2); <i>Cardaminopsis halleri</i> 6 ² (4); <i>Carex nigra</i> 2 ¹ (2), 6 ³ (3); <i>Cerastium cerastioides</i> 4 ³ (2), 9 ³ (3); <i>Cirsium heterophyllum</i> 6 ⁴ (3), 14 ¹ (5); <i>Cochlearia tatrae</i> 3 ² (4); <i>Coeloglossum viride</i> 4 ² (1); <i>Cruciata glabra</i> 3 ² (3); <i>Cystopteris fragilis</i> 6 ² (2), 3 ¹ (4); <i>Dryopteris dilatata</i> 4 ² (1);
1865	<i>Epilobium alpestre</i> 2 ³ (2); <i>E. montanum</i> 2 ² (2), 6 ² (4); <i>Euphrasia</i> sp. 2 ³ (2); <i>Festuca rubra</i> 3 ² (3); <i>Filipendula ulmaria</i> 3 ² (4); <i>Gentiana nivalis</i> 2 ¹ (2); <i>Gymnocarpium dryopteris</i> 3 ² (3); <i>Hieracium murorum</i> 3 ² (4); <i>Hieracium</i> sp. 3 ² (3); <i>Huperzia selago</i> 13 ¹ (1); <i>Hylotelephium argutum</i> 3 ² (4); <i>Juniperus sibirica</i> 3 ² (3); <i>Knautia kitaibelii</i> 2 ² (2); <i>Leontodon autumnalis</i> 3 ³ (3); <i>Leucanthemopsis *tatrae</i> 4 ¹ (2); <i>Leucanthemum vulgare</i> agg. 3 ¹ (3); <i>Lilium martagon</i> 3 ¹ (3); <i>Myosotis sylvatica</i> 12 ² (4); <i>Omalotheca norvegica</i> 6 ¹ (3); <i>Oxalis acetosella</i> 3 ¹ (3), 3 ³ (4); <i>Petasites albus</i> 3 ² (4); <i>Picea abies</i> 3 ² (3); <i>Pilosella officinarum</i> 3 ¹ (3); <i>Pinus mugo</i> 3 ² (4); <i>Polygonatum verticillatum</i> 4 ² (2); <i>Polystichum lonchitis</i> 2 ³ (2), 3 ² (3); <i>Primula farinosa</i> 3 ² (3); <i>Pseudorchis albida</i> 3 ¹ (3); <i>Ranunculus acris</i> 3 ² (3); <i>R. breyninus</i> 3 ² (3); <i>R. repens</i> 6 ² (3); <i>Rhinanthus pulcher</i> 10 ² (2), 12 ² (3); <i>Sagina saginoides</i> 2 ² (2), 3 ¹ (4); <i>Salix phyllicifolia</i> 8 ² (2); <i>Saxifraga carpatica</i> 4 ² (2), 3 ² (4); <i>S. rotundifolia</i> 6 ² (4); <i>Scrophularia nodosa</i> 2 ² (2); <i>S. scopolii</i> 3 ³ (4); <i>Selaginella selaginoides</i> 8 ¹ (2), 3 ² (3); <i>Silene pusilla</i> 2 ³ (2), 18 ⁴ (3); <i>Taraxacum fontanum</i> 6 ³ (2); <i>T. officinale</i> agg. 6 ² (3), 6 ³ (4); <i>Tephroseris crispa</i> 8 ¹ (2), 6 ² (4); <i>Trifolium repens</i> 3 ² (3); <i>Trisetum</i> sp. 3 ² (4); <i>Trollius altissimus</i> 8 ² (2), 9 ⁶ (4); <i>Urtica dioica</i> 3 ³ (4); <i>Valeriana tripteris</i> 12 ³ (2), 14 ² (5); <i>Veronica aphylla</i> 2 ² (2); <i>V. chamaedrys</i> 3 ² (3); <i>Veronica</i> sp. 4 ² (2); <i>Viola *sudetica</i> 3 ² (3).
1870	Eo: <i>Anomodon viticulosus</i> 2 ¹ (2); <i>Anthelia juratzkana</i> 9 ¹ (1); <i>Barbilophozia attenuata</i> 2 ⁵ (2), 3 ² (4); <i>B. hatcheri</i> 2 ² (2), 6 ² (4); <i>Barbilophozia</i> sp. 2 ² (2); <i>Bartramia ithyphylla</i> 2 ¹ (2), 14 ¹ (5); <i>Blepharostoma trichophyllum</i> 4 ² (1), 6 ² (2); <i>Blindia acuta</i> 6 ³ (3); <i>Blindia</i> sp. 2 ² (2); <i>Brachythecium glaciale</i> 2 ¹ (2); <i>B. rutabulum</i> 3 ² (4); <i>B. vanekii</i> 4 ³ (2); <i>B. velutinum</i> 4 ⁵ (1); <i>Bryum pallescens</i> 6 ¹ (2), 3 ¹ (4); <i>Cephalozia bicuspidata</i> 9 ¹ (1), 10 ² (2); <i>Chiloscyphus pallescens</i> 14 ¹ (5); <i>Cirriphyllum piliferum</i> 4 ¹ (1), 3 ² (3); <i>Climaciumpendroides</i> 6 ² (2); <i>Ctenidium molluscum</i> 2 ² (2); <i>Desmatodon latifolius</i> 2 ² (2), 3 ³ (3); <i>Dicranoweissia crispula</i> 3 ³ (3); <i>Dicranum bonjeanii</i> 4 ² (2), 3 ¹ (4); <i>D. polysetum</i> 3 ³ (3); <i>Dicranum</i> sp. 3 ⁶ (4); <i>Diplophyllum albicans</i> 2 ¹ (2); <i>D. taxifolium</i> 2 ¹ (2); <i>Drepanocladus exanulatus</i> 6 ⁴ (3), 3 ⁷ (4); <i>Fissidens osmundoides</i> 4 ¹ (2); <i>Grimmia hartmannii</i> 3 ² (3); <i>Heterocladium dimorphum</i> 4 ² (1), 2 ¹ (2); <i>H. heterophyllum</i> 2 ¹ (2); <i>Hylocomium pyrenaicum</i> 4 ² (1), 6 ² (2); <i>H. umbratum</i> 2 ¹ (2), 3 ¹ (3); <i>Jamesoniella autumnalis</i> 4 ³ (1); <i>Jungermannia atrovirens</i> 2 ³ (2); <i>J. sphaerocarpa</i> 4 ² (2), 3 ¹ (4); <i>Kiarea falcata</i> 9 ² (1); <i>Lescuraea mutabilis</i> 4 ¹ (1), 4 ⁵ (2); <i>Lescuraea</i> sp. 8 ³ (2); <i>Leskea polycarpa</i> 12 ⁴ (2); <i>Lophozia bantriensis</i> 4 ¹ (2); <i>L. sudetica</i> 13 ² (1); <i>Marsupella adusta</i> 2 ¹ (2); <i>M. emarginata</i> 9 ² (1), 4 ¹ (2); <i>Mnium ambiguum</i> 2 ⁵ (2); <i>M. marginatum</i> 9 ⁵ (1), 6 ² (2); <i>M. spinosum</i> 4 ³ (2), 3 ⁵ (4); <i>Mnium</i> sp. 4 ³ (2), 3 ² (4); <i>Nardia scalaris</i> 2 ¹ (2); <i>Oligotrichum hercynicum</i> 3 ¹ (3); <i>Oxystegus tenuirostris</i> 4 ² (1), 6 ² (2); <i>Paraleucobryum enerve</i> 6 ² (3); <i>P. longifolium</i> 3 ¹ (4); <i>Pellia</i> sp. 10 ² (2); <i>Philonotis tomentella</i> 4 ³ (1), 3 ¹ (4); <i>Plagiochila asplenoides</i> 10 ² (2); <i>Plagiognathum undulatum</i> 3 ⁵ (4); <i>Plagiothecium laetum</i> 3 ² (3); <i>Pohlia cruda</i> 2 ¹ (2); <i>P. elongata</i> 6 ³ (3); <i>P. filum</i> 3 ¹ (3); <i>P. nutans</i> 10 ³ (2); <i>Pohlia</i> sp. 4 ² (1); <i>Polytrichum longisetum</i> 3 ³ (3); <i>P. piliferum</i> 2 ² (2); <i>P. sexangulare</i> 9 ¹ (1), 6 ² (2); <i>Polytrichum</i> sp. 3 ² (4); <i>Porella</i> sp. 2 ² (2); <i>Pseudeoleskeia incurvata</i> 3 ¹ (3), 6 ² (4); <i>Ptilidium ciliare</i> 4 ² (1); <i>Racomitrium aciculare</i> 3 ² (3); <i>R. canescens</i> 6 ⁴ (3); <i>R. fasciculare</i> 2 ² (2); <i>R. heterostichum</i> 4 ² (1), 2 ² (2); <i>R. lanuginosum</i> 2 ² (2); <i>R. patens</i> 3 ² (4); <i>Rhizomnium pseudopunctatum</i> 4 ² (1); <i>Rhytidiodelphus triquetrus</i> 9 ³ (1), 2 ³ (2); <i>Rhytidiodelphus</i> sp. 6 ³ (3), 6 ⁵ (4); <i>Scapania compacta</i> 4 ⁵ (1); <i>S. degenerii</i> 2 ³ (2); <i>S. undulata</i> 8 ² (2); <i>Scapania</i> sp. 4 ² (2), 6 ³ (3); <i>Schistidium apocarpum</i> 4 ³ (1); <i>Sphagnum girgensohnii</i> 4 ¹ (1), 3 ³ (3); <i>S. palustre</i> 2 ² (2); <i>Sphagnum</i> sp. 3 ⁵ (3); <i>Tortella tortuosa</i> 6 ² (2); <i>Tortula norwegica</i> 4 ² (2). – <i>Cladonia bellidiflora</i> 13 ² (1), 2 ¹ (2); <i>C. digitata</i> 3 ² (3); <i>C. pyxidata</i> 2 ³ (2), 6 ² (3); <i>C. stricta</i> 8 ³ (2); <i>C. sulphurina</i> 4 ² (1); <i>Cladonia</i> sp. 4 ² (1), 4 ³ (2);
1880	
1885	
1890	
1895	
1900	
1905	
1910	

Icmadophila ericetorum 4² (1); *Peltigera aphthosa* 4² (2); *P. canina* 4² (1), 6² (4); *P. malacea* 4¹ (1), 2¹ (2); *P. rufescens* 2¹ (2), 3¹ (4).

Tab. 3. Communities of the alliances *Calamagrostion variae* (1, 2) and *Festucion carpaticaee* (3) in Slovakia.

1915

Convallario majalis-Calamagrostietum variae (1); *Geranio sylvatici-Calamagrostietum variae* (2); *Festucetum carpaticaee* (3)

	Community	1	2	3
1920	Number of relevés	37	26	97
	Average number of taxa	39	48	40
Diagnostic taxa of the associations				
	<i>Achillea stricta</i>	D1	78³	.
1925	QF <i>Convallaria majalis</i>	D1	65³	15 ²
	QF <i>Galium schultesii</i>	D1	49²	15 ³
	<i>Polygonatum odoratum</i>	C1	46²	.
	Sc <i>Pulsatilla slavica</i>	C1	43²	.
	<i>Brachypodium pinnatum</i>	D1	41³	8 ³
1930	<i>Bupleurum falcatum</i>	C1	38²	.
	<i>Campanula rapunculoides</i>	D1	38²	.
	<i>Securigera varia</i>	D1	38²	.
	<i>Anthericum ramosum</i>	C1	22²	.
	<i>Adenophora liliifolia</i>	C1	19¹	.
	<i>Knautia slovaca</i>	C1	19³	.
1935	ES <i>Carex *tatorum</i>	D2	3 ¹	77⁴
	ES <i>Helianthemum grandiflorum</i>	D2	.	69³
	Cv <i>Phleum hirsutum</i>	D2	.	58³
	MU <i>Geranium sylvaticum</i>	D2	3 ⁴	58²
1940	Cv <i>Achillea *alpestris</i>	D2	.	54³
	MU <i>Astrantia major</i>	D2	16 ²	54²
	Cv <i>Campanula elliptica</i>	D2	8 ²	54²
	Cv <i>Campanula serrata</i>	D2	3 ²	54²
	Cv <i>Knautia maxima</i>	D2	5 ²	50³
	<i>Leontodon hispidus</i>	D2	5 ¹	50³
1945	MU <i>Primula elatior</i>	D2	.	50²
	Cv <i>Crepis mollis</i>	D2	.	46²
	Cv <i>Silene vulgaris</i>	D2	3 ¹	46²
	Cv <i>Linum extraaxillare</i>	D2	.	42³
1950				3 ²
	<i>Ranunculus nemorosus</i>	D2	.	38²
	<i>Viola biflora</i>	D3	8 ³	19 ³
	MU <i>Bistorta major</i>	D3	.	56³
				52 ³
	Cv <i>Swertia *alpestris</i>	D3	11 ²	19 ²
				52 ³
1955	Cv <i>Luzula *rubella</i>	D3	5 ³	8 ³
	MU <i>Senecio subalpinus</i>	D3	.	51 ³
	NC <i>Ligusticum mutellina</i>	D3	.	47 ³
	<i>Myosotis alpestris</i>	D3	.	36 ³
<i>Calamagrostion variae</i>				
	<i>Calamagrostis varia</i>	C	99⁸	99⁸
1960	sa <i>Knautia kitaibelii</i>	C	41²	23 ²
	<i>Gymnadenia odoratissima</i>	C	16¹	12 ²
	<i>Epipactis atrorubens</i>	C	30¹	8 ²

	ES sa	<i>Carduus glaucinus</i> <i>Acinos alpinus</i>	D D	68³ 24²	27² 8²	14 ² . .
Festucion carpaticae						
1965		<i>Festuca carpatica</i>	C	. .	23 ²	99⁸
		<i>Bartsia alpina</i>	C	33²
		<i>Sesleria tatrae</i>	D	. .	15 ³	56³
		<i>Luzula sylvatica</i>	D	3 ²	15 ³	48³
		<i>Cortusa matthioli</i>	D	16 ³	8 ²	42³
1970		Calamagrostietalia villosae				
		<i>Laserpitium latifolium</i>		89 ⁴	81 ³	14 ²
		<i>Cirsium erisithales</i>		62 ²	81 ³	34 ³
		<i>Pimpinella *rhodochlamys</i>		49 ³	88 ³	52 ⁴
		<i>Pyrethrum clusii</i>		43 ²	50 ²	13 ²
1975		<i>Cyanus mollis</i>		35 ²	38 ³	18 ³
		<i>Vicia oreophila</i>		16 ³	35 ²	5 ²
		<i>Solidago *minuta</i>		19 ¹	27 ²	12 ³
		<i>Pleurospermum austriacum</i>		11 ¹	23 ²	4 ³
1980	ca	<i>Hieracium prenanthoides</i>		8 ³	23 ²	6 ²
	ca	<i>Dianthus *latifolius</i>		5 ²	35 ²	3 ²
	ca	<i>Calamagrostis arundinacea</i>		5 ³	31 ²	23 ²
	ca	<i>Anemone narcissiflora</i>		. .	31 ³	31 ³
	ca	<i>Vicia sylvatica</i>		. .	31 ³	4 ²
1985	ca	<i>Jacea pseudophrygia</i>		. .	27 ²	1 ³
		<i>Allium victorialis</i>		. .	19 ¹	1 ²
		<i>Bupleurum *vapincense</i>		. .	15 ³	15 ³
	JT	<i>Crepis conyzifolia</i>		. .	15 ²	7 ²
	tf	<i>Campanula tatrae</i>		. .	12 ²	15 ²
1990	tf	<i>Rhodiola rosea</i>		. .	4 ²	31 ³
		<i>Cerastium fontanum</i>		. .	4 ²	5 ²
		<i>Festuca picturata</i>		15 ³
	cv	<i>Calamagrostis villosa</i>		12 ⁴
	tf	<i>Trisetum fuscum</i>		11 ³
1995	tf	<i>Carex aterrima</i>		4 ³
	cv	<i>Sempervivum *carpathicum</i>		1 ³
		<i>Trommsdorffia uniflora</i>		1 ³
		<i>Gentiana punctata</i>		1 ²
Mulgedio-Aconitetea						
2000		<i>Aconitum variegatum</i>		22 ³	15 ⁴	4 ⁴
		<i>Gentiana asclepiadea</i>		16 ²	23 ²	29 ²
	aa	<i>Silene dioica</i>		16 ²	12 ²	26 ²
	aa	<i>Ranunculus platanifolius</i>		8 ³	4 ²	7 ³
		<i>Thalictrum aquilegiifolium</i>		5 ³	12 ²	21 ³
2005	Pc	<i>Chaerophyllum hirsutum</i>		5 ³	8 ²	38 ³
	de	<i>Delphinium elatum</i>		3 ²	15 ³	7 ³
		<i>Valeriana *sambucifolia</i>		. .	19 ²	16 ³
		<i>Aconitum firmum</i>		. .	15 ³	32 ³
		<i>Poa chaixii</i>		. .	8 ²	25 ²
		<i>Veratrum *lobelianum</i>		. .	8 ²	31 ³
2010	aa	<i>Adenostyles alliariae</i>		. .	8 ²	22 ³

	<i>Acetosa arifolia</i>	.	4 ³	37 ²
Pc	<i>Carduus personata</i>	.	4 ²	8 ³
Pc	<i>Geranium phaeum</i>	.	4 ²	1 ³
Pc	<i>Petasites hybridus</i>	3 ³	.	.
2015	<i>Milium effusum</i>	.	.	10 ²
Pc	<i>Stellaria nemorum</i>	.	.	5 ²
aa	<i>Doronicum austriacum</i>	.	.	3 ²
aa	<i>Cicerbita alpina</i>	.	.	2 ³
2020	<i>Chrysosplenium alternifolium</i>	.	.	2 ²
Pc	<i>Anthriscus nitida</i>	.	.	1 ³
Elyno-Seslerietea				
	<i>Phyteuma orbiculare</i>	54 ²	73 ²	44 ²
	<i>Thesium alpinum</i>	46 ²	46 ²	6 ²
	<i>Scabiosa lucida</i>	41 ³	88 ³	36 ³
2025	<i>Sesleria albicans</i>	38 ⁴	62 ³	16 ³
	<i>Festuca tatrae</i>	38 ²	35 ³	10 ⁴
	<i>Libanotis pyrenaica</i>	38 ³	4 ⁷	10 ³
	<i>Thymus pulcherimus</i>	32 ⁴	31 ²	10 ³
	<i>Polygala *brachyptera</i>	32 ²	15 ²	12 ³
2030	<i>Anthyllis *alpestris</i>	27 ²	23 ³	4 ⁴
	<i>Galium anisophyllum</i>	24 ³	62 ²	39 ²
	<i>Bellidiastrum michelii</i>	24 ²	15 ³	24 ³
	<i>Colymbada alpestris</i>	22 ¹	4 ²	2 ²
	<i>Ranunculus breyninus</i>	16 ³	27 ²	31 ²
2035	<i>Allium *montanum</i>	8 ²	12 ⁵	8 ²
	<i>Gentianella fatrae</i>	3 ²	15 ²	1 ²
	<i>Dianthus *praecox</i>	3 ²	8 ²	2 ²
	<i>Minuartia langii</i>	3 ²	4 ²	2 ³
	<i>Euphrasia salisburgensis</i>	11 ³	12 ²	.
2040	<i>Tephroseris capitata</i>	.	4 ²	16 ²
	<i>Erysimum witmannii</i>	22 ²	.	.
	<i>Leontodon incanus</i>	19 ¹	.	.
Querco-Fagetea				
2045	<i>Fs Mercurialis perennis</i>	51 ²	50 ²	8 ³
	<i>Fragaria vesca</i>	51 ²	54 ²	4 ²
	<i>Lilium martagon</i>	43 ²	27 ²	27 ²
	<i>Hieracium murorum</i>	41 ²	38 ²	6 ²
	<i>Poa nemoralis</i>	27 ³	38 ³	13 ³
	<i>Melica nutans</i>	24 ²	15 ²	1 ²
2050	<i>Polygonatum verticillatum</i>	14 ²	31 ²	19 ³
	<i>Tithymalus amygdalooides</i>	19 ²	27 ²	6 ²
	<i>Campanula trachelium</i>	22 ²	23 ²	7 ³
	<i>Acer pseudoplatanus</i>	19 ²	8 ²	1 ²
	<i>Myosotis sylvatica</i>	11 ²	12 ²	11 ²
2055	<i>Campanula persicifolia</i>	8 ¹	8 ²	1 ²
	<i>Lathyrus vernus</i>	8 ¹	8 ¹	5 ²
	<i>Dryopteris filix-mas</i>	5 ²	4 ²	4 ²
	<i>Asarum europaeum</i>	5 ¹	4 ²	7 ³
	<i>Pulmonaria obscura</i>	3 ²	4 ²	3 ²
2060	<i>Prenanthes purpurea</i>	22 ²	4 ²	.

	<i>Carex digitata</i>	19 ³	8 ²	.
	<i>Melittis melissophyllum</i>	14 ¹	15 ²	.
	<i>Phyteuma spicatum</i>	.	12 ²	26 ²
	<i>Carex alba</i>	27 ³	.	.
2065	<i>Aquilegia vulgaris</i>	27 ¹	.	.
	<i>Clematis alpina</i>	22 ²	.	.
Other taxa				
	<i>Rubus saxatilis</i>	59 ³	35 ²	7 ²
	<i>Digitalis grandiflora</i>	51 ²	50 ²	21 ³
2070	<i>Lotus corniculatus</i>	46 ²	77 ³	21 ²
	<i>Heracleum sphondylium</i>	46 ²	35 ²	73 ³
	<i>Carlina acaulis</i>	41 ²	81 ³	23 ³
	<i>Origanum vulgare</i>	41 ³	27 ³	10 ²
2075	<i>Leucanthemum vulgare</i> agg.	38 ³	65 ²	31 ²
	<i>Valeriana tripteris</i>	38 ³	8 ³	31 ³
	<i>Tithymalus cyparissias</i>	32 ²	12 ²	2 ³
	<i>Melampyrum sylvaticum</i>	32 ⁴	4 ²	1 ²
	<i>Cardaminopsis arenosa</i> agg.	24 ²	35 ²	13 ²
2080	<i>Linum catharticum</i>	24 ²	35 ²	3 ²
	<i>Senecio nemorensis</i> agg.	24 ²	19 ²	30 ³
	<i>Tragopogon orientalis</i>	16 ¹	38 ²	6 ²
	<i>Gymnadenia conopsea</i>	16 ³	27 ²	11 ³
	<i>Clinopodium vulgare</i>	16 ²	19 ⁵	5 ²
2085	<i>Jovibarba globifera</i>	16 ²	19 ²	4 ²
	<i>Laserpitium archangelica</i>	16 ³	4 ⁶	3 ²
	<i>Carex flacca</i>	14 ¹	31 ²	6 ³
	<i>Picea abies</i>	14 ²	4 ²	2 ²
	<i>Soldanella carpatica</i>	11 ³	23 ²	57 ³
2090	<i>Euphrasia rostkoviana</i>	11 ³	19 ⁴	1 ²
	<i>Arabis hirsuta</i> agg.	11 ²	12 ¹	6 ²
	<i>Veronica chamaedrys</i>	11 ²	4 ²	4 ²
	<i>Geranium robertianum</i>	11 ³	4 ³	2 ³
	<i>Cruciata glabra</i>	8 ¹	18 ²	3 ³
2095	<i>Asplenium viride</i>	8 ²	15 ²	3 ²
	<i>Cystopteris fragilis</i>	8 ²	8 ²	5 ²
	<i>Hylotelephium argutum</i>	8 ²	8 ³	2 ³
	<i>Sorbus aucuparia</i>	8 ²	4 ²	1 ¹
	<i>Hypericum maculatum</i>	5 ²	35 ²	54 ³
2100	<i>Vaccinium vitis-idaea</i>	5 ²	15 ²	8 ²
	<i>Ajuga reptans</i>	5 ²	12 ²	1 ²
	<i>Briza media</i>	3 ¹	38 ³	1 ²
	<i>Salix silesiaca</i>	3 ¹	27 ²	6 ²
	<i>Saxifraga paniculata</i>	3 ³	23 ³	5 ²
2105	<i>Vaccinium myrtillus</i>	3 ²	15 ²	18 ³
	<i>Rhinanthus pulcher</i>	3 ¹	15 ²	10 ²
	<i>Dactylis glomerata</i>	3 ²	12 ²	11 ³
	<i>Ranunculus pseudomontanus</i>	3 ²	8 ²	12 ³
	<i>Rubus idaeus</i>	3 ²	8 ²	8 ²
2110	<i>Polystichum lonchitis</i>	3 ²	8 ²	3 ²
	<i>Arabis alpina</i>	3 ²	4 ⁵	16 ³

	<i>Taraxacum officinale</i> agg.	3 ²	4 ²	4 ²
	<i>Ranunculus</i> sp.	3 ²	4 ²	1 ²
	<i>Galium album</i>	32 ³	.	4 ³
	<i>Carex ornithopoda</i>	27 ²	4 ²	.
2115	<i>Gymnocarpium robertianum</i>	22 ³	8 ²	.
	<i>Trommsdorffia maculata</i>	19 ¹	27 ²	.
	<i>Listera ovata</i>	3 ²	23 ²	.
	<i>Trifolium pratense</i>	.	35 ²	18 ³
2120	<i>Anthoxanthum alpinum</i>	.	31 ³	24 ²
	<i>Potentilla aurea</i>	.	31 ³	34 ²
	<i>Gentianella lutescens</i>	.	27 ²	6 ³
	<i>Agrostis capillaris</i>	.	27 ³	3 ²
	<i>Alchemilla</i> sp. div.	.	23 ³	62 ³
2125	<i>Crepis paludosa</i>	.	23 ³	18 ³
	<i>Parnassia palustris</i>	.	23 ³	36 ³
	<i>Poa alpina</i>	.	19 ³	26 ²
	<i>Homogyne alpina</i>	.	12 ²	37 ³
	<i>Geum rivale</i>	.	8 ²	36 ³
2130	<i>Epilobium alpestre</i>	.	4 ²	23 ²
	<i>Pedicularis hacquetii</i>	.	4 ²	13 ²
	<i>Carlina vulgaris</i>	24 ²	.	.
	<i>Bistorta vivipara</i>	.	.	19 ²
	<i>Deschampsia cespitosa</i>	.	.	19 ²
	<i>Oreogenum montanum</i>	.	.	18 ²
2135	Mosses			
	<i>Hylocomium splendens</i>	14 ³	4 ²	6 ³
	<i>Tortella tortuosa</i>	11 ³	35 ⁴	21 ³
	<i>Rhytidium rugosum</i>	11 ³	8 ²	1 ⁵
2140	<i>Ditrichum flexicaule</i>	5 ⁴	8 ³	1 ²
	<i>Ctenidium molluscum</i>	5 ³	4 ²	4 ²
	<i>Thuidium philiberti</i>	3 ²	12 ³	4 ³
	<i>Plagiochila asplenoides</i>	3 ⁴	8 ²	1 ²
	<i>Fissidens dubius</i>	3 ²	4 ²	4 ²

2145 Rarely occurring taxa with low constancy:

E₁: *Abies alba* 11¹ (1); *Acetosa scutata* 2⁶ (3); *Aconitum moldavicum* 8¹ (1); *Actaea spicata* 3² (1); *Adoxa moschatellina* 3² (3); *Aegopodium podagraria* 3² (1); *Ajuga genevensis* 3¹ (1); *Allium ochroleucum* 8¹ (1); *A. *sibiricum* 1¹ (3); *Allium* sp. 1² (3); *Androsace chamaejasme* 4² (2); *A. lactea* 3² (1); *Angelica sylvestris* 5³ (1), 1³ (3); *Archangelica officinalis* 4¹ (2), 2² (3); *Arctostaphylos uva-ursi* 3¹ (1); *Arenaria tenella* 4² (2); *Aruncus vulgaris* 4² (2); *Asperula cynanchica* 3¹ (1); *A. tinctoria* 16² (1); *Astragalus alpinus* 6⁴ (3); *A. australis* 4² (2); *A. frigidus* 4⁴ (3); *A. glycyphyllos* 3¹ (1); *A. norvegicus* 1⁶ (3); *Avenella flexuosa* 3² (3); *Avenula planiculmis* 4² (2), 1² (3); *A. versicolor* 4² (2), 3² (3); *Betonica officinalis* 3¹ (1), 4² (2); *Biscutella laevigata* 4² (2), 12² (3); *Botrychium lunaria* 8³ (2), 14² (3); *Bupleurum ranunculoides* 4² (2), 2³ (3); *Callianthemum coriandrifolium* 3⁴ (3); *Campanula carpatica* 5³ (1); *C. cochleariifolia* 3² (1), 4² (2); *C. moravica* 5² (1); *Campanula* sp. 11³ (1); *Cardamine impatiens* 3² (1), 1² (3); *C. pratensis* 5³ (3); *Cardamine* sp. 3² (1); *Cardaminopsis halleri* 4² (2), 16⁴ (3); *Cardaminopsis* sp. 8² (1); *Carex humilis* 5² (1); *C. muricata* 3² (1), 4¹ (2); *Carex* sp. 3² (1); *Cephalanthera rubra* 3¹ (1); *Cerastium *glandulosum* 32 (3); *C. holosteoides* 3¹ (1);

Chaerophyllum aromaticum 3¹ (1), 6³ (3); Chamaecytisus hirsutus 161 (1); Chamerion angustifolium 2² (3); Cherleria sedoides 4³ (2); Cimicifuga europaea 8¹ (1), 8³ (2); Cirsium eriophorum 2² (3); C. pannonicum 11¹ (1); Coeloglossum viride 7² (3); Colymbada scabiosa 3² (1); Coronilla vaginalis 14² (1); Cotoneaster sp. 3¹ (1); Crepis alpestris 11¹ (1); C. jacquinii 3¹ (1), 1² (3); C. praemorsa 1² (3); Cyanus *dominii 5¹ (1); Cypripedium calceolus 5¹ (1); Cystopteris montana 3² (2), 2² (3); Dactylorhiza sp. 4² (2); Daphne mezereum 15² (2), 4² (3); Delphinium oxysepalum 63 (3); Dentaria enneaphyllos 8² (2), 1² (3); Dianthus nitidus 4² (2), 9² (3); D. *alpestris 15³ (2), 3² (3); Dryopteris dilatata 2² (3); Epilobium montanum 8² (1), 3² (3); E. tetragonum 1² (3); Epipactis helleborine 5¹ (1); Equisetum palustre 1² (3); Erigeron acris 8² (2); E. alpinus 1³ (3); E. atticus 2³ (3); E. hungaricus 4² (2), 4³ (3); Erysimum wahlenbergii 6² (3); Eupatorium cannabinum 3² (1); Euphrasia kernerii 4¹ (2); E. picta 12² (2), 2² (3); E. tatrae 7² (3); Fagus sylvatica 3² (1), 4² (2); Festuca amethystina 3⁴ (1); F. pallens 11³ (1); F. rubra 12² (2); F. supina 2³ (3); F. versicolor 4⁵ (2), 3² (3); Filipendula ulmaria 4² (2), 3³ (3); Galeobdolon luteum s. l. 4¹ (2), 1² (3); Galeopsis speciosa 1² (3); Galeopsis sp. 1² (3); Galium glaucum 5¹ (1); Galium sp. 3³ (1); Genista pilosa 8² (1); G. tinctoria 3⁵ (1); Gentiana clusii 4³ (2); G. cruciata 3¹ (1); G. punctata 1² (3); G. verna 4² (2), 1² (3); Gentianella sp. 8² (1); Gentianopsis ciliata 3² (1); Geranium sanguineum 11² (1); Gymnocarpium dryopteris 4² (2), 2² (3); Hedysarum hedysaroides 11³ (3); Hesperis *nivea 4³ (2), 3³ (3); Hieracium bifidum 8² (1), 4¹ (2); H. bupleuroides 8² (1); H. dentatum 1² (3); H. epimedum 3³ (3); H. lachenalii 12² (2); H. laevigatum 16¹ (1); H. sabaudum 3² (1); H. stygium 1¹ (3); H. umbellatum 14¹ (1); H. villosum 4² (2), 9² (3); Hieracium sp. 8² (3); Hippocratea comosa 3² (1); Hordelymus europaeus 4² (2); Huperzia selago 1² (3); Hypericum hirsutum 4² (2); H. perforatum 11² (1); Hypochaeris glabra 3⁵ (1); H. radicata 5³ (1); Inula salicina 3² (1); Isopyrum thalictroides 3² (3); Jacea *oxylepis 1² (3); Juniperus sibirica 4² (2); Kernera saxatilis 8² (1); Knautia arvensis 15³ (3); Knautia sp. 1² (3); Lamium maculatum 4² (2), 1³ (3); L. purpureum 3³ (1); Lathyrus pratensis 8² (2), 8³ (3); Lembotropis nigricans 3² (1); Leontodon pseudotaraxaci 2³ (3); Leontopodium alpinum 8² (2); Leucanthemum rotundifolium 3³ (1), 10³ (3); Lunaria rediviva 4² (2), 2³ (3); Luzula sudetica 1³ (3); Maianthemum bifolium 4² (2), 1² (3); Melampyrum nemorosum 3² (1); M. pratense 3¹ (1); Moehringia muscosa 3² (1); Mycelis muralis 14² (1); Myosotis scorpioides agg. 92 (3); Omalotheca norvegica 4² (3); Orchis ustulata 4² (2); Orobanche alsatica 4² (2); O. caryophyllacea 3³ (1); O. reticulata 5¹ (1); Orthilia secunda 3³ (1); Oxalis acetosella 14⁴ (1), 2² (3); Oxytropis halleri 2³ (3); Paris quadrifolia 15² (2), 9² (3); Pedicularis verticillata 14² (3); Petasites albus 3⁵ (1), 2² (3); Phelipanche purpurea 5² (1); Phleum rhaeticum 4² (2), 11² (3); Ph. pratense 1³ (3); Picris *villarsii 1³ (3); Pilosella aurantiaca 3² (3); Pinus mugo 5² (3); Platanthera bifolia 5¹ (1), 4² (2); Poa stiriaca 11² (1); Polypodium vulgare 5² (1); Polystichum aculeatum 4³ (2), 1² (3); Potentilla crantzii 7² (3); Primula auricula 3² (1); Prunella vulgaris 4¹ (2), 1² (3); Pseudorchis albida 2² (3); Pteridium aquilinum 3¹ (1); Pulmonaria mollis 4² (2), 1² (3); Ranunculus alpestris 1² (3); R. auricomus agg. 4² (2); R. lanuginosus 4² (2); R. polyanthemos 3¹ (1); R. thora 1³ (3); Rhinanthus alectorolophus 16⁴ (1); Rh. serotinus 11² (1); Rhinanthus sp. 3² (1), 1² (3); Rhodax alpestris 11² (1), 1² (3); Ribes uva-crispa 5² (1); Rosa pendulina 19² (1), 1² (3); Salix alpina 10³ (3); S. caprea 5² (1), 4¹ (2); S. hastata 4³ (3); S. kitaibeliana 2³ (3); S. myrsinifolia 1² (3); S. reticulata 8² (3); S. retusa 1⁵ (3); Salix sp. 1³ (3); Salvia glutinosa 3³ (1); S. pratensis 3² (1); Sanguisorba minor 3¹ (1); Saussurea alpina 6³ (3); S. discolor 1² (3); Saxifraga adscendens 3² (1), 2² (3); S. aizoides 1² (3); S. moschata 1² (3); S. rotundifolia 8³ (3); Scorzonera hispanica 5¹ (1); Scrophularia scopolii 4² (2), 6² (3); Sedum album 3² (1); S. atratum 4² (2), 1¹ (3); Selaginella selaginoides 4² (2), 6² (3); Senecio sp. 12 (3); Seseli osseum 8² (1); Silene bupleuroides 5³ (1); S. *sillingeri 3¹ (1); Soldanella hungarica 7⁴ (3); Sorbus aria 11² (1); S. chamaemespilus 4² (2); Stachys

2165

2170

2175

2180

2185

2190

2195

2200

2205

2210

95

alpina 8² (2), 4² (3); *Symphytum tuberosum* 4² (2, 3); *Taraxacum fontanum* 3² (3); *T. nigricans* 2³ (3); *Taraxacum* sp. 7² (3); *Tephroseris crispa* 7³ (3); *Thalictrum minus* 11² (1), 5² (3); *Thymus alpestris* 8⁴ (2), 8² (3); *Th. pulegioides* 8² (2), 3³ (3); *Thymus* sp. 7² (3); *Tofieldia calyculata* 5² (1), 1² (3); *Traunsteinera globosa* 4¹ (2), 5³ (3); *Trifolium badium* 4² (2), 4³ (3); *T. flexuosum* 3¹ (1); *T. repens* 8¹ (2), 1² (3); *Trisetum alpestre* 3² (3); *Trollius altissimus* 12¹ (2), 15³ (3); *Tussilago farfara* 5³ (1), 4² (3); *Urtica dioica* 3³ (1), 2² (3); *Verbascum nigrum* 5³ (1); *Veronica fruticans* 4² (2), 1² (3); *V. teucrium* 3¹ (1); *Vicia cracca* 8¹ (1), 3³ (3); *V. sepium* 4² (2), 8² (3); *Vincetoxicum hirundinaria* 11¹ (1); *Viola hirta* 8² (1); *Viola *sudetica* 4² (2), 1² (3); *V. riviniana* 3¹ (1).
E₀: *Barbilophozia lycopodioides* 3² (3); *Brachythecium glareosum* 2² (3); *B. reflexum* 1⁵ (3); *B. rutabulum* 3² (3); *B. salebrosum* 3³ (2); *B. starkei* 2⁶ (3); *B. velutinum* 4³ (3); *Brachythecium* sp. 5³ (1), 1² (3); *Bryoerythrophyllum rubrum* 1² (3); *Bryum capillare* 4¹ (2), 1² (3); *B. elegans* 1³ (3); *B. schleicheri* 3³ (1); *Calliergonella cuspidata* 3² (3); *Callicladium haldanianum* 4² (2); *Campylium halleri* 1² (3); *C. stellatum* 4³ (2); *Campylium* sp. 1² (3); *Cirriphyllum piliferum* 2⁴ (3); *Desmatodon latifolius* 1² (3); *Dicranella subulata* 4² (2); *Drepanocladus uncinatus* 4² (2), 1² (3); *Encalypta streptocarpa* 4³ (2), 1² (3); *Entodon concinnus* 4² (2); *Errhynchium praelongum* 3² (3); *E. schleicheri* 4² (2), 23 (3); *E. striatum* 4² (2); *Fissidens taxifolius* 1² (3); *Fissidens* sp. 3² (1); *Homalothecium sericeum* 8⁴ (2); *Hylocomium umbratum* 1³ (3); *Hypnum cupressiforme* 8³ (2); *Isopterygiopsis pulchella* 1⁵ (3); *Lescuraea mutabilis* 4² (2); *L. plicata* 4² (3); *Metzgeria* sp. 4² (1); *M. spinosum* 6⁵ (3); *Mnium marginatum* 3³ (3); *Mnium* sp. 5⁴ (1), 4² (3); *Orthothecium intricatum* 4⁵ (2); *Plagiochila* sp. 5³ (1); *Plagiomnium affine* 8² (2), 9⁴ (3); *P. cuspidatum* 1³ (3); *P. rostratum* 2² (3); *P. undulatum* 8⁴ (1); *Plagiomnium* sp. 1² (3); *Plagiothecium denticulatum* 3² (3); *Plagiothecium* sp. 3⁵ (1); *Pleurozium schreberi* 5⁴ (1); *Polytrichum* sp. 3⁷ (1); *Porella cordeana* 1² (3); *Preissia quadrata* 3³ (1); *Pseudoleskeella catenulata* 4² (2); *Pseudoscleropodium purum* 3² (1); *Ptilium* sp. 3² (1); *Racomitrium canescens* 4³ (2); *Radula complanata* 4² (2), 1² (3); *Rhizomnium punctatum* 5³ (1), 6⁴ (3); *Rhizomnium* sp. 1³ (3); *Rhodobryum roseum* 3³ (1); *Rhynchosstegium murale* 2² (3); *Rhytidadelphus squarrosus* 12³ (3); *Rh. triquetrus* 12³ (2), 9⁴ (3); *Rhytidadelphus* sp. 3³ (1), 2⁶ (3); *Schistidium apocarpum* 4² (2); *Thuidium abietinum* 3³ (1); *Thuidium* sp. 3² (1); *Timmia bavarica* 1² (3); *Tortella* sp. 5⁵ (1); *Tortula ruralis* 1² (3); *T. subulata* 1² (3); *Tritomaria quinquentdentata* 2³ (3). – *Cetraria islandica* 4³ (2); *Cladonia fimbriata* 3² (1), 4³ (2); *Cladonia furcata* 4³ (2); *C. pyxidata* 11³ (1), 2³ (3); *C. squamosa* 3² (1); *C. symphytropa* 4³ (2); *Cladonia* sp. 1² (3); *Peltigera aphthosa* 1³ (3); *P. lepidophora* 4² (2); *P. polydactylon* 4² (2); *Peltigera* sp. 5⁴ (1), 4³ (2); *Solorina saccata* 3³ (1).

Tab. 4. Communities of the suballiances *Adenostylenion alliariae* (1–2) and *Delphinienion elati* (3–6) in Slovakia.

2250 *Ranunculo platanifolii-Adenostyletum alliariae* (1), subass. *milietosum alpicolae* (1a), subass. *ranunculetosum pseudoplatani* (1b); *Adenostylo alliariae-Athyrietum alpestris* (2), subass. *typicum* (2a), subass. *avenelletosum flexuosae* (2b); *Aconito firmi-*
Adenostyletum alliariae (3); *Petasito kablikiani-Senecietum hercynici* (4), subass. *doronicetosum austriaci* (4a), subass. *crepidetosum mollis* (4b); *Chaerophyllo hirsuti-Cicerbitetum alpiniae* (5); *Geranio robertianii-Delphinietum elati* (6), subass. *orobanchetosum flavae* (6a), subass. *ranunculetosum platanifolii* (6b)

2255	Community	1a	1b	1	2a	2b	2	3	4a	4b	4	5	6a	6b	6
	Number of relevés	56	37	93	15	9	14	54	5	8	13	5	5	5	10
	Average number of taxa	23	21	22	15	13	14	30	43	48	46	22	35	47	41
Diagnostic taxa of the associations															
	<i>Poa granitica</i>	C1,D1b	7 ⁴	41²	20³
	aa <i>Doronicum austriacum</i>	D1*,D4a	63 ⁵	62 ³	62⁴	7 ²	33 ²	17 ²	54 ³	99⁶	13 ³	46 ⁶	99 ⁵	.	.
2260	Cv <i>Festuca picturata</i>	D1	71 ³	78 ³	74³	20 ²	.	13 ²	4 ²
	aa <i>Silene dioica</i>	D1*	52 ²	46 ³	49³	13 ²	.	8 ²	44 ²	80 ²	75 ²	77 ²	40 ³	40 ²	80 ²
	MU <i>Aconitum firmum</i>	D1*	36 ³	65 ³	47³	20 ³	.	13 ³	61 ⁵	40 ³	13 ²	23 ²	80 ³	.	.
	<i>Viola biflora</i>	D1*	54 ³	38 ³	47³	20 ³	.	13 ³	67 ³	99 ³	88 ³	92 ³	99 ⁴	99 ²	40 ⁴
	<i>Phleum rhaeticum</i>	D1*	29 ²	51 ²	38²	.	.	.	15 ²	20 ²	.	8 ²	.	.	.
2265	<i>Potentilla aurea</i>	D1*	34 ²	22 ²	29²	.	.	.	17 ²	.	13 ²	8 ²	.	.	.
	aa <i>Athyrium distentifolium</i>	tC2,D5*	32 ⁴	24 ³	29 ⁴	99 ⁹	99 ⁹	99⁹	28 ³	.	.	.	99⁴	.	.
	<i>Saxifraga rotundifolia</i>	C3	28⁵
	<i>Alchemilla</i> sp. div.	D3	45 ³	27 ²	38 ³	7 ²	.	4 ²	94⁴	20 ²	25 ³	23 ²	40 ³	.	20 ²
	<i>Deschampsia cespitosa</i>	D3	9 ³	35 ³	19 ³	13 ³	.	8 ³	61³	20 ²	10 ²
2270	<i>Ligusticum mutellina</i>	D3*	93 ³	92 ³	92 ³	60 ²	22 ²	46 ²	54²
	tf <i>Rhodiola rosea</i>	D3*	25 ²	35 ²	29 ²	.	.	.	41³	.	13 ²	8 ²	.	.	.
	<i>Cardaminopsis halleri</i>	D4	11 ²	.	6 ²	20 ¹	.	13 ¹	.	80 ³	99 ⁴	92⁴	.	.	20 ²
	<i>Mnium spinosum</i> (E ₀)	D4	4 ⁵	14 ³	8 ³	.	.	.	8 ³	80 ⁴	88 ⁵	85⁵	.	.	.

	Cv	<i>Pimpinella *rhodochlamys</i>	D4	9 ²	40 ³	99 ⁴	77⁴	.	.	20 ²	10 ²	
2275	Cv	<i>Cirsium erisithales</i>	D4,D6b	9 ²	60 ³	75 ³	69³	.	.	40²	20 ²	
		<i>Dactylis glomerata</i>	D4	9 ³	40 ³	88 ³	69³	.	20 ²	40 ⁴	30 ³	
		<i>Myosotis alpestris</i>	D4	5 ²	.	3 ²	.	.	4 ²	40 ⁵	88 ³	69³	.	20 ³	40 ³	30 ³	
	Fs	<i>Pulmonaria obscura</i>	D4	60 ²	75 ²	69²	
	ca	<i>Calamagrostis arundinacea</i>	D4	2 ³	.	1 ³	.	33 ²	13 ²	9 ³	20 ³	88 ⁴	62⁴	.	20 ³	40 ³	30 ³
2280	QF	<i>Galium schultesii</i>	D4,D6b	4 ⁴	40 ²	75 ⁴	62³	.	.	60²	30 ²	
		<i>Hylotelephium argutum</i>	D4	7 ²	.	4 ²	7 ⁵	.	4 ⁵	7 ²	60 ³	63 ³	62³	.	20 ³	10 ³	
		<i>Trisetum flavescens</i>	D4	11 ³	40 ²	75 ³	62³	.	20 ³	10 ³	.	
		<i>Scrophularia scopolii</i>	D4	6 ²	60 ³	50 ²	54²	.	40 ³	20 ³	.	
		<i>Cystopteris fragilis</i>	D4	4 ²	40 ²	50 ²	46²	
2285	Fs	<i>Paris quadrifolia</i>	D4	4 ²	40 ³	50 ²	46²	.	20 ²	.	10 ²	
		<i>Eurhynchium praelongum (E₀)</i>	D4	2 ⁷	80 ⁴	25 ³	46³	
	aa	<i>Cicerbita alpina</i>	D1a,D5	54⁴	11 ³	37 ⁴	40 ⁴	33 ³	38 ⁴	37 ⁴	40 ⁷	38 ²	38 ⁴	99⁴	.	.	.
		<i>Leucanthemum rotundifolium</i>	D5	18 ³	5 ⁴	13 ³	7 ³	11 ¹	8 ²	17 ³	20 ⁵	13 ²	15 ⁴	99⁴	.	.	.
2290		<i>Petasites albus</i>	D5,D6a	4 ⁵	.	.	.	99⁶	80⁴	.	40 ⁴	
	fs	<i>Myosotis scorpioides</i> agg.	D5,D6a	5 ²	.	3 ²	.	.	15 ³	.	.	.	80³	60²	.	30 ²	
		<i>Prenanthes purpurea</i>	D5	60²	.	20 ²	10 ²	
		<i>Urtica dioica</i>	D6	17 ³	40 ²	.	15 ²	.	99 ²	99 ³	99³	
	Fs	<i>Geranium robertianum</i>	D6	99 ²	60 ³	80³	.	
2295	Fs	<i>Acer pseudoplatanus</i>	D6	2 ¹	.	25 ²	15 ²	.	40 ²	60 ¹	50²	
		<i>Fragaria vesca</i>	D6	2 ²	.	13 ²	8 ²	.	60 ²	40 ²	50²	
		<i>Carex muricata</i>	D6	20 ²	40 ²	30²	.	
		<i>Clinopodium vulgare</i>	D6	20 ²	40 ²	30²	.	

Differential taxa of the subassociations

aa	<i>Milium effusum</i>	D1a,D4a	68⁴	22 ³	49 ⁴	87 ³	11 ²	58 ³	50 ⁴	99³	.	38 ³	.	40 ³	20 ³		
2300	MU	<i>Geranium sylvaticum</i>	D1a,D6b	54³	5 ²	34 ³	20 ²	.	13 ²	81 ⁴	99 ⁴	88 ³	92 ³	99 ³	.	60³	30 ³

		<i>Hypericum maculatum</i>	D1a	48³	.	29 ³	7 ²	33 ²	17 ²	63 ³	99 ³	99 ³	99 ³	.	40 ²	60 ³	50 ³
		<i>Ranunculus pseudomontanus</i>	D1b	23 ²	62³	39 ²	7 ¹	.	4 ¹	2 ¹
		<i>Poa alpina</i>	D1b	7 ²	41³	20 ³	7 ²	.	4 ²	17 ³	20 ²	10 ²
		<i>Doronicum stiriacum</i>	D1b	.	27²	11 ²	27 ³	.	17 ³
2305		<i>Desmatodon latifolius</i> (E ₀)	D1b	4 ²	41³	18 ³
		<i>Rubus idaeus</i>	D2b	4 ²	.	2 ²	7 ³	99⁴	42 ⁴	26 ³	40 ³	.	15 ³	20 ³	60 ²	20 ³	40 ³
		<i>Oxalis acetosella</i>	D2b	4 ³	.	2 ³	7 ²	99³	42 ³	20 ³	20 ²	.	8 ²	40 ³	60 ³	20 ²	40 ³
		<i>Vaccinium myrtillus</i>	D2b	4 ²	.	2 ²	.	89³	33 ³	28 ³
		<i>Dryopteris dilatata</i>	D2b	4 ²	8 ²	5 ²	.	67³	25 ³	4 ³
2310	MU	<i>Gentiana asclepiadea</i>	D2b	13 ³	.	8 ³	13 ²	56³	29 ²	43 ²	20 ³	50 ³	38 ³	60 ³	20 ²	60 ²	40 ²
		<i>Filipendula ulmaria</i>	D4a	7 ²	80³	13 ³	38 ³
		<i>Ribes petraeum</i>	D4a	80²	13 ²	38 ²
	Cv	<i>Crepis mollis</i>	D4b	6 ²	.	88²	54 ²	.	.	20 ²	10 ²
	MU	<i>Astrantia major</i>	D4b	2 ³	.	1 ³	.	.	.	20 ²	.	75³	46 ³
2315	QF	<i>Poa nemoralis</i>	D4b	6 ²	.	75³	46 ³	.	20 ³	60 ⁶	40 ⁵
	Pc	<i>Petasites kablikianus</i>	D4b,D6a	2 ³	.	1 ³	.	.	.	2 ²	.	50⁷	31 ⁷	.	80⁶	.	40 ⁶
	fc	<i>Festuca carpatica</i>	D4b	2 ²	.	1 ²	.	.	.	31 ³	.	50⁴	31 ⁴	.	20 ²	10 ²	.
	Fs	<i>Daphne mezereum</i>	D4b	6 ²	.	50³	31 ³	.	20 ¹	40 ²	30 ¹
2320	Cv	<i>Linum extraaxillare</i>	D4b	50²	31 ²	.	.	20 ³	10 ³
		<i>Origanum vulgare</i>	D4b	50³	31 ³	.	40 ³	20 ³	.
	Cv	<i>Pleurospermum austriacum</i>	D4b	50²	31 ²	.	20 ²	.	10 ²
	Cv	<i>Campanula elliptica</i>	D4b	2 ²	.	38²	23 ²	.	20 ²	10 ²	.
		<i>Eurhynchium angustirete</i> (E ₀)	D4b	50³	31 ³
2325		<i>Crepis paludosa</i>	D6a	2 ²	.	1 ²	.	.	.	28 ³	20 ²	38 ³	31 ³	40 ³	80³	.	40 ³
		<i>Roegneria canina</i>	D6a	4 ³	.	13 ³	8 ³	.	60⁴	.	30 ⁴
	Fs	<i>Stachys sylvatica</i>	D6a	60³	.	30 ³
	Fs	<i>Orobanche flava</i>	D6a	60²	.	30 ²
	Fs	<i>Impatiens noli-tangere</i>	D6a	60²	.	30 ²

	aa	<i>Ranunculus platanifolius</i>	D6b	64 ³	49 ³	58 ³	53 ³	.	33 ³	15 ²	40 ³	13 ²	23 ³	99 ³	.	80 ²	40 ²
2330	fs	<i>Polygonatum verticillatum</i>	D6b	4 ⁴	.	2 ⁴	.	11 ²	4 ²	7 ²	99 ²	63 ⁴	77 ³	.	.	80 ³	40 ³
	Cv	<i>Luzula *rubella</i>	D6b	11 ²	.	6 ²	7 ³	.	4 ³	22 ³	40 ³	13 ³	23 ³	.	.	60 ²	30 ²
		<i>Adenostylenion alliariae</i>															
	SH	<i>Luzula alpinopilosa</i>	D	50 ³	95 ⁴	68 ³	53 ³	.	33 ³	11 ³
	Cv	<i>Gentiana punctata</i>	D	80 ²	76 ³	78 ³	60 ³	.	38 ³	4 ²
2335	JT	<i>Oreogenum montanum</i>	D	80 ³	78 ³	80 ³	67 ²	11 ²	46 ²
		<i>Delphinienion elati</i>															
		<i>Delphinium elatum</i>	t6	13 ⁴	60 ²	63 ³	62 ³	.	99 ⁷	99 ⁷	99 ⁷	
		<i>Epilobium alpestre</i>	D	5 ³	.	3 ³	.	.	50 ³	60 ³	63 ²	62 ³	40 ²	.	20 ³	10 ³	
	Pc	<i>Chaerophyllum hirsutum</i>	D	13 ²	.	8 ²	.	.	72 ⁴	80 ³	88 ⁴	85 ⁴	99 ⁶	99 ⁶	40 ⁵	70 ⁶	
2340	Fs	<i>Galeobdolon luteum</i> s. l.	D,D6b	4 ⁴	99 ³	38 ²	62 ³	80 ⁵	.	60 ²	30 ²	
		<i>Luzula sylvatica</i>	D	9 ³	.	5 ³	13 ³	11 ²	13 ²	54 ³	60 ²	50 ²	54 ²	80 ²	.	20 ²	10 ²
	Pc	<i>Stellaria nemorum</i>	D,D6a	14 ³	3 ²	10 ²	.	22 ³	8 ³	59 ⁴	20 ²	.	8 ²	80 ⁴	99 ⁵	.	50 ⁵
		<i>Cortusa matthioli</i>	D	24 ³	20 ²	50 ²	38 ²	.	40 ²	.	20 ²	.
		<i>Adenostylium alliariae, Mulgedio-Aconitetea</i>															
2345	MU	<i>Acetosa arifolia</i>		91 ³	68 ⁴	82 ⁴	93 ⁴	78 ³	88 ³	87 ⁴	60 ⁴	75 ³	69 ³	.	20 ⁵	10 ⁵	
	MU	<i>Primula elatior</i>		14 ³	14 ³	14 ³	7 ³	.	4 ³	65 ³	80 ²	63 ²	69 ²	.	40 ²	20 ²	
	aa	<i>Adenostyles alliariae</i>		99 ⁸	99 ⁸	99 ⁸	93 ⁴	33 ⁴	71 ⁴	96 ⁷	60 ⁷	50 ⁶	54 ⁶	99 ⁷	.	.	
	MU	<i>Veratrum *lobelianum</i>		95 ³	84 ³	90 ³	73 ³	67 ³	71 ³	52 ²	80 ³	63 ⁵	69 ⁴	40 ³	.	.	
	cv	<i>Calamagrostis villosa</i>		91 ⁴	43 ³	72 ⁴	60 ³	89 ²	71 ³	20 ³	20 ²	13 ³	15 ³	60 ²	.	.	
2350	MU	<i>Bistorta major</i>		46 ³	43 ³	45 ³	40 ³	11 ²	29 ²	33 ³	99 ²	99 ³	99 ³	.	.	.	
	MU	<i>Solidago *minuta</i>		36 ²	22 ²	30 ²	7 ¹	11 ²	8 ²	13 ²	
	MU	<i>Thalictrum aquilegiifolium</i>		13 ²	.	8 ²	7 ²	.	4 ²	31 ²	99 ²	75 ²	85 ²	40 ³	20 ¹	60 ²	40 ²
	MU	<i>Valeriana *sambucifolia</i>		2 ²	.	1 ²	.	.	24 ²	80 ³	88 ³	85 ³	20 ³	99 ³	40 ³	70 ³	
	MU	<i>Senecio subalpinus</i>		27 ³	5 ²	18 ³	.	.	56 ²	20 ²	50 ²	38 ²	60 ³	.	.	.	
2355	MU	<i>Poa chaixii</i>		21 ³	3 ²	14 ³	.	.	26 ³	40 ²	50 ³	46 ²	.	20 ²	10 ²	.	

	Pc	<i>Chrysosplenium alternifolium</i>	2 ³	3 ²	2 ³	.	.	.	28 ³	80 ³	63 ³	69 ³	20 ²	80 ³	40 ⁴	60 ³
	Pc	<i>Carduus personata</i>	2 ²	.	1 ²	.	.	.	30 ³	80 ⁴	63 ⁴	69 ⁴	.	60 ³	60 ²	60 ²
	Cv	<i>Achillea *alpestris</i>	2 ²	.	1 ²	.	.	.	13 ²	15 ²	8 ²	.	.	40 ²	20 ²	
	Cv	<i>Campanula serratula</i>	9 ¹	.	5 ¹	.	.	.	24 ²	40 ³	20 ³	
2360	Cv	<i>Crepis conyzifolia</i>	9 ²	.	5 ²	13 ²	8 ²	.	.	20 ⁵	10 ⁵	
	Cv	<i>Knautia maxima</i>	2 ¹	.	1 ¹	40 ⁴	.	20 ⁴
	Cv	<i>Anemone narcissiflora</i>	2 ²	5 ²	3 ²	.	.	.	2 ³	
	tf	<i>Trisetum fuscum</i>	4 ³	30 ²	14 ²	.	.	.	7 ³	
	tf	<i>Taraxacum alpinum</i>	4 ²	19 ²	10 ²	
2365	cv	<i>Sempervivum *carpathicum</i>	5 ²	3 ¹	4 ²	
	Cv	<i>Campanula tatrae</i>	20 ²	.	8 ²	.	.	.	6 ²	20 ³	25 ²	23 ²	.	60 ³	40 ³	50 ³
	MU	<i>Aconitum variegatum</i>	2 ²	.	13 ³	8 ³	.	20 ²	10 ²	
	Cv	<i>Phleum hirsutum</i>	6 ²	.	13 ²	8 ²	.	20 ⁶	60 ⁵	40 ⁵
2370	cr	<i>Calamagrostis varia</i>	6 ²	20 ⁶	38 ²	31 ³	.	20 ²	10 ²	
	Cv	<i>Cyanus mollis</i>	2 ⁶	.	25 ³	15 ³	.	20 ²	10 ²	
	ca	<i>Hieracium prenanthoides</i>	2 ²	40 ³	20 ³	
	Pc	<i>Geranium phaeum</i>	2 ²	20 ²	10 ²	
	Cv	<i>Vicia oreophila</i>	2 ²	20 ³	10 ³	
	Cv	<i>Bupleurum longifolium</i>	25 ³	15 ³	.	.	20 ²	10 ²	
2375	Cv	<i>Laserpitium latifolium</i>	25 ²	15 ²	.	.	20 ⁶	.	
	ca	<i>Vicia sylvatica</i>	20 ²	13 ³	15 ³	.	.	.	20 ²	10 ²
	Pc	<i>Petasites hybridus</i>	20 ⁶	.	10 ⁶
	Cv	<i>Pyrethrum clusii</i>	20 ²	10 ²
	Querco-Fagetea															
2380		<i>Phyteuma spicatum</i>	34 ²	11 ²	25 ²	.	.	.	19 ²	60 ²	38 ²	46 ²	.	20 ¹	10 ¹	
		<i>Dryopteris filix-mas</i>	.	3 ²	1 ²	.	.	.	19 ²	60 ²	13 ²	31 ²	.	60 ²	40 ⁴	50 ³
		<i>Mercurialis perennis</i>	2 ²	20 ⁶	38 ⁶	31 ⁶	.	40 ³	20 ³	
		<i>Myosotis sylvatica</i>	17 ³	40 ³	.	15 ³	.	60 ²	20 ²	40 ²
		<i>Lilium martagon</i>	2 ¹	20 ²	38 ²	31 ²	.	20 ¹	10 ¹	

2385	<i>Epilobium montanum</i>	7 ³	.	.	.	20 ²	20 ²	60 ²	40 ²
	<i>Lathyrus vernus</i>	2 ²	20 ²	40 ³	30 ²
	<i>Aegopodium podagraria</i>	2 ²	20 ²	20 ³	20 ³
	<i>Tithymalus amygdaloïdes</i>	2 ²	40 ³	20 ³	
	<i>Asarum europaeum</i>	13 ²	8 ²	.	.	20 ²	10 ²	
2390	<i>Cardamine impatiens</i>	40 ²	20 ³	30 ²
	<i>Aruncus vulgaris</i>	20 ¹	20 ²	20 ²
	<i>Lunaria rediviva</i>	20 ²	20 ³	20 ³
	<i>Mycelis muralis</i>	20 ²	20 ²	20 ²
	<i>Ranunculus lanuginosus</i>	20 ²	20 ²	20 ²
2395	<i>Hordelymus europaeus</i>	40 ³	.	20 ³
	<i>Campanula trachelium</i>	40 ²	.	20 ²
	<i>Melica nutans</i>	40 ³	20 ³	
	Other taxa														
2400	<i>Homogyne alpina</i>	68 ²	59 ³	65 ²	40 ²	89 ²	58 ²	41 ²	.	13 ²	8 ²	40 ³	.	20 ²	10 ²
	<i>Senecio nemorensis</i> agg.	25 ³	.	15 ³	13 ³	.	8 ³	74 ³	99 ⁶	99 ⁷	99 ⁷	40 ²	60 ²	40 ⁴	50 ³
	<i>Soldanella carpatica</i>	57 ²	73 ²	63 ²	53 ³	.	33 ³	56 ²	20 ²	50 ²	38 ²	.	.	.	
	<i>Avenella flexuosa</i>	21 ²	19 ²	20 ²	20 ²	11 ²	17 ²	19 ³	40 ³	20 ³
	<i>Anthoxanthum odoratum</i> agg.	23 ²	32 ²	27 ²	13 ²	.	8 ²	13 ²	.	13 ²	8 ²	.	.	20 ²	10 ²
2405	<i>Heracleum sphondylium</i>	9 ²	3 ³	6 ²	.	.	.	50 ²	40 ²	50 ²	46 ²	.	40 ²	80 ²	60 ²
	<i>Geum rivale</i>	7 ²	.	4 ²	.	.	.	56 ³	60 ³	75 ³	69 ³	.	40 ³	.	20 ³
	<i>Arabis alpina</i>	4 ³	.	2 ³	.	.	.	24 ³	.	25 ²	15 ²	.	80 ³	40 ⁶	60 ⁴
	<i>Chamerion angustifolium</i>	5 ²	.	3 ²	7 ²	11 ²	8 ²	4 ⁵	20 ²	.	10 ²
	<i>Streptopus amplexifolius</i>	4 ²	3 ¹	3 ²	7 ²	11 ²	8 ²	2 ²	
	<i>Sedum alpestre</i>	2 ²	11 ²	5 ²	7 ²	.	4 ²	2 ²	
2410	<i>Taraxacum</i> sp.	2 ¹	11 ²	5 ¹	.	.	.	11 ²	20 ²	.	10 ²
	<i>Caltha *laeta</i>	4 ⁴	5 ³	4 ³	.	.	.	15 ⁴	20 ²	.	
	<i>Melampyrum sylvaticum</i>	2 ¹	.	1 ¹	.	.	.	2 ²	20 ²	10 ²	
	<i>Leontodon hispidus</i>	4 ²	.	2 ²	.	.	.	4 ²	.	13 ²	8 ²	.	.	.	

	<i>Pedicularis hacquetii</i>	5 ¹	3 ¹	20 ²	13 ³	15 ³	.	.	.	20 ⁵	10 ⁵
2415	<i>Silene vulgaris</i>	2 ²	1 ²	2 ²	20 ¹	10 ¹
	<i>Soldanella hungarica</i>	4 ⁶	2 ⁶	2 ²	20 ²	20 ²
	<i>Archangelica officinalis</i>	4 ⁴	3 ⁷	3 ⁵	.	.	.	13 ⁴
	<i>Dryopteris carthusiana</i>	.	5 ²	2 ²	7 ²	.	.	4 ²	9 ²
	<i>Festuca supina</i>	.	3 ²	1 ²	.	11 ²	4 ²
2420	<i>Asplenium viride</i>	.	3 ²	1 ²	40 ²	20 ²
	<i>Pinus mugo</i>	11 ³	4 ³	2 ²	40 ⁴	.	15 ⁴
	<i>Sorbus aucuparia</i>	11 ²	4 ²	2 ²	20 ²	.	8 ²	.	.	40 ¹	20 ¹	
	<i>Angelica sylvestris</i>	2 ²	20 ²	50 ²	38 ²	.	40 ²	20 ²	30 ²	
2425	<i>Adoxa moschatellina</i>	11 ²	60 ³	13 ³	31 ³	.	.	20 ¹	10 ¹	
	<i>Polystichum lonchitis</i>	2 ²	.	25 ²	15 ²	.	.	20 ²	10 ²	
	<i>Valeriana tripteris</i>	9 ²	20 ²	.	8 ²	40 ³	.	20 ²	10 ²	
	<i>Salix silesiaca</i>	9 ²	20 ²	.	8 ²	.	20 ³	.	10 ³	
	<i>Lamium maculatum</i>	2 ²	.	13 ²	8 ²	.	40 ⁶	40 ³	40 ⁴	
	<i>Rubus saxatilis</i>	2 ³	.	38 ²	23 ²	.	.	20 ²	10 ²	
2430	<i>Cystopteris montana</i>	4 ²	40 ²	40 ³	40 ²	
	<i>Veronica chamaedrys</i>	4 ²	40 ²	20 ²	30 ²	
	<i>Poa trivialis</i>	2 ²	40 ²	.	20 ²	
	<i>Rosa pendulina</i>	2 ¹	20 ²	13 ²	15 ²	
	<i>Digitalis grandiflora</i>	25 ²	15 ²	.	40 ²	40 ⁴	40 ³	
2435	<i>Cimicifuga europaea</i>	25 ⁴	15 ⁴	.	.	20 ²	10 ²	
	<i>Ajuga reptans</i>	40 ⁴	.	20 ⁴	
	<i>Picea abies</i>	40 ²	.	20 ²	
	Mosses															
2440	<i>Brachythecium rutabulum</i>	4 ⁶	3 ³	3 ⁵	13 ²	.	8 ²	.	20 ⁵	.	8 ⁵	.	20 ⁸	.	10 ⁸	
	<i>Plagiothecium denticulatum</i>	7 ²	16 ²	11 ²	27 ³	11 ²	21 ³	6 ²	20 ¹	13 ²	15 ²
	<i>Rhytidadelphus squarrosus</i>	5 ²	3 ²	4 ²	7 ³	.	4 ³	4 ³	60 ²	25 ³	38 ²
	<i>Cirriphyllum piliferum</i>	2 ⁸	5 ³	3 ⁴	13 ¹	8 ¹

	<i>Drepanocladus uncinatus</i>	4 ²	5 ²	4 ²	.	11 ²	4 ²	.	20 ¹	.	8 ¹	.	.	.
	<i>Tortella tortuosa</i>	4 ²	.	2 ²	.	.	.	2 ²	.	25 ²	15 ²	.	.	.
2445	<i>Brachythecium reflexum</i>	18 ³	41 ⁵	27 ⁴	13 ²	11 ²	13 ²	4 ⁶
	<i>Brachythecium salebrosum</i>	9 ⁴	3 ²	6 ⁴	7 ²	.	4 ²	2 ⁶
	<i>Brachythecium starkei</i>	9 ⁴	22 ³	14 ³	.	.	.	13 ⁴
	<i>Mnium</i> sp.	4 ²	.	2 ²	7 ¹	.	4 ¹	11 ²
2450	<i>Polytrichum</i> sp.	2 ²	.	1 ²	7 ²	.	4 ²	11 ²
	<i>Plagiomnium affine</i>	2 ³	.	1 ³	.	.	.	4 ⁵	.	.	.	20 ²	.	10 ²
	<i>Plagiochila asplenoides</i>	2 ²	.	1 ²	.	.	.	20 ²	13 ²	15 ²
	<i>Rhodobryum roseum</i>	4 ²	.	2 ²	38 ²	23 ²
	<i>Brachythecium glareosum</i>	2 ⁵	.	1 ⁵	25 ³	15 ³
2455	<i>Brachythecium velutinum</i>	2 ²	.	1 ²	25 ²	15 ²
	<i>Conocephalum conicum</i>	2 ⁶	20 ³	25 ³	23 ²	.	.	.
	<i>Plagiomnium cuspidatum</i>	2 ²	40 ³	.	15 ³	.	.	.
	<i>Eurhynchium schleicheri</i>	40 ³	13 ²	23 ²
	<i>Rhytidadelphus triquetrus</i>	20 ³	25 ⁴	23 ⁴

D1* differential taxa against the association *Adenostylo-Athyrietum alpestris*

2460 D3*, D5* differential taxa against the other communities of the suballiance *Delphinienion elati*

Rarely occurring taxa with low constancy:

E.: *Acetosa scutata* 5² (1a), 3² (1); *Aconitum vulparia* 20⁶ (6b), 10⁶ (6); *Agrostis capillaris* 2² (3); *A. stolonifera* 6² (3); *Ajuga genevensis* 20² (6a), 10² (6); *Alliaria petiolata* 20³ (6b), 10³ (6); *Allium *sibiricum* 2² (3); *A. *montanum* 20² (6b), 10² (6); *Pc Anthriscus nitida* 2² (3); *Anthriscus* sp. 20² (6a), 10² (6); *Anthyllis *alpestris* 20³ (6b), 10³ (6); *Asplenium ruta-muraria* 20² (6b), 10² (6); *A. trichomanes* 20² (6b), 10² (6); *Avenula versicolor* 3² (1b), 1² (1); *Botrychium lunaria* 20¹ (6b), 10¹ (6); *Calamagrostis* sp. 4³ (3); *Callianthemum coriandrifolium* 2² (1a), 1² (1); *Cardamine amara* 2² (3), 20² (6a), 10² (6); *C. pratensis* 2² (1a), 1² (1); *Cardaminopsis arenosa* agg. 17² (3), 40⁴ (6b), 20⁴ (6); tf *C. neglecta* 2² (1a), 1² (1); tf *Carex aterrima* 4² (1a), 2² (1); *C. buxbaumii* 2² (3); Cv *C. *silicicola* 5² (1a), 3² (1); *C. *tatarorum* 20² (6b), 10² (6); *C. sylvatica* 20² (6a), 10² (6); *Carlina acaulis* 20² (6b), 10² (6); *Cerastium *glandulosum* 4² (3); *C. holosteoides* 2¹ (3); *Cerinthe glabra* 13⁵ (4b), 8⁵ (4); *Chaerophyllum aromaticum* 2³ (3), 20² (6a), 10² (6); *Chelidonium majus* 20² (6b), 10² (6); *Cirsium oleraceum* 4³ (3), 20³ (6a), 10³ (6); *C. palustre* 2¹ (3); *Clematis alpina* 20² (6b), 10² (6);

- 2475 *Coeloglossum viride* 2¹ (3); *Conioselinum tataricum* 2¹ (3); *Crepis jacquinii* 2² (3); *Dactylorhiza fuchsii* 40¹ (6a), 20¹ (6); *Delphinium oxysepalum* 11⁴ (3); *Delphinium* sp. 2³ (3); *Epilobium alsinifolium* 15² (3); *E. anagallidifolium* 2² (1a, 1), 3¹ (1b); *E. collinum* 2¹ (3); *Equisetum arvense* 20² (6a), 10² (6); *Erysimum wahlenbergii* 13² (4b), 8² (4); *Euphrasia salisburgensis* 2² (3); *E. tatrae* 2² (3); *Fagus sylvatica* 20¹ (6a), 10¹ (6); *Festuca gigantea* 20² (6a), 10² (6); *F. rubra* 2² (3); *F. tatrae* 20³ (6b), 10³ (6); *Festuca* sp. 2² (1a), 1² (1); *Galeopsis speciosa* 20² (6b), 10² (6); *G. tetrahit* 20² (6a), 10² (6); *Galeopsis* sp. 20¹ (6b), 10¹ (6); *Galium album* 4² (3), 40⁴ (6b), 20⁴ (6); *G. anisophyllum* 4² (3), 40³ (6b), 20³ (6); *Gentiana frigida* 3² (1b), 1² (1); *Glechoma hederacea* s. l. 20² (6a), 20³ (6b, 6); *Gymnadenia conopsea* 2² (3), 20¹ (6b), 10¹ (6); *Gymnocarpium dryopteris* 2² (3), 20² (6b), 10² (6); *G. robertianum* 20³ (6b), 10³ (6); cv *Hieracium alpinum* agg. 5² (1a), 3² (1); *H. murorum* 20² (6b), 10² (6); *Hieracium* sp. 2² (3); *Huperzia selago* 2¹ (1a), 1¹ (1), 2² (3); *Hylotelephium maximum* 20² (6b), 10² (6); *Hypericum hirsutum* 20² (6b), 10² (6); *Jovibarba globifera* 20³ (6b), 10³ (6); *Juniperus communis* 11² (2b), 4² (2); *Knautia arvensis* 20² (6b), 10² (6); *Lapsana communis* 20² (6b), 10² (6); *Lathyrus pratensis* 20² (6b), 10² (6); *Leucanthemum margaritae* 20² (6b), 10² (6); *Libanotis pyrenaica* 13² (4b), 8² (4); *Linum catharticum* 20¹ (6b), 10¹ (6); *Lonicera nigra* 20² (4a), 8² (4); *L. xylosteum* 20¹ (6b), 10¹ (6); *Lotus corniculatus* 20² (6b), 10² (6); *Maianthemum bifolium* 2¹ (3); *Omalotheca norvegica* 5¹ (1a), 3² (1b), 4¹ (1), 4² (3); *O. supina* 3² (1b), 1² (1); *O. sylvatica* 2¹ (3); *Oxyria digyna* 2² (1a, 1, 3), 3² (1b); *Parnassia palustris* 6² (3); *Phyteuma orbiculare* 13² (4b), 8² (4); *Poa laxa* 2² (1a), 1² (1); *P. pratensis* 7¹ (2a), 4¹ (2); *P. stiriaca* 20² (6a), 10² (6); *Polemonium caeruleum* 20² (6a), 10² (6); *Polygala *brachyptera* 20² (6b), 10² (6); *Polypodium vulgare* 20² (6b), 10² (6); *Polystichum aculeatum* 20³ (4a), 8³ (4); *Primula minima* 3² (1b), 1² (1); *Pseudorchis albida* 2¹ (1a), 1¹ (1); *Pulsatilla scherfelii* 3² (1b), 1² (1); *Ranunculus acris* 2³ (1a, 1), 3² (1b); *R. alpestris* 20² (6b), 10² (6); *R. breyninus* 4² (3), 20³ (6b), 10³ (6); *R. nemorosus* 11² (1b), 4² (1), 2¹ (3); *R. polyanthemos* 2² (3), 20² (6b), 10² (6); *Rhinanthus pulcher* 5² (1a), 2² (1), 13² (4b), 8² (4); *Rhinanthus* sp. 20¹ (6b), 10¹ (6); *Rubus caesius* 20² (6b), 10² (6); *Rumex alpinus* 9³ (3); *Salix caprea* 20¹ (6b), 10¹ (6); *Salvia glutinosa* 20² (6a), 10² (6); *Sanicula europaea* 20² (6a), 10² (6); *Saxifraga androsacea* 2³ (3); *S. carpatica* 3² (1b), 1² (1); *S. hieraciifolia* 2¹ (3); *S. paniculata* 4² (3), 20³ (6b), 10³ (6); *S. wahlenbergii* 4² (3); *Scabiosa lucida* 2² (3), 20² (6b), 10² (6); *Scrophularia nodosa* 20² (6a), 10² (6); *Sesleria albicans* 20² (6b), 10² (6); *S. tatrae* 9² (3), 13³ (4b), 8³ (4); *Silene pusilla* 4² (3); *Stachys alpina* 20⁵ (6b), 10⁵ (6); *Swertia *alpestris* 3² (1b), 1² (1), 6³ (3); *Sympyton tuberosum* 2² (3); *Thesium alpinum* 20² (6a), 10² (6); *Thlaspi *tatrense* 20¹ (6b), 10¹ (6); *Thymus pulcherrimus* 20³ (6b), 10³ (6); *Tragopogon orientalis* 20¹ (6b), 10¹ (6); *Trifolium badium* 2² (3); *T. orbelicum* 2² (3); *Trollius altissimus* 7³ (3); Cv *Trommsdorffia uniflora* 2¹ (1a), 1¹ (1); *Tussilago farfara* 4² (3), 20² (6a), 10² (6); *Vaccinium vitis-idaea* 2³ (3); *Veronica alpina* 5¹ (1b), 2¹ (1), 4² (3); *Veronica* sp. 2¹ (3), 20² (6b), 10² (6); *Vicia cracca* 20² (6b), 10² (6); *V. sepium* 7² (3), 40² (6b), 20² (6); *Viola tricolor* 20¹ (6b), 10¹ (6).
 E₀: *Amblystegium serpens* 2³ (1a), 1³ (1); *Anomodon viticulosus* 2³ (1a), 1³ (1); *Barbilophozia lycopodioides* 13² (2b), 8² (2); *Bartramia ithyphylla* 3³ (1b), 1³ (1); *Brachythecium rivulare* 2³ (3); *Brachythecium* sp. 3³ (1b), 1³ (1), 13⁶ (3); *Bryum capillare* 13¹ (4b),

8¹ (4); *B. pallescens* 2³ (1a), 1³ (1); *B. weigelii* 2³ (1a), 1³ (1); *Bryum* sp. 4³ (1a, 1), 5³ (1b), 4² (3); *Calypogeia azurea* 22² (2b), 8² (2); *Campylium stellatum* 13⁴ (2a), 8⁴ (2); *Ceratodon purpureus* 2² (1a), 1² (1); *Chiloscyphus polyanthos* 2² (1a), 1² (1); *Cirriphyllum cirrosum* 3¹ (1b), 1¹ (1), 2² (3); *Cratoneuron* sp. 2³ (3); *Dicranella heteromalla* 3³ (1b), 1³ (1); *Dicranum scoparium* 2² (3), 20³ (4a), 8³ (4); *Dicranum* sp. 2² (3); *Distichium capillaceum* 2² (3); *Eurhynchium striatum* 13³ (4b), 8³ (4); *Eurhynchium* sp. 2³ (3); *Fissidens dubius* 13² (4b), 8² (4); *F. taxifolius* 2² (1a), 1² (1); *Fissidens* sp. 20² (4a), 8² (4); *Grimmia* sp. 2² (3); *Hylocomium splendens* 7² (2a), 4² (2), 20² (4a), 8² (4); *Hypnum cupressiforme* 33⁴ (2b), 13⁴ (2); *Hypnum* sp. 2⁷ (3); *Jungermannia obovata* 2³ (3); *Kiarea starkei* 7² (2a), 4² (2); *Kiarea* sp. 3² (1b), 1² (1); *Lescuraea atrovirens* 5¹ (1a), 11² (1b), 8² (1); *L. mutabilis* 2² (1a), 5³ (1b), 3³ (1); *Leskea polycarpa* 11⁵ (1a), 11³ (1b), 11⁴ (1); *Marchantia polymorpha* 2³ (3); *Marsupella sphacelata* 3² (1b), 1² (1); *Mnium marginatum* 13⁵ (4b), 8⁵ (4); *M. spinulosum* 2² (3), 20² (4a), 8² (4); *Oligotrichum hercynicum* 3² (1b), 1² (1); *Palustriella decipiens* 2⁵ (3); *Pellia neesiana* 5³ (1b), 2³ (1); *Pellia* sp. 2² (1a), 1² (1); *Philonotis fontana* 3² (1b), 1² (1); *Ph. seriata* 2¹ (1a), 1¹ (1); *Philonotis* sp. 6² (3); *Plagiomnium rostratum* 3¹ (1b), 1¹ (1), 6² (3); *P. undulatum* 2² (3); *Plagiomnium* sp. 2³ (1a), 1³ (1); *Plagiothecium cavifolium* 2⁶ (1a), 1⁶ (1); *P. curvifolium* 2⁶ (1a), 1⁶ (1), 11² (2b), 4² (2); *P. laetum* 25² (4b), 15² (4); *P. nemorale* 2² (1a), 3³ (1b), 2³ (1); *P. platyphyllum* 2³ (1a, 1), 3³ (1b); *Plagiothecium* sp. 5³ (1a), 2³ (1); *Pleurozium schreberi* 3² (1b), 1² (1), 11² (2b), 4² (2); *Pohlia drummondii* 8⁶ (1b), 3⁶ (1); *P. ludwigii* 8³ (1b), 3³ (1), 4² (3); *P. nutans* 2³ (1a), 5³ (1b), 3³ (1); *P. wahlenbergii* 3⁵ (1b), 1⁵ (1); *Pohlia* sp. 5² (1b), 2² (1); *Polytrichum alpinum* 5² (1a), 8² (1b), 6² (1); *P. formosum* 7³ (1a), 4³ (1), 4⁵ (3); *P. juniperinum* 7³ (2a), 4³ (2); *P. piliferum* 3² (1b), 1² (1); *Pseudoleskeia incurvata* 2³ (1a), 14⁸ (1b), 6⁷ (1); *Pseudoleskeella nervosa* 3¹ (1b), 1¹ (1); *Racomitrium heterostichum* 5³ (1b), 2³ (1); *Rhizomnium magnifolium* 2⁵ (3); *Rh. punctatum* 2⁵ (1a), 1⁵ (1), 20³ (2a), 13³ (2); *Rhynchostegium murale* 13³ (4b), 8³ (4); *Rhytidadelphus subpinnatus* 2¹ (1a), 3⁵ (1b), 2³ (1); *Scapania uliginosa* 3¹ (1b), 1¹ (1); *Schistidium strictum* (Turn.) Loeske ex Mart. 7¹ (2a), 4¹ (1); *Splachnum sphaericum* 3¹ (1b), 1¹ (1); *Tritomaria exsecta* 3² (1b), 1² (1). – *Cetraria islandica* 3⁵ (1b), 1⁵ (1); *Cladonia coccifera* 3³ (1b), 1³ (1); *C. mitis* 3² (1b), 1² (1).

Tab. 5. Geranio robertianii-Delphinietum elati ass. nov.

subass. *orobanchetosum flavae* subass. nov. (r. 1–5), subass. *ranunculetosum platanifolii* subass. nov. (r. 6–10)

2525	Number of relevé		1	2	3	4	5	6	7	8	9	10	St		
			32	24	40	40	38	46	46	39	45	57	%		
Diagnostic taxa of the association															
	de	<i>Delphinium elatum</i>	t		2a	2b	4	4	4	2b	3	4	2b	3	100
		<i>Urtica dioica</i>	D	+	+	1	+	1	+	1	2m+	+	1	100	
2530	Fs	<i>Geranium robertianum</i>	D	+	1	+	+	+	.	+	2a	+	.	80	
	EA	<i>Fragaria vesca</i>	D	+	+	.	.	+	+	.	.	.	+	50	
	Fs	<i>Acer pseudoplatanus</i>	D	+	.	+	.	.	r	+	r	.	.	50	
		<i>Carex muricata</i>	D	+	+	+	.	.	30		
		<i>Clinopodium vulgare</i>	D	.	.	+	.	.	+	.	.	+	30		
2535	Differential taxa of the subassociations														
	Pc	<i>Stellaria nemorum</i>		1	1	2b	2a	2b	50	
	Pc	<i>Petasites kablodianus</i>		.	2a	2b	3	3	40	
	fs	<i>Petasites albus</i>		2a	2a	.	+	1	40	
	ai,cl	<i>Crepis paludosa</i>		+	+	+	2a	40	
2540	cl	<i>Myosotis nemorosa</i>		+	.	+	+	+	30	
	Fs	<i>Stachys sylvatica</i>		.	+	2a	+	30	
	Pc	<i>Orobanche flava</i>		.	.	+	+	+	30	
	ai	<i>Roegneria canina</i>		.	.	2b	1	+	30	
	Fs	<i>Impatiens noli-tangere</i>		.	.	+	+	1	30	
2545	aa	<i>Ranunculus platanifolius</i>		+	+	+	+	+	.	40	
	fs	<i>Polygonatum verticillatum</i>		+	+	.	1	1	1	40	
	Cv	<i>Luzula luzuloides</i>		1	+	.	.	+	30	
	MU	<i>Geranium sylvaticum</i>		+	.	.	+	.	2b	30	
	QF	<i>Galium schultesii</i>		+	+	1	.	.	30		
2550	Fs	<i>Galeobdolon montanum</i>		+	+	+	.	30		
	Cv	<i>Cirsium erisithales</i>		+	.	.	+	20		
Differential taxa of the suballiance Delphinienion elati															
	Pc	<i>Valeriana *sambucifolia</i>		+	+	1	1	2a	.	+	1	.	.	70	
		<i>Heracleum sphondylium</i>		.	.	+	+	.	.	+	+	+	1	60	
2555	Pc	<i>Carduus personata</i>		.	.	1	+	1	+	.	+	.	+	60	
	Fs,EA	<i>Senecio nemorensis</i> agg.		.	.	+	+	+	+	2a	.	.	.	50	
		<i>Cortusa matthioli</i>		+	.	.	+	20	
	cl	<i>Geum rivale</i>		.	.	.	+	1	20	
	ac	<i>Luzula sylvatica</i>		+	10	
2560		<i>Epilobium alpestre</i>		1	10		
Adenostylium alliaiae, Mulgedio-Aconitetea															
	Pc	<i>Chaerophyllum hirsutum</i>		4	2a	2b	2a	3	+	4	.	.	.	70	
	Pc,ai	<i>Chrysosplenium alternifolium</i>		.	+	+	1	1	+	2b	.	.	.	60	
	aa	<i>Silene dioica</i>		.	.	+	+	.	+	+	1	.	+	60	
		<i>Aconitum variegatum</i>		.	.	+	+	1	.	+	.	1	.	50	
2565	MU,cr	<i>Calamagrostis varia</i>		2b	.	.	.	4	.	+	.	2a	40		
		<i>Gentiana asclepiadea</i>		+	1	+	.	.	+	40	

		<i>Thalictrum aquilegiifolium</i>	.	.	.	r	.	.	1	.	+	+	40
		<i>Knautia maxima</i>	.	.	.	1	2a	20
2570	aa,Fs	<i>Milium effusum</i>	+	1	.	.	.	20
	Pc	<i>Geranium phaeum</i>	+	.	1	.	.	20
		<i>Primula elatior</i>	+	.	.	+	.	20
	Cv	<i>Achillea *alpestris</i>	+	.	+	.	20
	Cv	<i>Campanula serratula</i>	1	+	20	
2575	Cv	<i>Pleurospermum austriacum</i>	+	10
	Pc	<i>Petasites hybridus</i>	.	.	2b	10
		<i>Acetosa arifolia</i>	2a	10
	Cv	<i>Crepis conyzifolia</i>	2a	10
	Cv	<i>Pyrethrum clusii</i>	+	10
2580	Cv	<i>Vicia oreophila</i>	+	10
	Cv	<i>Laserpitium latifolium</i>	+	.	.	.	10
	Cv	<i>Bupleurum *vapincense</i>	1	10		
	Cv	<i>Campanula elliptica</i>	+	10		
	Cv	<i>Crepis mollis</i>	+	10		
2585	Cv	<i>Cyanus mollis</i>	+	10		
	fc	<i>Festuca carpatica</i>	+	10		
	Cv	<i>Hieracium prenanthoides</i>	+	10		
	Cv	<i>Linum extraaxillare</i>	+	10		
	Cv	<i>Phleum hirsutum</i>	+	10		
2590	Cv	<i>Pimpinella *rhodochlamys</i>	+	10		
		<i>Poa chaixii</i>	+	10		
		Querco-Fagetea											
	Fs	<i>Dryopteris filix-mas</i>	.	.	+	+	+	2a	.	+	.	.	50
	QF	<i>Poa nemoralis</i>	1	2b	2b	2a	.	40
2595	Fs	<i>Myosotis sylvatica</i>	.	+	1	.	+	+	40
	ai	<i>Oxalis acetosella</i>	.	+	.	+	2m	+	40
	ai	<i>Lamium maculatum</i>	.	.	2b	2a	.	.	1	.	+	40	
	Fs,EA	<i>Epilobium montanum</i>	.	.	.	+	+	+	+	.	.	.	40
	Fs	<i>Cardamine impatiens</i>	+	.	+	.	.	.	1	.	.	30	
2600	Fs	<i>Lathyrus vernus</i>	+	1	1	.	30	
	Fs	<i>Pulmonaria obscura</i>	.	.	1	.	.	+	.	1	.	.	30
	ai	<i>Angelica sylvestris</i>	.	.	+	+	.	.	.	+	.	.	30
	Fs	<i>Daphne mezereum</i>	.	.	.	r	.	r	.	.	+	.	30
	fs	<i>Hordelymus europaeus</i>	1	+	20
2605	Fs	<i>Campanula trachelium</i>	.	.	.	+	+	20
	ac	<i>Scrophularia scopolii</i>	+	.	.	.	1	20	
	Fs	<i>Mercurialis perennis</i>	+	1	20
	Fs	<i>Melica nutans</i>	+	1	.	.	.	20
	Fs	<i>Tithymalus amygdalooides</i>	1	+	20	
2610		Other taxa											
		<i>Viola biflora</i>	+	+	+	+	+	1	.	.	2m.	70	
		<i>Arabis alpina</i>	+	+	.	1	2a	.	.	.	2b	2a	60
		<i>Hypericum maculatum</i>	+	.	.	.	+	1	.	.	2a	+	50
		<i>Cystopteris montana</i>	+	+	1	.	.	+	40
2615		<i>Digitalis grandiflora</i>	+	.	.	.	r	.	.	+	.	2a	40
		<i>Rubus idaeus</i>	.	.	+	1	+	1	40
		<i>Veronica chamaedrys</i>	+	.	.	.	+	.	+	.	.	.	30

	<i>Dactylis glomerata</i>	.	.	.	r	.	+	.	.	.	2a	30
	<i>Dactylorhiza fuchsii</i>	r	r	20	
2620	<i>Picea abies</i>	+	.	.	.	r	20	
	<i>Ajuga reptans</i>	.	.	.	2b	+	20	
	<i>Poa trivialis</i>	.	.	+	.	+	20	
	<i>Sorbus aucuparia</i>	r	r	.	.	20	
2625	<i>Asplenium viride</i>	+	.	.	+	20	
	<i>Cardaminopsis arenosa</i> agg.	+	2b	.	.	20	
	<i>Galium anisophyllum</i>	+	.	1	.	20	
	<i>Origanum vulgare</i>	1	.	+	20	
	<i>Galium album</i>	+	.	2a	20	
2630	<i>Vicia sepium</i>	+	.	+	20	
	<i>Avenella flexuosa</i>	+	+	20	
	<i>Thesium alpinum</i>	+	r	20	

Taxa occurring in 1–2 relevés:

E₁: *Aconitum vulparia* 2b (9); *Adoxa moschatellina* r (6); *Aegopodium podagraria* + (3), 1

2635 (8); *Ajuga genevensis* + (1); *Alchemilla* sp. + (10); *Alliaria petiolata* + (8); *Allium *montanum* + (10); *Anthoxanthum alpinum* + (10); *Anthriscus* sp. + (1); *Anthyllis *alpestris* 1 (9); *Aruncus vulgaris* r (4), + (7); *Asarum europaeum* + (10); *Asplenium ruta-muraria* + (9); *A. trichomanes* + (9); *Botrychium lunaria* r (7); *Cardamine amara* + (5); *Cardaminopsis halleri* + (6); *Carex *tatarorum* + (9); *C. sylvatica* + (3); *Carlina acaulis* + (9); *Chaerophyllum aromaticum* + (4); *Chamerion angustifolium* + (3); *Chelidonium majus* + (8); *Cimicifuga europaea* + (7); *Cirsium oleraceum* 1 (3); *Clematis alpina* + (6); *Deschampsia cespitosa* + (5); *Equisetum arvense* + (5); *Fagus sylvatica* r (2); *Festuca gigantea* + (5); *Festuca tatrae* 1 (9); *Galeopsis speciosa* + (10); *G. tetrahit* + (5); *Galeopsis* sp. r (6); *Glechoma hederacea* + (4), 1 (7); *Gymnadenia conopsea* r (9); *Gymnocarpium dryopteris* + (6); *G. robertianum* 1 (9); *Hieracium murorum* + (9); *Homogyne alpina* + (10); *Hylotelephium argutum* 1 (6); *H. maximum* + (8); *Hypericum hirsutum* + (7); *Jovibarba globifera* 1 (9); *Knautia arvensis* + (10); *Lapsana communis* + (8); *Lathyrus pratensis* + (10); *Leucanthemum margaritae* + (10); *Lilium martagon* r (10); *Linum catharticum* + (9); *Lonicera xylosteum* r (6); *Lotus corniculatus* + (9); *Lunaria rediviva* + (2), 1 (8); *Melampyrum sylvaticum* + (6); *Mycelis muralis* + (4, 9); *Paris quadrifolia* + (2); *Phyteuma spicatum* r (6); *Poa alpina* + (9); *P. stiriaca* + (2); *Polemonium caeruleum* + (4); *Polygala *brachyptera* + (9); *Polypodium vulgare* + (6); *Polystichum lonchitis* + (9); *Prenanthes purpurea* + (7); *Ranunculus alpestris* + (9); *R. breyninus* 1 (7); *R. lanuginosus* + (2, 8); *R. polyanthemos* + (9); *Rhinanthus* sp. r (7); *Rubus caesius* + (8); *R. saxatilis* + (7); *Salix caprea* r (7); *S. silesiaca* 1 (1); *Salvia glutinosa* + (3); *Sanicula europaea* + (4); *Saxifraga paniculata* 1 (9); *Scabiosa lucida* + (9); *Scrophularia nodosa* + (3); *Sesleria albicans* + (9); *Silene vulgaris* 2a (10); *Soldanella hungarica* r (6); *Stachys alpina* 2a (10); *Taraxacum* sp. + (2); *Thlaspi *tatrense* r (6); *Thymus pulcherrimus* 1 (9); *Tragopogon orientalis* r (10); *Trisetum flavescens* 1 (10); *Tussilago farfara* + (1); *Valeriana tripteris* + (6); *Veronica* sp. + (8); *Vicia cracca* + (10); *Viola tricolor* r (6).

E₀: *Brachythecium rutabulum* 4, *Plagiognathus affine* +.

**Tab. 6. Comparison of the floristic composition of the higher syntaxa of the class
Mulgedio-Aconitetea.**

2665

(abbreviated synoptic table)

Number of column	1	2	3	4	5	6a	6b	7
Number of relevés	210	147	148	63	97	117	81	353
Average number of taxa	24	26	41	43	40	21	34	29
2670	Calamagrostion villosae							
	<i>Calamagrostis villosa</i>	C1	99⁸	39 ³	34 ⁸	.	12 ⁴	72 ³
	<i>Carex *silicicola</i>	t1	47³	24 ²	.	.	3 ²	.
	<i>Hieracium alpinum</i> agg.	C1	33²	5 ²	.	.	5 ²	3 ²
	<i>Avenula versicolor</i>	C1	28²	7 ²	.	2 ²	3 ²	1 ²
2675	<i>Sempervivum *carpathicum</i>	C1	22²	4 ²	.	.	1 ²	3 ²
	<i>Pulsatilla scherfelii</i>	D1	30³	8 ²	.	.	1 ²	.
	Trisetion fuscī							
	<i>Trisetum fuscum</i>	C2	3 ²	31⁴	.	.	11 ³	11 ²
	<i>Rhodiola rosea</i>	C2	13 ²	60⁴	1 ²	2 ²	31 ³	23 ²
2680	<i>Taraxacum alpinum</i>	C2	2 ²	41²	.	.	8 ²	.
	<i>Carex aterrima</i>	C2	3 ²	27²	.	.	2 ²	2 ²
	<i>Cerastium fontanum</i>	C2	1 ²	21²	.	2 ²	5 ²	.
	<i>Cardaminopsis neglecta</i>	C2	.	14²	.	.	1 ²	.
2685	<i>Caltha *laeta</i>	D2	+ ²	44³	.	.	3 ³	11 ⁴
	<i>Cardamine amara</i> s. l.	D2	.	13²	.	.	.	17 ²
	<i>Bryum pseudotriquetrum</i> (E ₀)	C2	.	20²	.	.	.	1 ³
	Calamagrostion arundinaceae							
2690	<i>Calamagrostis arundinacea</i>	C3	4 ²	1 ⁵	87⁶	16 ³	22 ²	3 ²
	<i>Knautia maxima</i>	t3	+ ²	.	59³	24 ²	12 ³	1 ¹
	<i>Vicia oreophila</i>	t3	.	.	41²	24 ²	5 ²	2 ²
	<i>Hieracium prenanthoides</i>	C3	2 ²	.	39³	14 ³	6 ²	.
2695	<i>Allium victorialis</i>	C3	.	.	32³	8 ¹	1 ²	.
	<i>Dianthus *latifolius</i>	C3	.	.	30²	17 ²	3 ²	.
	<i>Vicia sylvatica</i>	C3	.	.	27⁴	13 ³	4 ²	.
	<i>Jacea pseudophrygia</i>	C3	.	.	22⁴	11 ²	1 ³	.
2700	<i>Agrostis capillaris</i>	D3	.	7 ²	55⁴	11 ³	3 ²	.
	<i>Cruciata glabra</i>	D3	.	1 ²	47²	13 ²	3 ³	.
	<i>Briza media</i>	D3	.	1 ¹	32²	17 ²	1 ²	.
	<i>Avenula planiculmis</i>	D3	.	.	24⁴	2 ²	1 ²	.
	Calamagrostion variae							
	<i>Calamagrostis varia</i>	C4	.	.	2 ²	99⁸	23 ⁴	.
	<i>Knautia kitaibelii</i>	C4	.	1 ²	2 ²	33²	6 ³	.
2705	<i>Epipactis atrorubens</i>	C4	.	.	.	21¹	.	.
	<i>Gymnadenia odoratissima</i>	C4	.	.	.	14¹	1 ²	.
	<i>Carduus glaucinus</i>	D4	.	.	1 ²	51³	14 ²	.
	<i>Mercurialis perennis</i>	D4	.	.	13 ²	51²	8 ³	.
2710	<i>Acinos alpinus</i>	D4	.	.	.	17²	.	.
	Festucion carpaticaæ							
	<i>Festuca carpatica</i>	C5	.	.	7 ³	10 ³	99⁸	1 ²
	<i>Bartsia alpina</i>	C5	7 ²	3 ²	1 ²	.	33²	26 ³

125

	<i>Sesleria tatrae</i>	D5	2 ²	.	7 ³	6 ³	56³	.	7 ²	.
	<i>Luzula sylvatica</i>	D5,D6b	7 ³	3 ²	13 ²	8 ²	48³	7 ³	51³	5 ²
	<i>Cortusa matthioli</i>	D5,D6b	.	.	1 ²	13 ³	42³	.	25³	9 ²
2715	<i>Calamagrostietalia villosae</i>									
	<i>Luzula luzuloides</i>	t	59³	12²	86³	6³	51³	6 ²	21 ³	3 ²
	<i>Solidago virgaurea</i>		59²	18²	43²	22²	12³	26 ²	9 ²	1 ²
	<i>Campanula serrata</i>		9 ²	2 ²	71 ²	24 ²	28 ²	4 ¹	16 ²	1 ²
2720	<i>Achillea *alpestris</i>		8 ²	7 ²	67 ³	22 ³	46 ³	1 ²	11 ²	.
	<i>Anemone narcissiflora</i>		3 ²	3 ³	43 ³	13 ³	31 ³	3 ²	1 ³	.
	<i>Crepis conyzifolia</i>		25 ³	7 ²	36 ³	6 ²	7 ²	4 ²	2 ⁴	.
	<i>Festuca picturata</i>		70 ⁴	63³	.	.	15 ³	62 ³	2 ²	0 ²
2725	<i>Gentiana punctata</i>	D6a	57 ³	30²	.	.	1 ²	70³	2 ²	.
	<i>Trommsdorffia uniflora</i>		27 ²	1 ²	11 ²	.	1 ³	1 ¹	.	.
	<i>Campanula tatrae</i>		37 ²	20²	1 ²	5 ²	15²	.	1 ²	.
	<i>Crepis mollis</i>		+ ²	.	48 ²	19 ²	48 ³	.	12 ²	1 ³
	<i>Linum extraaxillare</i>		+ ⁴	.	39 ³	17 ³	38 ³	.	5 ²	.
2730	<i>Cirsium erisithales</i>		.	.	53 ³	70 ²	34 ³	.	20 ³	3 ²
	<i>Cyanus mollis</i>		.	.	27 ⁴	37 ²	18 ³	.	9 ³	0 ¹
	<i>Laserpitium latifolium</i>		.	.	33 ⁴	86 ⁴	14 ²	.	4 ²	0 ²
	<i>Pimpinella major</i>		.	.	60 ³	65 ³	52 ⁴	.	19 ³	1 ²
	<i>Pleurospermum austriacum</i>		.	.	11 ²	16 ²	4 ³	.	6 ²	0 ²
2735	<i>Bupleurum longifolium</i>		.	.	14 ³	6 ³	15 ³	.	2 ³	.
	<i>Campanula elliptica</i>		.	.	50 ³	27 ²	32 ²	.	5 ²	.
	<i>Phleum hirsutum</i>		.	.	49 ³	24 ³	39 ³	.	2 ³	.
	<i>Pyrethrum clusii</i>		.	.	66 ³	46 ²	13 ²	.	1 ²	.
	<i>Adenostylium alliariae</i>									
	<i>Adenostyles alliariae</i>	C6a,b	25 ²	25 ³	9 ³	3 ²	22 ³	94⁷	79⁷	3 ²
2740	<i>Ranunculus platanifolius</i>	t6a,b	20 ²	10 ³	19 ²	6 ³	7 ³	53³	23³	3 ²
	<i>Doronicum austriacum</i>	C6a,b	10 ²	14 ²	2 ²	.	3 ²	53⁴	49³	6 ³
	<i>Milium effusum</i>	t6a,b	9 ²	12 ³	3 ²	.	10 ²	51³	42⁴	15 ²
	<i>Silene dioica</i>	t6a,b	7 ²	9 ³	5 ²	.	26 ²	41³	52²	32 ²
2745	<i>Athyrium distentifolium</i>	C6a,b	9 ²	14 ²	1 ²	2 ²	.	44⁶	25³	4 ²
	<i>Cicerbita alpina</i>	C6a,b	1 ³	2 ³	3 ²	.	2 ²	37⁴	37⁴	2 ²
	<i>Adenostylenion alliariae</i>									
	<i>Luzula alpinopilosa</i>	D6a	29 ³	51²	.	.	.	61³	7 ³	.
	<i>Oreogenum montanum</i>	D6a	80 ³	68 ³	.	.	18 ²	73³	6 ²	1 ²
	<i>Delphinienion elati</i>									
2750	<i>Delphinium elatum</i>	C6b	.	.	6 ³	8 ³	7 ³	.	30⁵	5 ³
	<i>Epilobium alpestre</i>	D6b	.	1 ³	6 ²	2 ²	23 ²	2 ²	46³	5 ²
	<i>Petasition officinalis</i>									
	<i>Chaerophyllum hirsutum</i>	D6b,C7	1 ³	31 ⁴	6 ³	8 ³	6 ³	6 ²	78⁵	87⁴
	<i>Stellaria nemorum</i>	D6b,C7	.	29 ⁴	5 ²	.	5 ²	9 ²	53⁴	83⁴
2755	<i>Petasites hybridus</i>	C7	.	.	.	2 ³	.	.	14 ⁵	70⁸
	<i>Petasites kablikianus</i>	C7	1 ³	11 ⁶	32⁸
	<i>Chrysosplenium alternifolium</i>	C7	+ ²	20 ³	1 ²	.	2 ²	2 ³	38 ³	47⁴
	<i>Carduus personata</i>	C7	.	2 ³	8 ²	2 ²	8 ³	1 ²	37 ²	51³
	<i>Geranium phaeum</i>	C7	.	.	1 ²	2 ²	1 ³	.	4 ²	51³
2760	<i>Orobanche flava</i>	C7	4 ²	23²
	<i>Anthriscus nitida</i>	C7	1 ³	.	1 ²	16³

	<i>Impatiens noli-tangere</i>	D7	4 ²	54³
	<i>Mentha longifolia</i>	D7	47³
	<i>Chaerophyllum aromaticum</i>	D7	.	.	1 ²	2 ¹	6 ³	.	2 ³	46³
2765	<i>Poa trivialis</i>	D7	.	.	1 ²	1 ³	.	.	4 ²	46³
	<i>Ranunculus repens</i>	D7	.	1 ²	1 ³	.	.	.	1 ⁵	45²
	<i>Aegopodium podagraria</i>	D7	.	.	5 ²	2 ²	.	.	4 ²	39³
	<i>Lamium maculatum</i>	D7	.	.	1 ²	2 ²	1 ³	.	6 ⁴	37³
	<i>Filipendula ulmaria</i>	D7	.	1 ²	2 ³	2 ²	3 ³	.	11 ³	35²
2770	<i>Mulgedio-Aconitetea</i>									
	<i>Acetosa arifolia</i>		29 ³	30 ³	40 ³	2 ³	37 ²	83 ⁴	70 ⁴	18 ²
	<i>Aconitum firmum</i>	D2	14 ²	76⁴	2 ²	6 ³	32 ³	40 ³	47 ⁴	11 ³
	<i>Gentiana asclepiadea</i>		23 ²	7 ³	38 ²	19 ²	29 ²	12 ³	42 ²	6 ²
	<i>Geranium sylvaticum</i>		20 ²	36 ⁴	61 ³	25 ²	84 ⁴	30 ³	77 ³	15 ²
2775	<i>Poa chaixii</i>		22 ²	5 ²	16 ²	3 ²	25 ²	11 ³	25 ³	3 ²
	<i>Primula elatior</i>		2 ²	10 ²	25 ²	21 ²	70 ³	12 ³	57 ³	38 ²
	<i>Senecio subalpinus</i>		7 ³	31 ³	22 ²	3 ²	47 ³	15 ³	47 ²	5 ²
	<i>Thalictrum aquilegiifolium</i>		3 ²	8 ³	3 ²	8 ²	21 ³	7 ²	40 ²	19 ²
	<i>Valeriana *sambucifolia</i>		1 ²	1 ³	4 ²	8 ²	16 ³	1 ²	40 ³	29 ²
2780	<i>Veratrum *lobelianum</i>		61 ³	31 ²	12 ²	3 ²	31 ³	86 ³	48 ³	3 ²
	<i>Bistorta major</i>		40 ³	55 ³	4 ³	.	52 ³	42 ³	38 ³	+ ²
	<i>Astrantia major</i>		.	.	28 ²	33 ²	32 ²	1 ³	21 ²	11 ²
	<i>Aconitum variegatum</i>		.	.	9 ³	19 ³	4 ⁴	.	14 ³	14 ²

+ occurrence with frequency lower than 0,5%

2785